

Convex polygons in Cartesian products

Citation for published version (APA):

De Carufel, J. L., Dumitrescu, A., Meulemans, W., Ophelders, T., Pennarun, C., Tóth, C. D., & Verdonschot, S. (2019). Convex polygons in Cartesian products. In G. Barequet, & Y. Wang (Eds.), *35th International Symposium on Computational Geometry, SoCG 2019* [22] (Leibniz International Proceedings in Informatics (LIPIcs); Vol. 129). Schloss Dagstuhl - Leibniz-Zentrum für Informatik.
<https://doi.org/10.4230/LIPIcs.SocG.2019.22>

DOI:

[10.4230/LIPIcs.SocG.2019.22](https://doi.org/10.4230/LIPIcs.SocG.2019.22)

Document status and date:

Published: 01/06/2019

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Convex Polygons in Cartesian Products

Jean-Lou De Carufel

School of Electrical Engineering and Computer Science, University of Ottawa, Canada
jdecaruf@uottawa.ca

Adrian Dumitrescu

Department of Computer Science, University of Wisconsin-Milwaukee, USA
dumitres@uwm.edu

Wouter Meulemans

Department of Mathematics and Computer Science, TU Eindhoven, The Netherlands
w.meulemans@tue.nl

Tim Ophelders

Department of Computational Mathematics, Science and Engineering,
Michigan State University, East Lansing, MI, USA
ophelder@egr.msu.edu

Claire Pennarun

LIRMM, CNRS & Université de Montpellier, France
claire.pennarun@gmail.com

Csaba D. Tóth

Department of Mathematics, California State University Northridge, Los Angeles, CA, USA
Department of Computer Science, Tufts University, Medford, MA, USA
csaba.toth@csun.edu

Sander Verdonschot

School of Computer Science, Carleton University, Ottawa, ON, Canada
sander@cg.scs.carleton.ca

Abstract

We study several problems concerning convex polygons whose vertices lie in a Cartesian product of two sets of n real numbers (for short, *grid*). First, we prove that every such grid contains a convex polygon with $\Omega(\log n)$ vertices and that this bound is tight up to a constant factor. We generalize this result to d dimensions (for a fixed $d \in \mathbb{N}$), and obtain a tight lower bound of $\Omega(\log^{d-1} n)$ for the maximum number of points in convex position in a d -dimensional grid. Second, we present polynomial-time algorithms for computing the longest convex polygonal chain in a grid that contains no two points with the same x - or y -coordinate. We show that the maximum size of such a convex polygon can be efficiently approximated up to a factor of 2. Finally, we present exponential bounds on the maximum number of convex polygons in these grids, and for some restricted variants. These bounds are tight up to polynomial factors.

2012 ACM Subject Classification Theory of computation \rightarrow Computational geometry

Keywords and phrases Erdős–Szekeres theorem, Cartesian product, convexity, polyhedron, recursive construction, approximation algorithm

Digital Object Identifier 10.4230/LIPIcs.SoCG.2019.22

Related Version A full version of this paper is available at <https://arxiv.org/abs/1812.11332>.

Funding *Wouter Meulemans*: Partially supported by the Netherlands eScience Center (NLeSC, 027.015.G02)

Csaba D. Tóth: Supported in part by the NSF awards CCF-1422311 and CCF-1423615.

Acknowledgements This work was initiated at the 2017 Fields Workshop on Discrete and Computational Geometry (Carleton University, Ottawa, ON, July 31–August 4, 2017).



© Jean-Lou De Carufel, Adrian Dumitrescu, Wouter Meulemans, Tim Ophelders,
Claire Pennarun, Csaba D. Tóth, and Sander Verdonschot;
licensed under Creative Commons License CC-BY


35th International Symposium on Computational Geometry (SoCG 2019).

Editors: Gill Barequet and Yusu Wang; Article No. 22; pp. 22:1–22:17



Leibniz International Proceedings in Informatics

Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



1 Introduction

Can a convex polygon P in the plane be reconstructed from the projections of its vertices to the coordinate axes? Assuming that no two vertices of P share the same x - or y -coordinate, we arrive at the following problem: given two sets, X and Y , each containing n real numbers, does the Cartesian product $X \times Y$ support a convex polygon with n vertices? We say that $X \times Y$ *contains* a polygon P if every vertex of P is in $X \times Y$; and $X \times Y$ *supports* P if it contains P and no two vertices of P share an x - or y -coordinate. For short, we call the Cartesian product $X \times Y$ an $n \times n$ *grid*.

Not every $n \times n$ grid supports a convex n -gon. This is the case already for $n = 5$ (Figure 1). Several interesting questions arise: can we decide efficiently whether an $n \times n$ -grid supports a convex n -gon? How can we find the largest k such that it supports a convex k -gon? What is the largest k such that *every* $n \times n$ grid supports a convex k -gon? How many convex polygons does an $n \times n$ grid support, or contain? We initiate the study of these questions for convex polygons, and their higher dimensional variants for convex polyhedra.

Our results. We first show that every $n \times n$ grid supports a convex polygon with $(1 - o(1)) \log n$ vertices¹; this bound is tight up to a constant factor: there are $n \times n$ grids that do not support convex polygons with more than $4(\lceil \log n \rceil + 1)$ vertices. We generalize our upper and lower bounds to higher dimensions, and show that every d -dimensional Cartesian product $\prod_{i=1}^d Y_i$, where $|Y_i| = n$ and d is constant, contains $\Omega(\log^{d-1} n)$ points in convex position; this bound is also tight apart from constant factors (Section 2). Next, we present polynomial-time algorithms to find a maximum supported convex polygon that is x - or y -monotone. We show how to efficiently approximate the maximum size of a supported convex polygon up to a factor of two (Section 3). Finally, we present tight asymptotic bounds for the maximum number of convex polygons supported by an $n \times n$ grid (Section 4).

Related work. Erdős and Szekeres proved, as one of the first Ramsey-type results in combinatorial geometry [16], that for every $k \in \mathbb{N}$, a sufficiently large point set in the plane in general position contains k points in convex position. The minimum cardinality of a point set that guarantees k points in convex position is known as the Erdős–Szekeres number, $f(k)$. They proved that $2^{k-2} + 1 \leq f(k) \leq \binom{2k-4}{k-2} + 1 = 4^{k(1-o(1))}$, and conjectured that the lower bound is tight [14]. The current best upper bound, due to Suk [29], is $f(k) \leq 2^{k(1+o(1))}$. In other words, every set of n points in general position in the plane contains $(1 - o(1)) \log n$ points in convex position, and this bound is tight up to lower-order terms.

In dimension $d \geq 3$, the asymptotic growth rate of the Erdős–Szekeres number is not known. By the Erdős–Szekeres theorem, every set of n points in general position in \mathbb{R}^d contains $\Omega(\log n)$ points in convex position (it is enough to find points whose projections onto a generic plane are in convex position). For every constant $d \geq 2$, Károlyi and Valtr [19] and Valtr [30] constructed n -element sets in general position in \mathbb{R}^d in which no more than $O(\log^{d-1} n)$ points are in convex position. Both constructions are recursive, and one of them is related to high-dimensional Horton sets [30]. These bounds are conjectured to be optimal apart from constant factors. Our results establish the same $O(\log^{d-1} n)$ upper bound for Cartesian products $\prod_{i=1}^d Y_i$, where $|Y_i| = n$, for which it is tight apart from constant factors. However our results do not improve the bounds for points in general position.

¹ Throughout this paper all logarithms are in base 2.

Algorithmically, one can find a largest convex cap in a given set of n points in \mathbb{R}^2 in $O(n^2 \log n)$ time by dynamic programming [10], and a largest subset in convex position in $O(n^3)$ time [7, 10]. The same approach can be used for counting the number of convex polygons contained in a given point set [20]. While this approach applies to grids, it is unclear how to include the restriction that each coordinate is used at most once. On the negative side, finding a largest subset in convex position in a point set in \mathbb{R}^d for $d \geq 3$ was recently shown to be NP-hard [15].

There has been significant interest in *counting* the number of convex polygons in various point sets. Answering a question of Hammer, Erdős [13] proved that every set of n points in general position in \mathbb{R}^2 contains $\exp(\Theta(\log^2 n))$ subsets in convex position, and this bound is the best possible. Bárány and Pach [2] showed that the number of convex polygons in an $n \times n$ section of the integer lattice is $\exp(O(n^{2/3}))$. Bárány and Vershik [3] generalized this bound to d -dimensions and showed that there are $\exp(O(n^{(d-1)/(d+1)}))$ convex polytopes in an $n \times \dots \times n$ section of \mathbb{Z}^d . Note that the exponent is sublinear in n for every $d \geq 2$. We prove that an $n \times n$ Cartesian product can contain $\exp(\Theta(n))$ convex polygons, significantly more than integer grids; our bounds are tight up to polynomial factors.

Motivated by integer programming and geometric number theory, lattice polytopes (whose vertices are in \mathbb{Z}^d) have been intensely studied; refer to [1, 4]. However, results for lattices do not extend to arbitrary Cartesian products. Recently, several deep results have been established for Cartesian products in incidence geometry and additive combinatorics [23, 24, 25, 28], while the analogous statements for points sets in general position remain elusive.


Definitions. A polygon P in \mathbb{R}^2 is *convex* if all of its internal angles are strictly smaller than π . A point set in \mathbb{R}^2 is in *convex position* if it is the vertex set of a convex polygon; and it is in *general position* if no three points are collinear. Similarly, a polyhedron P in \mathbb{R}^d is *convex* if it is the convex hull of a finite set of points. A point set in \mathbb{R}^d is in *convex position* if it is the vertex set of a convex polytope; and it is in *general position* if no $d + 1$ points lie on a hyperplane. In \mathbb{R}^d , we say that the x_d -axis is *vertical*, hyperplanes orthogonal to x_d are *horizontal*, and understand the above-below relationship with respect to the x_d -axis. Let \mathbf{e}_d be a standard basis vector parallel to the x_d -axis. A point set P in \mathbb{R}^d is *full-dimensional* if no hyperplane contains P .

We consider special types of convex polygons. Let P be a convex polygon with vertices $((x_1, y_1), \dots, (x_k, y_k))$ in clockwise order. We say that P is a *convex cap* if the x - or y -coordinates are strictly monotonic, and a *convex chain* if both the x - and y -coordinates are strictly monotonic. We distinguish four types of convex caps (resp., chains) based on the monotonicity of the coordinates as follows:

- *convex caps* come in four types $\{\curvearrowright, \curvearrowleft, \smile, \frown\}$. We have
 - $P \in \curvearrowright$ if and only if $(x_i)_{i=1}^k$ strictly increases;
 - $P \in \curvearrowleft$ if and only if $(y_i)_{i=1}^k$ strictly increases;
 - $P \in \smile$ if and only if $(x_i)_{i=1}^k$ strictly decreases;
 - $P \in \frown$ if and only if $(y_i)_{i=1}^k$ strictly decreases;
- *convex chains* come in four types $\{\curvearrowright, \curvearrowleft, \smile, \frown\}$. We have
 - $\curvearrowright = \curvearrowleft \cap \smile, \quad \curvearrowleft = \curvearrowright \cap \frown, \quad \smile = \smile \cap \curvearrowleft, \quad \frown = \frown \cap \smile.$

Initial observations. It is easy to see that for $n = 3, 4$, every $n \times n$ grid supports a convex n -gon. However, there exists a 5×5 grid that does not support any convex pentagon (cf. Fig. 1). Interestingly, every 6×6 grid supports a convex pentagon.

► **Lemma 1.** *Every 6×6 grid $X \times Y$ supports a convex polygon of size at least 5.*



■ **Figure 1** Maximum-size supported convex polygons of respective sizes 3, 4, 4, and 5 in $n \times n$ grids, where n is between 3 and 6.

Proof. Let $X' = X \setminus \{\min(X), \max(X)\}$ and $Y' = Y \setminus \{\min(Y), \max(Y)\}$. The 4×4 grid $X' \times Y'$ supports a convex chain P' of size 3 between two opposite corners of $X' \times Y'$. Then one x -coordinate $x' \in X'$ and one y -coordinate $y' \in Y'$ are not used by P' . Without loss of generality, assume that $P' \in \rho$. Then the convex polygon containing the points of P' and $(x', \min(Y))$ and $(\max(X), y')$ is a supported convex polygon of size 5 on $X \times Y$. ◀

2 Extremal bounds for convex polytopes in Cartesian products

2.1 Lower bounds in the plane

In this section, we show that for every $n \geq 3$, every $n \times n$ grid supports a convex polygon with $\Omega(\log n)$ vertices. The results on the Erdős–Székere number cannot be used directly, since they crucially use the assumption that the given set of points is in general position. An $n \times n$ section of the integer lattice is known to contain $\Theta(n)$ points in general position [12], and this number is conjectured to be $\frac{\pi}{\sqrt{3}}n(1 + o(1))$ [17, 31]. However, this result does not apply to arbitrary Cartesian products. It is worth noting that higher dimensional variants for the integer lattice are poorly understood: it is known that an $n \times n \times n$ section of \mathbb{Z}^3 contains $\Theta(n^2)$ points no three of which are collinear [22], but no similar statements are known in higher dimensions. We use a recent result from incidence geometry.

► **Lemma 2.** (Payne and Wood [21]) *Every set of N points in the plane with at most ℓ collinear, $\ell \leq O(\sqrt{N})$, contains a set of $\Omega(\sqrt{N/\log \ell})$ points in general position.*

► **Lemma 3.** *Every $n \times n$ grid supports a convex polygon of size $(1 - o(1)) \log n$.*


Proof. Every $n \times n$ grid contains a set of $\Omega(\sqrt{n^2/\log n}) = \Omega(n/\sqrt{\log n})$ points in general position by applying Lemma 2 with $N = n^2$ and $\ell = n$. Discarding points with the same x - or y -coordinate reduces the size by a factor at most $\frac{1}{4}$, so this asymptotic bound also holds when coordinates in X and Y are used at most once. By Suk’s result [29], the grid supports a convex polygon with at least $(1 - o(1))(\log(n/\sqrt{\log n})) = (1 - o(1)) \log n$ vertices. ◀

2.2 Upper bounds in the plane

For the upper bound, we construct $n \times n$ Cartesian products that do not support large convex chains. For $n = 8$, such a grid is depicted in Figure 2.

► **Lemma 4.** *For every $n \in \mathbb{N}$, there exists an $n \times n$ grid that contains at most $4(\lceil \log n \rceil + 1)$ points in convex position.*

Proof. Let $g(n)$ be the maximum integer such that for all n -element sets $X, Y \subset \mathbb{R}$, the grid $X \times Y$ supports a convex polygon of size $g(n)$; clearly $g(n)$ is nondecreasing. Let k be the minimum integer such that $n \leq 2^k$; thus $\lceil \log n \rceil \leq k$ and $g(n) \leq g(2^k)$. We show that $g(2^k) \leq 4(k + 1)$ and thereby establish that $g(n) \leq 4(k + 1)$.



■ **Figure 2** An 8×8 grid with no convex chains of size greater than $4 = \log 8 + 1$, where $X = \{0, 1, \dots, 7\}$ and $Y = \{0, 1, 16, 17, 256, 257, 272, 273\}$. Two lines determined by grid points are drawn in blue.

Assume w.l.o.g. that $n = 2^k$, and let $X = \{0, \dots, n - 1\}$. For a k -bit integer m , let m_i be the bit at its i -th position, such that $m = \sum_{i=0}^{k-1} m_i 2^i$. Let $Y = \{\sum_{i=0}^{k-1} m_i (2n)^i \mid 0 \leq m \leq n - 1\}$ (see Fig. 2). Both X and Y are symmetric: $X = \{\max(X) - x \mid x \in X\}$ and $Y = \{\max(Y) - y \mid y \in Y\}$. Thus, it suffices to show that no convex chain $P \in \mathcal{C}$ of size greater than $k + 1$ exists.

Consider two points, $p = (x, y)$ and $p' = (x', y')$, in $X \times Y$ such that $x < x'$ and $y < y'$. Assume $y = \sum_{i=0}^{k-1} m_i (2n)^i$ and $y' = \sum_{i=0}^{k-1} m'_i (2n)^i$. The slope of the line spanned by p and p' is $\text{slope}(p, p') = \frac{\sum_{i=0}^{k-1} (m'_i - m_i) (2n)^i}{x' - x}$. Let j be the largest index such that $m_j \neq m'_j$. Then $y < y'$ implies $m_j < m'_j$, and we can bound the slope as follows:

$$\text{slope}(p, p') \geq \frac{(2n)^j - \sum_{i=0}^{j-1} (2n)^i}{x' - x} > \frac{(2n)^j - 2(2n)^{j-1}}{n - 1} = 2 \cdot (2n)^{j-1},$$

$$\text{slope}(p, p') \leq \frac{\sum_{i=0}^j (2n)^i}{x' - x} \leq \frac{\sum_{i=0}^j (2n)^i}{1} = \frac{(2n)^{j+1} - 1}{2n - 1} < 2 \cdot (2n)^j.$$

Hence, $\text{slope}(p, p') \in I_j = (2 \cdot (2n)^{j-1}, 2 \cdot (2n)^j)$. Let us define the family of intervals I_0, I_1, \dots, I_{k-1} analogously, and note that these intervals are pairwise disjoint. Suppose that some convex chain $P \in \mathcal{C}$ contains more than $k + 1$ points. Since the slopes of the first $k + 1$ edges of P decrease monotonically, by the pigeonhole principle, there must be three consecutive vertices $p = (x, y)$, $p' = (x', y')$, and $p'' = (x'', y'')$ of P such that both $\text{slope}(p, p')$ and $\text{slope}(p', p'')$ are in the same interval, say I_j . Assume that $y = \sum_{i=0}^{k-1} m_i (2n)^i$, $y' = \sum_{i=0}^{k-1} m'_i (2n)^{i+1}$, and $y'' = \sum_{i=0}^{k-1} m''_i (2n)^{i+1}$. Then j is the largest index such that $m_j \neq m'_j$, and also the largest index such that $m'_j \neq m''_j$. Because $m < m' < m''$, we have $m_j < m'_j < m''_j$, which is impossible since each of m_j, m'_j and m''_j is either 0 or 1.

Hence, $X \times Y$ does not contain any convex chain in \curvearrowright of size greater than $k+1$. Analogously, every convex chain in \curvearrowleft , \curvearrowright , or \curvearrowleft has at most $k+1$ vertices. Consequently, $X \times Y$ contains at most $4(k+1)$ points in convex position. ◀

2.3 Upper bounds in higher dimensions


We construct Cartesian products in \mathbb{R}^d , for $d \geq 3$, that match the best known upper bound $O(\log^{d-1} n)$ for the Erdős–Szekeress numbers in d -dimensions for points in general position. Our construction generalizes the ideas from the proof of Lemma 4 to d -space.

► **Theorem 5.** *Let $d \geq 2$ be fixed. For every $n \geq 2$, there exist n -element sets $Y_i \subseteq \mathbb{R}$ for $i = 1, \dots, d$, such that the Cartesian product $Y = \prod_{i=1}^d Y_i$ contains at most $O(\log^{d-1} n)$ points in convex position.*

Proof. We construct point sets recursively. For $d = 2$, the result follows from Lemma 4. For $d \geq 3$ and $0 \leq i \leq j$, we define $S_d(i, j)$ as a Cartesian product of d sets, where the first $d-1$ sets have 2^j elements and the last set has 2^i elements. We then show that $S_d(i, j)$ does not contain the vertex set of any full-dimensional convex polyhedron with more than $2^{d^2-d+1} \cdot i \cdot j^{d-2}$ vertices (there is no restriction on lower-dimensional convex polyhedra).

To initialize the recursion, we define boundary values as follows: For every $j \geq 0$, let $S_2(j, j)$ be the $2^j \times 2^j$ grid defined in the proof of Lemma 4 that does not contain more than $4(j+1)$ points in convex position. Note that every line that contains 3 or more points from $S_2(j, j)$ is axis-parallel (this property was not needed in the proof of Lemma 4). Assume now that $d \geq 3$, and $S_{d-1}(j, j)$ has been defined for all $j \geq 0$; and for all $k = 1, \dots, d$, every k -dimensional flat containing $2^k + 1$ or more points is parallel to at least one coordinate axis. We now construct $S_d(i, j)$ for all $0 \leq i \leq j$ as follows.

Let $S_d(0, j) = S_{d-1}(j, j) \times \{0\}$. For $i = 1, \dots, j$, we define $S_d(i, j)$ as the disjoint union of two translates of $S_d(i-1, j)$. Specifically, let $S_d(i, j) = A \cup B$, where $A = S_d(i-1, j)$ and $B = A + \lambda_d^i \mathbf{e}_d$, where $\lambda_d^i > 0$ is sufficiently large (as specified below) and algebraically independent from the coordinates of $S_d(i-1, j)$, and for all $k = 1, \dots, d$, every k -dimensional flat containing $2^k + 1$ or more points is parallel to at least one coordinate axis. .



■ **Figure 3** A polyhedron $\text{conv}(P)$ in \mathbb{R}^3 , whose projection $\text{conv}(P)^{\text{proj}}$ is a rectangle. Seven points in P are projected onto the four vertices of $\text{conv}(P)^{\text{proj}}$. Overall the *silhouette* of P contains 12 points. Red (blue) vertices are upper (lower); the purple point is both upper and lower.

Let $P \subset S_d(i, j)$ be a full-dimensional set in convex position. The orthogonal projection of $\text{conv}(P)$ to the horizontal hyperplane $x_d = 0$ is a convex polytope in \mathbb{R}^{d-1} that we denote by $\text{conv}(P)^{\text{proj}}$; refer to Fig. 3. The *silhouette* of P is the subset of vertices whose orthogonal

projection to $x_d = 0$ lies on the boundary of $\text{conv}(P)^{\text{proj}}$. Since no three points in P are collinear, at most two points in P are projected to the same point. A point $p \in P$ is an upper (resp., lower) vertex if P lies in the closed halfspace below (resp., above) some tangent hyperplane of $\text{conv}(P)$ at p (a point in P may be both upper and lower vertex).

We prove, by double induction on d and i , the following:

▷ **Claim 6.** If $P \subset S_d(i, j)$ is a full-dimensional set in convex position, then P contains at most $2^{d(d-1)} \cdot i \cdot j^{d-2}$ upper (resp., lower) vertices of $\text{conv}(P)$.

For $d = 2$ and $i = j$, this holds by definition (cf. Lemma 4). For $i = 0$, the set $S_d(0, j) = S_{d-1}(j, j) \times \{0\}$ lies in a horizontal hyperplane in \mathbb{R}^d , and so it is not full-dimensional, hence the claim vacuously holds. By induction, $S_{d-1}(j, j)$ contains at most $2^{(d-1)(d-2)} \cdot j \cdot j^{d-3}$ upper (resp., lower) vertices in \mathbb{R}^{d-1} , hence $S_d(0, j)$ has at most $2 \cdot 2^{(d-1)(d-2)} \cdot j \cdot j^{d-3}$ extreme points in \mathbb{R}^d . Assume that $d \geq 3$, $1 = i \leq j$, and the claim holds for $S_{d-1}(j, j)$. We prove the claim for $S_d(i, j)$. The set $S_d(1, j)$ is the disjoint union of $A = S_d(0, j)$ and $B = S_d(0, j) + \lambda_d^1 \mathbf{e}_d$. Every upper (resp., lower) vertex of $S_d(1, j)$ is an extreme vertex in A or B , hence $S_d(1, j)$ contains at most $4 \cdot 2^{(d-1)(d-2)} \cdot j^{d-2} = 2^{d^2-3d+4} \cdot j^{d-2} < 2^{d^2-d} \cdot 1 \cdot j^{d-2}$ upper (resp., lower) vertices, as required, where we used that $d \geq 3$.

In the general case, we assume that $d \geq 3$, $2 \leq i \leq j$, and the claim holds for $S_{d-1}(j, j)$ and $S_d(i-1, j)$. We prove the claim for $S_d(i, j)$. Recall that $S_d(i, j)$ is the disjoint union of two translates of $S_d(i-1, j)$, namely $A = S_d(i-1, j)$ and $B = S_d(i-1, j) + \lambda_d^i \mathbf{e}_d$. Let $P \subset S_d(i, j)$ be a full-dimensional set in convex position. We partition the upper vertices in P as follows. Let $P_0 \subset P$ be the set of upper vertices whose orthogonal projections to $x_d = 0$ are vertices of $\text{conv}(P)^{\text{proj}}$. For $k = 1, \dots, d-1$, let $P_k \subset P$ be the set of upper vertices whose orthogonal projections to $x_d = 0$ lie in the relative interior of a k -face of $\text{conv}(P)^{\text{proj}}$. By construction, only axis-aligned faces can contain interior points. For the $(d-1)$ -dimensional polytope $\text{conv}(P)^{\text{proj}}$, the axis-aligned k -faces ($k = 1, \dots, d-1$) can be partitioned into $\binom{d-1}{k}$ equivalence classes, based on the set of parallel coordinate axes.

The orthogonal projection of $S_d(i, j)$ to $x_d = 0$ is $S_{d-1}(j, j)$, and the orthogonal projection of P_0 , denoted P_0^{proj} , is the vertex set of a $(d-1)$ -dimensional convex polyhedron in $S_{d-1}(j, j)$. By induction, $|P_0| \leq 2 \cdot 2^{(d-1)(d-2)} \cdot j \cdot j^{d-3} = 2^{d^2-3d+3} j^{d-2}$. We show the following.

▷ **Claim 7.** For every axis-aligned face F of $\text{conv}(P)^{\text{proj}}$, the set of upper vertices that project to the interior of F is contained in either A or B .

Let F be an axis-aligned k -face of $\text{conv}(P)^{\text{proj}}$ for $k \in \{1, 2, \dots, d-1\}$. Let $P(F) \subset P$ be the set of upper vertices whose orthogonal projections lie in the interior of F , and let $P(\partial F)$ be the set of upper vertices whose orthogonal projections lie in the boundary of F . Let $P(\partial F)^{\text{proj}}$ be the orthogonal projection of $P(\partial F)$ to the hyperplane $x_d = 0$. Consider the point set $P' = P(\partial F)^{\text{proj}} \cup P(F)$, and observe that if $P(F) \neq \emptyset$, then it is a vertex set of a $(k+1)$ -dimensional polytope in which all vertices are upper. It remains to show that $P(F) \subseteq A$ or $P(F) \subseteq B$. Suppose, for the sake of contradiction, that $P(F)$ contains points from both A and B . Let p_a be a vertex in $P(F)$ with the maximum x_d -coordinate. The 1-skeleton of $\text{conv}(P')$ contains a x_d -monotonically decreasing path from p_a to an x_d -minimal vertex in P' . Let p_b be the neighbor of p_a along such a path. Then $p_b \in B$ by the choice of p_a . Every hyperplane containing p_a and p_b , in particular the tangent hyperplane of P' containing the edge $p_a p_b$, partitions $P(\partial F)^{\text{proj}}$, which is a contradiction.

We can now finish the proof of Claim 6. By induction on $S_d(i-1, j)$, we have

$$\begin{aligned} |P_k| &\leq \binom{d-1}{k} \cdot 2^{(d-k-1)(d-k-2)+1} j^{d-k-1} \cdot 2^{k(k-1)} \cdot (i-1) \cdot j^{k-1} \\ &\leq \binom{d-1}{k} \cdot 2^{(d-1)(d-2)} \cdot (i-1) \cdot j^{d-2}. \end{aligned}$$

Altogether, the number of upper vertices is

$$\sum_{k=0}^{d-1} |P_k| \leq 2^{d^2-3d+3} j^{d-2} + \sum_{k=1}^{d-1} \binom{d-1}{k} 2^{(d-1)(d-2)} \cdot (i-1) \cdot j^{d-2} < 2^{d(d-1)} \cdot i \cdot j^{d-2},$$

as required, where we used the binomial theorem and the inequality $d \geq 3$. \blacktriangleleft

2.4 Lower bounds in higher dimensions

The proof technique in Section 2.1 is insufficient for establishing a lower bound of $\Omega(\log^{d-1} n)$ for $d \geq 3$. Whereas a d -dimensional $n \times \dots \times n$ grid contains $\Omega(n^\delta)$ points in general position for some $\delta = \delta(d) > 0$ [6], the current best lower bound on the number of points in convex position in any set of n points in general position in \mathbb{R}^d is $\Omega(\log n)$; the conjectured value is $\Omega(\log^{d-1} n)$. Instead, we rely on the structure of Cartesian products and induction on d . Our main result in this section is the following.

► **Theorem 8.** *Every d -dimensional Cartesian product $\prod_{i=1}^d Y_i$, where $|Y_i| = n$ and d is fixed, contains $\Omega(\log^{d-1} n)$ points in convex position.*

We say that a strictly increasing sequence of real numbers $A = (a_1, \dots, a_n)$, has the *monotone differences* property (for short, A is *MD*), if either $a_{i+1} - a_i > a_i - a_{i-1}$, or $a_{i+1} - a_i < a_i - a_{i-1}$ for $i = 2, \dots, n-1$. Furthermore, the sequence A is *r -MD* for some $r > 1$ if either $a_{i+1} - a_i \geq r(a_i - a_{i-1})$, or $a_{i+1} - a_i \leq (a_i - a_{i-1})/r$ for $i = 2, \dots, n-1$.

A finite set $X \subseteq \mathbb{R}$ is *MD* (resp., *r -MD*) if its elements arranged in increasing order form an MD (resp., *r -MD*) sequence. These sequences are intimately related to convexity: a strictly increasing sequence $A = (a_1, \dots, a_n)$ is MD if and only if there exists a monotone (increasing or decreasing) convex function $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $a_i = f(i)$ for all $i = 1, \dots, n$. MD sets have been studied in additive combinatorics [11, 18, 23, 27].

We first show that every n -element set $X \subseteq \mathbb{R}$ contains an MD subset of size $\Omega(\log n)$, and this bound is the best possible (Lemma 9). In contrast, every n -term arithmetic progression contains an MD subsequence of $\Theta(\sqrt{n})$ terms: for example $(0, \dots, n-1)$ contains the subsequence $(i^2 : i = 0, \dots, \lfloor \sqrt{n-1} \rfloor)$. We then show that for constant $d \geq 2$, the d -dimensional Cartesian product of MD sets of size n contains $\Theta(n^{d-1})$ points in convex position. The combination of these results immediately implies that every $n \times \dots \times n$ Cartesian product in \mathbb{R}^d contains $\Omega(\log^{d-1} n)$ points in convex position.

The following lemma gives a lower bound for MD sequences. It is known that a monotone sequence of n reals contains a 2-MD sequence (satisfying the so-called *doubling differences condition* [26]) of size $\Omega(\log n)$ [5, Lemma 4.1]; see also [8] for related recent results.


► **Lemma 9.** *Every set of n real numbers contains an MD subset of size $\lfloor (\log n)/2 \rfloor + 1$. For every $n \in \mathbb{N}$, there exists a set of n real numbers in which the size of every MD subset is at most $\lceil \log n \rceil + 1$.*

Proof. Let $X = (x_0, \dots, x_{n-1})$ be a strictly increasing sequence. Assume w.l.o.g. that $n = 2^\ell + 1$ for some $\ell \in \mathbb{N}$. We construct a sequence of nested intervals

$$[a_0, b_0] \supset [a_1, b_1] \supset \dots \supset [a_\ell, b_\ell]$$

such that the endpoints of the intervals are in X and the lengths of the intervals decrease by factors of 2 or higher, that is, $b_i - a_i \leq (b_{i-1} - a_{i-1})/2$ for $i = 1, \dots, \ell$.

We start with the interval $[a_0, b_0] = [x_0, x_{n-1}]$; and for every $i = 0, \dots, \ell - 1$, we divide $[a_i, b_i]$ into two intervals at the median, and recurse on the shorter interval.



■ **Figure 4** A sequence X of 17 elements and nested intervals $[a_0, b_0] \supset \dots \supset [a_4, b_4]$.

By partitioning $[a_i, b_i]$ at the median, the algorithm maintains the invariant that $[a_i, b_i]$ contains $2^{\ell-i} + 1$ elements of X . Note that for every $i = 1, \dots, \ell$, we have either $(a_{i-1} = a_i$ and $b_{i-1} < b_i)$ or $(a_{i-1} = a_i$ and $b_i < b_{i-1})$. Consequently, the sequences $A = (a_0, a_1, \dots, a_\ell)$ and $B = (b_\ell, b_{\ell-1}, \dots, b_0)$ both increase (not necessarily strictly), and at least one of them contains at least $1 + \ell/2$ distinct terms. Assume w.l.o.g. that A contains at least $1 + \ell/2$ distinct terms. Let $C = (c_0, \dots, c_k)$ be a maximal strictly increasing subsequence of A . Then $k \geq \ell/2 = \lfloor (\log n)/2 \rfloor$.

We show that C is an MD sequence. Let $i \in \{1, \dots, k - 1\}$. Assume that $c_i = a_j = \dots = a_{j'}$ for consecutive indices j, \dots, j' . Then $c_{i-1} = a_{j-1}$, $c_i = a_j$, and $c_{i+1} = a_{j'+1}$. By construction, $c_i \in [a_{j-1}, b_{j-1}] = [c_{i-1}, b_j]$ such that $c_i - c_{i-1} \geq b_j - c_i$. Similarly, $c_{i+1} \in [a_{j'}, b_{j'}] = [c_i, b_{j'}]$ such that $c_{i+1} - c_i \geq b_{j'} - c_{i+1}$. However, $[a_j, b_j] \subset [a_{j-1}, b_{j-1}]$. As required, this yields

$$c_{i+1} - c_i = a_{j'+1} - a_j < b_j - a_j \leq \frac{b_{j-1} - a_{j-1}}{2} \leq a_j - a_{j-1} = c_i - c_{i-1}.$$


The upper bound construction is the point set Y defined in the proof of Lemma 4, for which every chain in \sphericalangle or \searrow supported by $\{0, \dots, n - 1\} \times Y$ has at most $\lceil \log n \rceil + 1$ vertices. Let $\{b_0, \dots, b_{\ell-1}\} \subset Y$ be an MD subset such that $b_0 < \dots < b_{\ell-1}$. Then $\{(i, b_i) : i = 0, \dots, \ell - 1\} \subset X \times Y$ is in \sphericalangle or \searrow . Consequently, every MD subset of Y has at most $\lceil \log n \rceil + 1$ terms, as claimed. ◀

We show how to use Lemma 9 to establish a lower bound in the plane. While this approach yields worse constant coefficients than Lemma 3, its main advantage is that it generalizes to higher dimensions (see Lemma 12 below).

► **Lemma 10.** *The Cartesian product of two MD sets, each of size n , supports n points in convex position.*

Proof. Let $A = \{a_1, \dots, a_n\}$ and $B = \{b_1, \dots, b_n\}$ be MD sets such that $a_i < a_{i+1}$ and $b_i < b_{i+1}$ for $i = 1, \dots, n - 1$. We may assume, by applying a reflection if necessary, that $a_{i+1} - a_i < a_i - a_{i-1}$ and $b_{i+1} - b_i < b_i - b_{i-1}$, for $i = 2, \dots, n - 1$ (see Fig. 5).

We define $P \subset A \times B$ as the set of n points (a_i, b_j) such that $i + j = n + 1$. By construction, every horizontal (vertical) line contains at most one point in P . Since the differences $a_i - a_{i-1}$ are positive and strictly decrease in i ; and the differences $b_{\ell-i} - b_{\ell-i-1}$ are negative and their absolute values strictly increase in i , the slopes $(b_{\ell-i} - b_{\ell-i-1}) / (a_i - a_{i-1})$ strictly decrease, which proves the convexity of P . ◀



■ **Figure 5** A 7×7 grid $\{a_1, \dots, a_7\} \times \{b_1, \dots, b_7\}$, where the differences between consecutive x -coordinates (resp., y -coordinates) decrease by factors of 2 or higher. The point sets $\{(0, 0)\} \cup \{(a_i, b_j) : i + j = k\}$, for $k = 2, \dots, 8$, form nested convex chains.

► **Lemma 11.** *The Cartesian product of three MD sets, each of size n , contains $\binom{n+1}{2}$ points in convex position.*

Proof. Let $A = \{a_1, \dots, a_n\}$, $B = \{b_1, \dots, b_n\}$, and $C = \{c_1, \dots, c_n\}$ be MD sets, where the elements are labeled in increasing order. We may assume, by applying a reflection in the x -, y -, or z -axis if necessary, that $a_{i+1} - a_i < a_i - a_{i-1}$, $b_{i+1} - b_i < b_i - b_{i-1}$, and $c_{i+1} - c_i < c_i - c_{i-1}$, for $i = 2, \dots, n-1$. For $i, j, k \in \{1, \dots, n\}$, let $p_{i,j,k} = (a_i, b_j, c_k) \in A \times B \times C$. We now let $P = \{p_{i,j,k} : i+j+k = n+2\}$. It is clear that $|P| = \sum_{i=1}^n i = \binom{n+1}{2}$. We let $P' = P \cup \{p_{1,1,1}\}$ and show that the points in P' are in convex position.

By Lemma 10, the points in P' lying in the planes $x = a_1$, $y = b_1$, and $z = c_1$ are each in convex position. These convex $(n+1)$ -gons are faces of the convex hull of P , denoted $\text{conv}(P)$. We show that the remaining faces of $\text{conv}(P)$ are the triangles $T'_{i,j,k}$ spanned by $p_{i,j,k}$, $p_{i,j+1,k-1}$, and $p_{i+1,j,k-1}$; and the triangles $T''_{i,j,k}$ spanned by $p_{i,j,k}$, $p_{i,j-1,k+1}$, and $p_{i-1,j,k+1}$.

The projection of these triangles to the xy -plane is shown in Fig. 5. By construction, the union of these faces is homeomorphic to a sphere. It suffices to show that the dihedral angle between any two adjacent triangles is convex. Without loss of generality, consider the triangle $T'_{i,j,k}$, which is adjacent to at most three other triangles: $T''_{i+1,j,k-1}$, $T''_{i,j+1,k-1}$, and $T''_{i+1,j+1,k-2}$ (if they exist). Consider first the triangles $T'_{i,j,k}$ and $T''_{i+1,j+1,k-2}$. They share the edge $p_{i+1,j-1,k+1}p_{i-1,j+1,k+1}$, which lies in the plane $z = c_{k+1}$. The orthogonal projections of these triangles to this plane are congruent, however their extents in the z -axis are $c_{i+1} - c_i$ and $c_i - c_{i-1}$, respectively. Since $c_{i+1} - c_i < c_i - c_{i-1}$, their dihedral angle is convex. Similarly, the dihedral angles between $T'_{i,j,k}$ and $T''_{i+1,j,k-1}$ (resp., $T''_{i,j+1,k-1}$) is convex because $a_{i+1} - a_i < a_i - a_{i-1}$ and $b_{i+1} - b_i < b_i - b_{i-1}$. ◀

The proof technique of Lemma 11 generalizes to higher dimensions (details are provided in the full version):

► **Lemma 12.** *For every constant $d \geq 2$, the Cartesian product of d MD sets, each of size n , contains $\Omega(n^{d-1})$ points in convex position.*

Now Theorem 8 follows from Lemma 9 and Lemma 12.

3 Algorithms

In this section, we describe polynomial-time algorithms for (i) finding convex chains and caps of maximum size; and (ii) approximating the maximum size of a convex polygon; where these structures are *supported* by a given grid. The main challenge is to ensure that the vertices of the convex polygon (resp., cap or chain) have distinct x - and y -coordinates. The coordinates of a point $p \in X \times Y$ are denoted by $x(p)$ and $y(p)$.

As noted in Section 1, efficient algorithms are available for finding a largest convex polygon or convex cap *contained* in a planar point set. Edelsbrunner and Guibas [10, Thm. 5.1.2] use the dual line arrangement of N points in the plane and dynamic programming to find the maximum size of a convex cap in \curvearrowright in $O(N^2)$ time and $O(N)$ space; the same bounds hold for \curvearrowleft , \curvearrowright , and \curvearrowright . A longest convex cap can be also returned in $O(N^2 \log N)$ time and $O(N \log N)$ space. It is straightforward to adapt their algorithm to find the maximum size of a convex cap in \curvearrowright , and report a longest such chain within the same time and space bounds. Since x - and y -coordinates do not repeat in a convex chain, we obtain the following.

► **Theorem 13.** *In a given $n \times n$ grid, the maximum size of a supported convex chain can be computed in $O(n^4)$ time and $O(n^2)$ space; and a supported convex chain of maximum size can be computed in $O(n^4 \log n)$ time and $O(n^2 \log n)$ space.*


We make use of the following general observation:

► **Observation 14.** *If a supported convex polygon P is in a set \curvearrowright , \curvearrowleft , \curvearrowright , \curvearrowleft , \curvearrowright , \curvearrowright , \curvearrowleft , or \curvearrowright , then every subsequence of P is in the same set. That is, these classes are hereditary.*

3.1 Convex caps

For computing the maximum size of a convex cap in \curvearrowright , we need to be careful to use each y -coordinate at most once. We design an algorithm that finds the maximum size of *two* convex chains that use distinct y -coordinates by dynamic programming. Specifically, for two edges $l = (l_1, l_2)$ and $r = (r_1, r_2)$, we compute the maximum total size $C(l, r)$ of a pair of chains $A \in \curvearrowright$ and $B \in \curvearrowleft$ such that their vertices use distinct y -coordinates and such that the last two vertices of A are l_1 and l_2 (or $A = (l_1)$ if $l_1 = l_2$), and the first two vertices of B are r_1 and r_2 (or $B = (r_1)$ if $r_1 = r_2$). We use the dynamic programming algorithm of [10] to find $L(p_1, p_2)$ (resp., $R(p_1, p_2)$), the size of a largest convex chain P in \curvearrowright (resp., \curvearrowleft), ending (resp., starting) with vertices p_1 and p_2 , or $P = (p_1)$ if $p_1 = p_2$.

The desired quantity $C(l, r)$ can be computed by dynamic programming. By Observation 14, we can always safely eliminate the highest vertex of the union of the two chains, to find a smaller subproblem, as this vertex cannot be (implicitly) part of the optimal solution to a subproblem. In particular, if l is a single vertex and it is highest, we can simply use the value of $R(r_1, r_2)$, incrementing it by one for the one vertex of l . Analogously, we handle the case if r is or both l and r are a single vertex. The interesting case is when both chains end in an edge. Here, we observe that we can easily check whether l and r use unique coordinates.



■ **Figure 6** Illustration for the cases of $C(l, r)$. (a) Invalid configuration, as $y(r_2) = y(l_2)$. (b) r is a single point above l_2 , so we look for the longest chain ending in l . (c) Removing the topmost point (from l in this case), testing all valid possible v to find the longest chain. Note that the left and right chain may not complete to a cap – this is checked separately. (d) We only need to test whether l and r make a cap (purple dotted line) to check whether the $C(l, r)$ entry should be considered.

If not, then this subproblem is invalid; otherwise, we may find a smaller subproblem by eliminating the highest vertex and comparing all possible subchains that could lead to it.

With the reasoning above, we obtain the recurrence below; see Fig. 6(a–c) for illustration. The first case eliminates invalid edges, and edge pairs that use a y -coordinate more than once. In all remaining cases, we assume that $l \in \nearrow$, $r \in \searrow$, and l and r use distinct y -coordinates.

$$C(l, r) = \begin{cases} -\infty & \text{if } l_1 \neq l_2 \text{ and } l \notin \nearrow, \text{ or} \\ & r_1 \neq r_2 \text{ and } r \notin \searrow, \text{ or} \\ & \{y(l_1), y(l_2)\} \cap \{y(r_1), y(r_2)\} \neq \emptyset \\ 2 & \text{otherwise, if } l_1 = l_2 \text{ and } r_1 = r_2 \\ L(l_1, l_2) + 1 & \text{otherwise, if } r_1 = r_2 \text{ and } y(l_2) < y(r_1) \\ R(r_1, r_2) + 1 & \text{otherwise, if } l_1 = l_2 \text{ and } y(l_2) > y(r_1) \\ \max_{(v, l_1, l_2) \in \nearrow \text{ or } v = l_1} C((v, l_1), r) + 1 & \text{otherwise, if } y(l_2) > y(r_1) \\ \max_{(r_1, r_2, v) \in \searrow \text{ or } v = r_1} C(l, (r_2, v)) + 1 & \text{otherwise, } y(l_2) < y(r_1). \end{cases}$$

Let $E = (X \times Y)^2$ denote the number of pairs (edges) in the grid, from which we take l and r . As $|E| = O(n^4)$, we can compute $C(l, r)$ for all l and r in $O(|E|^2 |X \times Y|) = O(n^{10})$ time and $O(|E|^2) = O(n^8)$ space. With $C(l, r)$, we can easily find the size of a maximum size cap P in \nearrow , using the observation below, and analogous observations for the special case $k = 1$ and/or $\ell = 1$ (see Fig. 6(d)).

► **Observation 15.** If $A = (a_1, \dots, a_k) \in \curvearrowright$ and $B = (b_1, \dots, b_\ell) \in \curvearrowleft$ with $k \geq 2, \ell \geq 2$ and $(a_{k-1}, a_k, b_1, b_2) \in \curvearrowright$ and A and B use distinct y -coordinates, then $(a_1, \dots, a_k, b_1, \dots, b_\ell)$ lies in \curvearrowright and has size $k + \ell$.

Note that the condition $(a_{k-1}, a_k, b_1, b_2) \in \curvearrowright$ implies that the x -coordinates are disjoint.

► **Theorem 16.** For a given $n \times n$ grid, a supported convex cap of maximum size can be computed in $O(n^{10})$ time and $O(n^8)$ space.

Although computing the maximum size of a supported convex polygon remains elusive, we can easily devise a constant-factor approximation algorithm by eliminating duplicate coordinates as described in Section 3.3 of the full version.

4 The maximum number of convex polygons


Let $F(n)$ be the maximum number of convex polygons that can be present in an $n \times n$ grid, with no restriction on the number of times each coordinate is used. Let $G(n)$ be this number where all $2n$ grid lines are used (i.e., each grid line contains at least one vertex of the polygon). Let $\bar{F}(n)$ and $\bar{G}(n)$ be the corresponding numbers where each grid line is used at most once (so $\bar{F}(n)$ counts the maximum number of supported convex polygons). By definition, we have $F(n) \geq G(n) \geq \bar{G}(n)$ and $F(n) \geq \bar{F}(n) \geq \bar{G}(n)$ for all $n \geq 2$. We prove the following theorem, in which the $\Theta^*(\cdot)$ notation hides polynomial factors in n .

► **Theorem 17.** The following bounds hold:

$$F(n) = \Theta^*(16^n), \quad \bar{F}(n) = \Theta^*(9^n), \quad G(n) = \Theta^*(9^n), \quad \bar{G}(n) = \Theta^*(4^n).$$

4.1 Upper bounds

We first prove that $F(n) = O(n \cdot 16^n)$ by encoding each convex polygon in a unique way, so that the total number of convex polygons is bounded by the total number of encodings. Recall that a convex polygon P can be decomposed into four convex chains $\curvearrowright_P, \curvearrowleft_P, \curvearrowdown_P, \curvearrowup_P$, with only extreme vertices of P appearing in multiple chains. Let $\zeta_P = \curvearrowright_P \cup \curvearrowleft_P$ and $\upsilon_P = \curvearrowleft_P \cup \curvearrowdown_P$. To encode P , we assign the following number to each of the $2n$ grid lines ℓ (see Fig. 7): 0 if ℓ is not incident on any vertex of P , 3 if ℓ is incident on multiple vertices of P , 1 if ℓ is incident on one vertex of P and that vertex lies on ζ_P if ℓ is horizontal, or on υ_P if ℓ is vertical, and 2 otherwise. We also record the index of the horizontal line containing the leftmost vertex of P (pick the topmost of these if there are multiple leftmost points).



■ **Figure 7** Encoding the grid lines.

22:14 Convex Polygons in Cartesian Products

Since each of the $2n$ grid lines is assigned one of 4 possible values, and there are n horizontal lines, the total number of encodings is $O(n \cdot 4^{2n}) = O(n \cdot 16^n)$. All that is left to show is that each encoding corresponds to at most one convex polygon.

First, observe that if P is a convex chain, say in \nearrow , then the set of grid lines containing a vertex of P uniquely defines P : since both coordinates change monotonically, the i -th vertex of P must be the intersection of the i -th horizontal and vertical lines. So all we need to do to reconstruct P is to identify the set of lines that make up each convex chain.

Since we know the location of the (topmost) leftmost vertex of P , we know where \nearrow_P starts. Every horizontal line above this point labelled with a 1 or 3 must contain a vertex of \nearrow_P ; let k be the number of such lines. Since the x -coordinates are monotonic as well, \nearrow_P ends at the k -th vertical line labelled with a 2 or 3. The next chain, \searrow_P , starts either at the end of \nearrow_P , if the horizontal line is labelled with a 1, or at the intersection of this horizontal line with the next vertical line labelled with a 2 or 3, if this horizontal line is labelled with a 3. We can find the rest of the chains in a similar way. Thus, $F(n) = O(n \cdot 16^n)$.

The upper bounds for $\bar{F}(n)$, $G(n)$, and $\bar{G}(n)$ are analogous, except that certain labels are excluded. For the number of supported convex polygons $\bar{F}(n)$, each grid line is used at most once, which means that the label 3 cannot be used. Thus, $\bar{F}(n) = O(n \cdot 3^{2n}) = O(n \cdot 9^n)$. Similarly, for $G(n)$, all grid lines contain at least one vertex of the polygon, so the label 0 cannot be used. Therefore $G(n) = O(n \cdot 3^{2n}) = O(n \cdot 9^n)$. Finally, for $\bar{G}(n)$, every grid line contains exactly one vertex of the polygon, so neither 0 nor 3 can be used as labels. This gives $\bar{G}(n) = O(n \cdot 2^{2n}) = O(n \cdot 4^n)$ possibilities.


4.2 Lower bounds

Assume that $n = 2m + 3$, where $m \in \mathbb{N}$ satisfies suitable divisibility conditions, as needed. All four lower bounds use the same grid, constructed as follows (see Fig. 8).

$$\begin{aligned} X &= \{1, \dots, n-1\} & Y^- &= \{y_1, \dots, y_{m+2}\}, \text{ where } y_i = n^i \\ Y &= Y^- \cup Y^+ & Y^+ &= \{z_1, \dots, z_{m+2}\}, \text{ where } z_i = 2 \cdot y_{m+2} - y_i \end{aligned}$$

Note that this results in an $(n-1) \times (n-1)$ grid, since $y_{m+2} = z_{m+2}$. To obtain an $n \times n$ grid, we duplicate the median grid lines in both directions and offset them by a sufficiently small distance $\varepsilon > 0$. The resulting grid has the property that any three points p, q, r in the lower half $X \times Y^-$ with $x(p) < x(q) < x(r)$ and $y(p) < y(q) < y(r)$ make a left turn at q . To see this, suppose that $y(p) = n^i$, $y(q) = n^j$, and $y(r) = n^k$, for some $1 \leq i < j < k \leq n$. Then the slope of pq is strictly smaller than the slope of qr , since

$$\text{slope}(qr) = \frac{n^k - n^j}{x(r) - x(q)} \geq \frac{n^{j+1} - n^j}{n-1} = n^j > \frac{n^j - n^i}{1} \geq \frac{n^j - n^i}{x(q) - x(p)} = \text{slope}(pq).$$



■ **Figure 8** The $n \times n$ grid defined in Section 4.2, with $n = 9 = 2m + 3$ for $m = 3$, before doubling the median lines. The segments (parts of grid lines) incident to vertices are drawn in blue.

Thus, any sequence of points with increasing x - and y -coordinates in the lower half is in \swarrow . By symmetry, such a sequence in the upper half $X \times Y^+$ is in \nearrow . Analogously, points with increasing x -coordinates and decreasing y -coordinates are in \searrow if they are in the lower half and \nwarrow if they are in the upper half.

We first derive lower bounds on $\bar{G}(n)$ and $G(n)$ by constructing a large set of convex polygons that use each grid line at least once. Then we use these bounds to derive the bounds on $\bar{F}(n)$ and $F(n)$. The polygons we construct all share the same four extreme vertices, which lie on the intersections of the grid boundary with the duplicated median grid lines. Specifically, the leftmost and rightmost vertices are the intersections of the duplicate horizontal medians with the left and right boundary, and the highest and lowest vertices are the intersections of the duplicate vertical medians with the top and bottom boundary. Since each of these median lines now contains a vertex, we can choose additional vertices from the remaining $2m$ grid lines in each direction.

To construct each polygon, select $m/2$ vertical grid lines left of the median to participate in the bottom chain, and do the same right of the median. Likewise, select $m/2$ horizontal grid lines above and below the median, respectively, to participate in the left chain. The remaining grid lines participate in the other chain (top or right). This results in a polygon with $m/2$ vertices in each quadrant of the grid (excluding the extreme vertices). The convexity follows from our earlier observations. The total number of such polygons is

$$\binom{m}{\frac{m}{2}}^4 = \Theta\left(\left(m^{-\frac{1}{2}}2^m\right)^4\right) = \Theta(m^{-2}2^{4m}) = \Theta(n^{-2}2^{2n}) = \Theta(n^{-2}4^n) = \Theta^*(4^n).$$

The first step uses the following estimate, which can be derived from Stirling’s formula for the factorial [9]. Let $0 < \alpha < 1$, then

$$\binom{n}{\alpha n} = \Theta(n^{-\frac{1}{2}}2^{H(\alpha)n}), \text{ where } H(\alpha) = -\alpha \log \alpha - (1 - \alpha) \log(1 - \alpha).$$

For the lower bound on $G(n)$, the only difference is that we now allow grid lines to contain vertices in two chains. We obtain a maximum when we divide the grid lines evenly between the three groups (bottom chain, top chain, both chains). Thus, we select $m/3$ vertical grid lines left of the median to participate in the bottom chain, another $m/3$ to participate in the top chain and the remaining $m/3$ participate in both. We repeat this selection to the right of the median and on both sides of the median horizontal line. As before, this results in a convex polygon with the same number of vertices in each quadrant of the grid – exactly $2m/3$ this time. The number of such polygons is

$$\begin{aligned} \binom{m}{\frac{m}{3}}^4 \binom{\frac{2m}{3}}{\frac{m}{3}}^4 &= \Theta\left(\left(m^{-\frac{1}{2}}2^{H(\frac{1}{3})m} \cdot m^{-\frac{1}{2}}2^{H(\frac{1}{2})\frac{2m}{3}}\right)^4\right) \\ &= \Theta\left(m^{-4}2^{4m(\log 3 - \frac{2}{3} + \frac{2}{3})}\right) = \Theta(n^{-4}2^{2n \log 3}) = \Theta(n^{-4}9^n) = \Theta^*(9^n). \end{aligned}$$

We translate these bounds to bounds on $\bar{F}(n)$ and $F(n)$ in the full version.

4.3 The maximum number of weakly convex polygons

A polygon P in \mathbb{R}^2 is *weakly convex* if all of its internal angles are less than or equal to π . We summarize the bounds we obtain in the following (details are provided in the full version):

► **Theorem 18.** *Let $W(n)$ denote the maximum number of weakly convex polygons that can be present in an $n \times n$ grid. Then $W(n) = \Omega(16^n)$ and $W(n) = O^*(16^n)$.*

References

- 1 Imre Bárány. Random points and lattice points in convex bodies. *Bulletin of the American Mathematical Society*, 45:339–365, 2008.
- 2 Imre Bárány and János Pach. On the number of convex lattice polygons. *Combinatorics, Probability and Computing*, 1(4):295–302, 1992.
- 3 Imre Bárány and Anatoly Moiseevich Vershik. On the number of convex lattice polytopes. *Geometric and Functional Analysis*, 2(4):381–393, 1992.
- 4 Alexander Barvinok. Lattice points and lattice polytopes. In Jacob E. Goodman, Joseph O’Rourke, and Csaba D. Tóth, editors, *Handbook of Discrete and Computational Geometry*, chapter 7, pages 185–210. CRC Press, Boca Raton, FL, 3rd edition, 2017.
- 5 Boris Bukh and Jiří Matoušek. Erdős–Szekeres-type statements: Ramsey function and decidability in dimension 1. Manuscript, 2012. Available from: <https://arxiv.org/abs/1207.0705>.
- 6 Jean Cardinal, Csaba D. Tóth, and David R. Wood. A note on independent hyperplanes and general position subsets in d -space. *Journal of Geometry*, 108(1):33–43, 2017.
- 7 Václav Chvátal and G. T. Klincksiek. Finding largest convex subsets. *Congressus Numerantium*, 29:453–460, 1980.
- 8 David Conlon, Jacob Fox, János Pach, Benny Sudakov, and Andrew Suk. Ramsey-type results for semi-algebraic relations. *Transactions of the American Mathematical Society*, 366:5043–5065, 2014.
- 9 Adrian Dumitrescu, André Schulz, Adam Sheffer, and Csaba D. Tóth. Bounds on the maximum multiplicity of some common geometric graphs. *SIAM Journal on Discrete Mathematics*, 27(2):802–826, 2013.
- 10 Herbert Edelsbrunner and Leonidas J. Guibas. Topologically Sweeping an Arrangement. *Journal of Computer and System Sciences*, 38(1):165–194, 1989.
- 11 György Elekes, Melvyn B. Nathanson, and Imre Z. Ruzsa. Convexity and Sumsets. *Journal of Number Theory*, 83(2):194–201, 2000.
- 12 Paul Erdős. Appendix, in Klaus F. Roth, On a problem of Heilbronn. *Journal of the London Mathematical Society*, 26:198–204, 1951.
- 13 Paul Erdős. Some more problems on elementary geometry. *Gazette of the Australian Mathematical Society*, 5(2):52–54, 1978.
- 14 Paul Erdős and György Szekeres. On some extremum problems in elementary geometry. *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae Sectio Mathematica*, 3-4:53–62, 1960/1961.
- 15 Panos Giannopoulos, Christian Knauer, and Daniel Werner. On the Computational Complexity of Erdős–Szekeres and Related Problems in \mathbb{R}^3 . In *Proc. 21st European Symposium on Algorithms*, volume 8125 of *LNCS*, pages 541–552. Springer, 2013.
- 16 Ronald L. Graham. Euclidean Ramsey theory. In Jacob E. Goodman, Joseph O’Rourke, and Csaba D. Tóth, editors, *Handbook of Discrete and Computational Geometry*, chapter 11, pages 281–297. CRC Press, Boca Raton, FL, 3rd edition, 2017.
- 17 Richard K. Guy and Patrick A. Kelly. The No-Three-in-Line-Problem. *Canadian Mathematical Bulletin*, 11:527–531, 1968.
- 18 Norbert Hegyvári. On consecutive sums in sequences. *Acta Mathematica Hungarica*, 48(1-2):193–200, 1986.
- 19 Gyula Károlyi and Pavel Valtr. Configurations in d -space without large subsets in convex position. *Discrete & Computational Geometry*, 30(2):277–286, 2003.
- 20 Joseph S.B. Mitchell, Günter Rote, Gopalakrishnan Sundaram, and Gerhard Woeginger. Counting convex polygons in planar point sets. *Information Processing Letters*, 56(1):45–49, 1995.
- 21 Michael S. Payne and David R. Wood. On the general position subset selection problem. *SIAM Journal on Discrete Mathematics*, 27(4):1727–1733, 2013.
- 22 Attila Pór and David R. Wood. No-three-in-line-in-3D. *Algorithmica*, 47(4):481–488, 2007.

- 23 Orit E. Raz, Micha Sharir, and Ilya D. Shkredov. On the number of unit-area triangles spanned by convex grids in the plane. *Computational Geometry: Theory and Applications*, 62:25–33, 2017.
- 24 Orit E. Raz, Micha Sharir, and József Solymosi. Polynomials vanishing on grids: The Elekes–Rónyai problem revisited. *American Journal of Mathematics*, 138(4):1029–1065, 2016.
- 25 Orit E. Raz, Micha Sharir, and Frank De Zeeuw. Polynomials vanishing on Cartesian products: The Elekes–Szabó theorem revisited. *Duke Mathematical Journal*, 165(18):3517–3566, 2016.
- 26 David Alan Rosenthal. *The classification of the order indiscernibles of real closed fields and other theories*. PhD thesis, University of Wisconsin–Madison, 1981.
- 27 Tomasz Schoen and Ilya D. Shkredov. On Sumsets of Convex Sets. *Combinatorics, Probability and Computing*, 20(5):793–798, 2011.
- 28 Ryan Schwartz, József Solymosi, and Frank de Zeeuw. Extensions of a result of Elekes and Rónyai. *Journal of Combinatorial Theory, Series A*, 120(7):1695–1713, 2013.
- 29 Andrew Suk. On the Erdős–Szekeres convex polygon problem. *Journal of the American Mathematical Society*, 30:1047–1053, 2017.
- 30 Pavel Valtr. Sets in \mathbb{R}^d with no large empty convex subsets. *Discrete Mathematics*, 108(1–3):115–124, 1992.
- 31 Eric W. Weisstein. No-Three-in-a-Line-Problem. online, 2005. Available from: <http://mathworld.wolfram.com/No-Three-in-a-Line-Problem.html>.