

Hamiltonicity below Dirac's condition

Citation for published version (APA):

Jansen, B. M. P., Kozma, L., & Nederlof, J. (2019). Hamiltonicity below Dirac's condition. *arXiv*, 2019, [1902.01745v1]. <https://doi.org/10.48550/arXiv.1902.01745>

DOI:

[10.48550/arXiv.1902.01745](https://doi.org/10.48550/arXiv.1902.01745)

Document status and date:

Published: 01/01/2019

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl


providing details and we will investigate your claim.

Hamiltonicity below Dirac's condition

Bart M.P. Jansen¹

Eindhoven University of Technology

b.m.p.jansen@tue.nl

 <http://orcid.org/0000-0001-8204-1268>

László Kozma²

Freie Universität Berlin

laszlo.kozma@fu-berlin.de

Jesper Nederlof³

Eindhoven University of Technology

j.nederlof@tue.nl

Abstract

Dirac's theorem (1952) is a classical result of graph theory, stating that an n -vertex graph ($n \geq 3$) is Hamiltonian if every vertex has degree at least $n/2$. Both the value $n/2$ and the requirement for *every vertex* to have high degree are necessary for the theorem to hold.

In this work we give efficient algorithms for determining Hamiltonicity when either of the two conditions are relaxed. More precisely, we show that the Hamiltonian cycle problem can be solved in time $c^k \cdot n^{O(1)}$, for some fixed constant c , if at least $n - k$ vertices have degree at least $n/2$, or if all vertices have degree at least $n/2 - k$. The running time is, in both cases, asymptotically optimal, under the exponential-time hypothesis (ETH).

The results extend the range of tractability of the Hamiltonian cycle problem, showing that it is fixed-parameter tractable when parameterized below a natural bound. In addition, for the first parameterization we show that a kernel with $O(k)$ vertices can be found in polynomial time.

2012 ACM Subject Classification Theory of computation \rightarrow Graph algorithms analysis, Theory of computation \rightarrow Parameterized complexity and exact algorithms

Keywords and phrases Hamiltonian cycle, fixed-parameter tractability, kernelization.

1 Introduction

The HAMILTONIAN CYCLE problem asks whether a given undirected graph has a cycle that visits each vertex exactly once. It is a central problem of graph theory, operations research, and computer science, with an early history that well predates these fields (see e.g. [27]).

Several conditions that guarantee the existence of a Hamiltonian cycle in a graph are known. Perhaps best known among these is Dirac's theorem from 1952 [14]. It states that a graph with n vertices ($n \geq 3$) is Hamiltonian if every vertex has degree at least $n/2$. Various extensions and refinements of Dirac's theorem have been obtained, often involving further graph parameters besides minimum degree (see e.g. the book chapters [13, § 10], [29, § 11] and survey articles [17, 28, 30] for an overview). We remark that a polynomial-time verifiable condition for Hamiltonicity cannot be both necessary and sufficient, unless $P = NP$ [25]. In its stated form, Dirac's theorem is as strong as possible. In particular, if we replace $n/2$ by

¹ Supported by NWO Gravitation grant "Networks".

² Supported by ERC Consolidator Grant No 617951.

³ Supported by NWO Gravitation grant "Networks" and NWO Grant No 639.021.438.

$\lfloor n/2 \rfloor$, the graph may fail to be two-connected—a precondition for Hamiltonicity. (Consider two $\lceil n/2 \rceil$ -cliques with a common vertex.)

In this paper we relax the conditions of Dirac's theorem and consider input graphs in which (1) at least $n - k$ vertices have degree at least $n/2$ (the degrees of the remaining vertices can be arbitrarily small), or (2) all vertices have degree at least $n/2 - k$.

For both relaxations we show that HAMILTONIAN CYCLE can be solved deterministically, in time $c^k \cdot n^{O(1)}$, for some fixed constant c . This establishes the fixed-parameter tractability of HAMILTONIAN CYCLE when parameterized by the distance from Dirac's bound, for two natural ways of measuring this distance.

The known exact algorithms for HAMILTONIAN CYCLE in general graphs have exponential running time (the problem is one of the original 21 NP-hard problems [25]). The best deterministic running time of $O(2^n \cdot n^2)$ is achieved by the dynamic programming algorithm of Bellman [4], and Held and Karp [23], and has not been improved since the 1960s. Among randomized algorithms, the current best running time of $O(1.657^n)$ is achieved by the more recent algorithm of Björklund [6] based on determinants. Improving these bounds remains a central open question of the field.

Assuming the exponential-time hypothesis (ETH) [24], there is no algorithm for HAMILTONIAN CYCLE with running time $2^{o(n)}$. In both parameterizations considered in this paper, $k \leq n$ holds. Thus, under ETH, a running time of the form $2^{o(k)} \cdot n^{O(1)}$ is ruled out, and our algorithms are optimal, up to the base of the exponential. Furthermore, there exists a fixed constant $\alpha > 0$, such that our parameterized bounds asymptotically improve the current best bounds for HAMILTONIAN CYCLE, if the value of k is at most $\alpha \cdot n$.

For the first parameterization, we show that HAMILTONIAN CYCLE admits a kernel with $O(k)$ vertices, computable in polynomial time. In other words, the input graph can be compressed (roughly) to the order of its sparse part, while preserving Hamiltonicity.

Our results show that checking Hamiltonicity becomes tractable as we approach the degree-bound of Dirac's theorem. The crude intuition behind Dirac's theorem (and many of its generalizations) is that *having many edges* makes a graph Hamiltonian. It is a priori far less obvious why approaching the Dirac bound would make the *algorithmic problem* easier; one may even expect that the more edges there are, the harder it becomes to certify *non-Hamiltonicity*. To provide some intuition why this is not the case, we give a brief informal summary of the arguments.

When $n - k$ vertices have degree at least $n/2$, i.e. in the first case, our algorithm takes advantage of the fact that, by a result of Bondy and Chvátal, the subgraph induced by the high-degree vertices can be completed to a clique without changing the Hamiltonicity of the graph; all relevant structure is thus in the sparse part and its interconnection with the dense part. Then, we find a subset of the vertices in the clique that are well-connected to the sparse part (by solving a matching problem in an auxiliary graph), and we ignore the remainder of the clique. Finally, we show how a Hamiltonian cycle on this smaller, well-connected subgraph, can be extended to a Hamiltonian cycle of the entire graph, guided by the alternating paths of the matching. For this parameterization we are not aware of a comparable result in the literature.

When all vertices have degree at least $n/2 - k$, i.e. in the second case, a result of Nash-Williams implies that either a Hamiltonian cycle, or a sufficiently large independent set can be found in polynomial time. In the latter case, we certify non-Hamiltonicity by showing (roughly) that the complement of the independent set is not coverable by a certain number of disjoint paths. This argument is essentially the same as the one given by Häggkvist [22] towards his algorithm with running time $O(n^{5k})$ for the same parameterization. (Häggkvist

states this algorithmic result as a corollary of structural theorems. He does not describe the details of the algorithm or its analysis, but these are not hard to reconstruct.) Here we improve the running time of Häggkvist’s algorithm to the stated (asymptotically optimal) $c^k \cdot n^{O(1)}$ by more efficiently solving the arising path-cover subproblem.

1.1 Statement of results

Our first result shows that if a graph has a “relaxed” Dirac property, it can be compressed while preserving its Hamiltonicity.

► **Theorem 1.1.** *Let G be an n -vertex graph such that at least $n - k$ vertices of G have degree at least $n/2$. There is a deterministic algorithm that, given G , constructs in time $O(n^3)$ a $3k$ -vertex graph G' , such that G is Hamiltonian if and only if G' is Hamiltonian.*

Equivalently stated in the language of parameterized complexity, the Hamiltonian cycle problem parameterized by k has a *kernel* with a linear number of vertices. To determine the Hamiltonicity of a graph G , we simply apply the algorithm of Theorem 1.1 to compress G , and use an exponential-time algorithm (for instance, the Held-Karp algorithm) to solve HAMILTONIAN CYCLE directly on the compressed graph. We thus obtain the following result.

► **Corollary 1.2.** *If at least $n - k$ vertices of an n -vertex graph G have degree at least $n/2$, then HAMILTONIAN CYCLE with input G can be solved in deterministic time $O(8^k \cdot k^2 + n^3)$.*

As an alternative, we may also use an approach based on inclusion-exclusion [26] to solve the reduced HAMILTONIAN CYCLE instance, achieving the overall running time $O(8^k \cdot k^3 + n^3)$, with *polynomial space*.

Our result for the second relaxation of Dirac’s theorem is as follows.

► **Theorem 1.3.** *If every vertex of an n -vertex graph G has degree at least $n/2 - k$, then HAMILTONIAN CYCLE with input G can be solved in deterministic time $O(30^{6k} \cdot n^3)$.*

The running time of the Bellman-Held-Karp algorithm for HAMILTONIAN CYCLE is $O(2^n \cdot n^2)$. Denoting $\alpha = k/n$, our results represent an asymptotic improvement if $\alpha < 1/3$ in the first parameterization, and if $\alpha < 0.0339$ in the second parameterization.

As a counterpoint to our results, we mention that HAMILTONIAN CYCLE remains hard (in both parameterizations) for arbitrarily small values of α .

► **Theorem 1.4.** *Assuming ETH, HAMILTONIAN CYCLE cannot be solved in time $2^{o(n)}$ in n -vertex graphs with at least $(1 - \alpha) \cdot n$ vertices of degree at least $n/2$, and in n -vertex graphs with minimum degree $(1 - \alpha) \cdot n/2$, for arbitrary fixed $0 < \alpha < 1/2$.*

Proof. In both cases we construct a graph with the given degree-requirements that embeds a hard instance of HAMILTONIAN PATH with $\alpha \cdot n$ vertices. For the second statement we can use the construction from the NP-hardness proof of Dahlhaus, Hajnal, and Karpinski [12]. For the first statement, consider an $\alpha \cdot n$ -vertex instance of HAMILTONIAN PATH, connected by two disjoint edges to an $(1 - \alpha) \cdot n$ -vertex clique. ■

1.2 Related work

In general, parameterized complexity [11, 16] allows a finer-grained understanding of algorithmic problems than classical, univariate complexity. No new insight is gained, however, if the chosen parameter k is large in all interesting cases. For example, in planar graphs, the Four Color Theorem guarantees the existence of an independent set of size $n/4$. As a

consequence, any exponential-time algorithm for maximum independent set trivially achieves fixed-parameter tractability in terms of the solution size.

To deal with this issue, Mahajan and Raman [31] introduced the method of parameterizing problems *above* or *below* a guaranteed bound. (Similar considerations motivate the “distance from triviality” framework of Guo, Hüffner, and Niedermeier [18].) In the example of planar independent set, an interesting parameter is the amount by which the solution size exceeds $n/4$. Similar ideas have successfully been applied to several problems (see e.g. [2, 5, 10, 19, 20, 32]). Our results also fall in the framework of “above/below” parameterization, with the remark that our parameter of interest is not the value to be optimized but a structural property of the input, which we parameterize near its “critical value”.

Perhaps closest to our work is the recent result of Gutin and Patel [21] on the Traveling Salesman problem, parameterized below the cost of the *average* tour. Although it concerns Hamiltonian cycles (in an edge-weighted complete graph), the result of Gutin and Patel is not directly comparable with our results. In particular, averaging arguments do not seem to help when studying the *existence* of Hamiltonian cycles, which is often determined by local structure in the graph. For instance, HAMILTONIAN CYCLE remains NP-hard even in graphs with average degree αn for any constant $\alpha < 1$. (Consider a clique of $\sqrt{\alpha n}$ vertices, connected by two non-incident edges to the remaining graph that encodes a hard instance of HAMILTONIAN PATH.)

2 Preliminaries

We use standard graph-theoretic notation (see e.g. [13]). An edge between vertices u and v is written simply as uv or vu . The *neighborhood* of a vertex v in graph G is denoted by $N_G(x)$. The *degree* of v in G is $d_G(v) = |N_G(v)|$, and the minimum degree of G is $\delta_G = \min_{v \in V(G)} d_G(v)$. We conveniently omit the subscript G whenever possible. For a set $S \subseteq V(G)$ of vertices, $G[S]$ denotes the subgraph induced by S on G .

We state Dirac's theorem and a strengthened statement due to Ore. Let G be an n -vertex undirected graph, with $n \geq 3$.

► **Lemma 2.1** (Dirac [14]). *If $\delta \geq n/2$, then G is Hamiltonian.*

► **Lemma 2.2** (Ore [34]). *If $d(u) + d(v) \geq n$ for every non-adjacent pair of vertices u, v of G , then G is Hamiltonian.*

We state a theorem of Bondy and Chvátal that we use in the proofs of both Theorem 1.1 and Theorem 1.3.

► **Lemma 2.3** (Bondy-Chvátal [7]). *Let G be an n -vertex graph, and let G' be obtained from G by adding an edge uv to G for some pair of non-adjacent vertices u, v such that $d_G(u) + d_G(v) \geq n$. Then G' is Hamiltonian if and only if G is Hamiltonian. Moreover, given a Hamiltonian cycle of G' , a Hamiltonian cycle of G can be obtained in linear time.*

It is easy to see that Lemma 2.3 implies both Lemma 2.1 and Lemma 2.2, as in both cases we can iterate the edge-augmentation step until obtaining a complete graph.

Finally, we state yet another strengthening of Dirac's theorem, due to Nash-Williams [33]. We write this result in a slightly non-standard, explicitly algorithmic form. Our use of this result in proving Theorem 1.3 is the same as in the argument of Häggkvist [22].

► **Lemma 2.4** (Nash-Williams [33]). *Let G be a 2-connected graph with n vertices, with $\delta \geq (n+2)/3$. Then, we can find in G , in time $O(n^3)$, either a Hamiltonian cycle, or an independent set of size $\delta + 1$.*

The following proof of Lemma 2.4 is due to Bondy [8], sketched in [29, §11]. We spell it out fully to make our discussion self-contained and to provide an explicitly algorithmic form (this requires only minor changes compared to the presentation in [29]).

Proof. All cycles considered in the proof are simple. Denote $V = V(G)$. Start with an arbitrary cycle C of G (the fact that G is 2-connected guarantees the existence of a cycle, and we can easily find one in linear time). We extend C into successively longer cycles until we either (1) reach a Hamiltonian cycle, or (2) find an independent set of the required size.

Unless we have already found a Hamiltonian cycle, $|C| \leq n - 1$ holds. Suppose $V \setminus C$ is an independent set, and let $v \in V \setminus C$ be an arbitrary vertex. Due to the independence of $V \setminus C$, we have $N(v) \subseteq C$. If two neighbors of v are connected by an edge of C , then we can immediately extend C via v . Assume therefore, that this is not the case. Fix an arbitrary orientation of C , and let N^+ be the set of successors in C of the vertices $N(v)$. Then, $|N^+| \geq \delta$. Again, if two vertices $x, y \in N^+$ are connected, then C can be further extended (see Figure 1(a) for illustration). Thus, we can assume that $N^+ \cup \{v\}$ is an independent set of G , of the required size.

It remains to show that C can be extended whenever $V \setminus C$ is *not* an independent set. Then, $|V \setminus C| \geq 2$ must hold. Let us orient C arbitrarily and label its vertices accordingly as x_1, \dots, x_k . Let $P = (x_1, p_1, p_2, \dots, x_{t+1})$, with $1 \leq t < k$, be a simple path of length at least 3, that intersects C only at the endpoints x_1 and x_{t+1} (see Figure 1(c)). We claim that the existence of such a path can be assumed without loss of generality (by suitably choosing the starting label x_1).

To see this, consider a path (u, v, w) in G , where $u \in C$ and $v, w \in V \setminus C$. (Such a path must exist by the assumption that $V \setminus C$ is not independent and the fact that G is 2-connected: start with an arbitrary edge outside C and consider a path from one of its endpoints to a vertex of C).

If there is a path, vertex-disjoint from $\{u, v\}$, from w to an arbitrary vertex $x \in C \setminus \{u\}$, then we obtain the desired structure by labeling $x_1 = u$, $p_1 = v$, $p_2 = w$, and $x_{t+1} = x$.

If there is no such path, then there must be a path R , internally vertex-disjoint from $\{u, v\}$, from v to an arbitrary vertex $x \in C \setminus \{u\}$ (otherwise, deleting u would disconnect G , contradicting its 2-connectivity). Furthermore, there must exist a path Q connecting u to w , not containing v (otherwise, deleting v would separate u and w) and internally vertex-disjoint from C and R (otherwise, we would have a path from w to a vertex in $C \setminus \{u\}$, ruled out previously). Now, we obtain the desired structure by setting $x_1 = u$, $x_{t+1} = x$, and p_1 and p_2 the second and third vertices on the path consisting of Q , and the edge wv (Figure 1(b)).

Let $d_t^\uparrow, d_k^\uparrow, d_2^\uparrow$ denote the number of neighbors x_i with $1 \leq i \leq t$ of x_t, x_k , resp. p_2 . Similarly, let $d_t^\downarrow, d_k^\downarrow, d_2^\downarrow$ denote the number of neighbors x_i with $t < i \leq k$ of x_t, x_k , resp. p_2 . Let $d_t^\circ, d_k^\circ, d_2^\circ$ denote the number of neighbors in $V \setminus C$ of x_t, x_k , resp. p_2 .

► **Claim 2.5.** *At least one of the following three inequalities holds:*

- (1) $d_t^\uparrow + d_k^\uparrow + d_2^\uparrow > t + 1$,
- (2) $d_t^\downarrow + d_k^\downarrow + d_2^\downarrow > k - t + 1$.
- (3) $d_t^\circ + d_k^\circ + d_2^\circ > n - k - 1$

Proof. Suppose otherwise. Then the sum of degrees of x_t, x_k, p_2 is at most $(t + 1) + (k - t + 1) + (n - k - 1) \leq n + 1$, and thus at least one of them has degree at most $(n + 1)/3 < \delta$, a contradiction. ◻

In the following, we assume $d_2^\uparrow \leq 1$, since, if p_2 is connected to some x_i with $1 < i \leq t$, then we may choose a different index t in our construction.

Suppose inequality (1) holds. Then, $d_t^\uparrow + d_k^\uparrow > t$, and by the pigeonhole-principle, there is some $1 \leq i < t$ such that $x_k x_{i+1}$ and $x_t x_i$ are edges of G . Then, C can be extended by adding these two edges and the path $(x_1, p_1, \dots, x_{t+1})$ and removing the edges $x_k x_1$, $x_t x_{t+1}$, and $x_i x_{i+1}$. (See Figure 1(c).)

Suppose inequality (2) holds. Then, there is some $t+1 \leq i < k$ such that one of the following is true: (a) $x_k x_i$ and $x_t x_{i+1}$ are edges of G , (b) $x_t x_i$ and $p_2 x_{i+1}$ are edges of G , (c) $i < k-1$, and $x_k x_i$ and $p_2 x_{i+2}$ are edges of G , (d) $x_k x_t$ or $p_2 x_k$ is an edge of G . In all these cases C can be extended similarly to the previous case. (See Figure 1(d) for an illustration of cases (a), (b), (c).)

To see that one of the four cases must hold, we can argue by contradiction. Apart from the boundaries, every vertex x_i that is connected to x_t rules out x_{i-1} from being connected to x_k . Similarly, every vertex x_i that is connected to p_2 rules out x_{i-2} from being connected to x_k , as well as x_{i-1} from being connected to x_t . Thus, fixing the connections from x_t and p_2 first, we rule out $d_t^\downarrow + d_k^\downarrow - 2$ possible neighbors of x_k .

Suppose inequality (3) holds. Then, by a similar pigeonhole-argument, one of the following is true: (a) $x_t p_j$ or $x_k p_j$ is an edge of G for some internal vertex p_j on path P , or (b) at least two of $x_t w$, $x_k w$, and $p_2 w$ are edges of G , for some $w \in V \setminus (C \cup P)$. In all these cases C can be extended similarly to the previous cases.

The claimed running time can be achieved via a straightforward implementation. \blacksquare

3 Relaxing the cardinality-constraint (proof of Theorem 1.1)

Let $C \subseteq V(G)$ denote the set of high-degree vertices of G (those with degree at least $n/2$), and let $S = V(G) \setminus C$ denote the remaining (i.e. low-degree) vertices.

Observe that $|S| \leq k$. By Lemma 2.3, we may add all edges between vertices in C , without changing the Hamiltonicity of G , assume therefore that C is a clique.

The proof of the following theorem is inspired by the *crown reductions* [1, 9, 15] used to obtain kernels for VERTEX COVER and SAVING k COLORS.


► **Theorem 3.1.** *There is a polynomial-time algorithm that, given a graph G and a nonempty set $S \subseteq V(G)$ such that $G - S$ is a clique, outputs an induced subgraph G' of G on at most $3|S|$ vertices such that G is Hamiltonian if and only if G' is Hamiltonian.*

Proof. Given a graph G let $S \subseteq V$, such that $C := V(G) \setminus S$ is the vertex set of a clique in G . If $C \leq 2|S|$ then $G' := G$ suffices, so we assume $C > 2|S|$ in the remainder. Let $S' := \{v_1, v_2 \mid v \in S\}$ be a set containing two representatives for each vertex of S . Construct a bipartite graph H on vertex set $C \cup S'$. For each edge $cv \in E(G)$ with $c \in C$ and $v \in S$, add the edges cv_1, cv_2 to H . Compute a maximum matching $M \subseteq E(H)$ in graph H , for example using the Edmonds-Karp algorithm. Let C^* be the vertices of C saturated (matched) by M . If $|C^*| \geq |S| + 1$ then let $C' := C^*$, and otherwise let $C' \subseteq C$ be a superset of C^* of size $|S| + 1$. Output the graph $G' := G[C' \cup S]$ as the result of the reduction.

► **Claim 3.2.** *Graph G' has at most $3|S|$ vertices.*

Proof. Since each vertex of C^* is matched to a distinct vertex in S' , with $|S'| = 2|S|$, it follows that $|C^*| \leq 2|S|$ which implies $|C'| \leq 2|S|$. As $V(G') = C' \cup S$, the claim follows. \square

The output graph G' therefore satisfies the size bound. It remains to prove that it is equivalent to G with respect to Hamiltonicity. We first prove the simpler implication.



■ **Figure 1** Illustration of the proof of Lemma 2.4. (a) cycle C with sets $N(v)$ and N^+ of vertices shown as solid, resp. hollow circles; (b) constructing a path between two vertices of C ; (c) one case of extending C ; (d) extending C in the first three cases, when inequality (2) holds. Vertices x_t, x_k, p_2 shown as hollow dots.

► **Claim 3.3.** *If G' is Hamiltonian, then G is Hamiltonian.*

Proof. Suppose that G' is Hamiltonian, and let $F \subseteq E(G)$ be a Hamiltonian cycle in G' . Fix an arbitrary orientation of F . As each vertex from C' has a unique successor on F , while $|C'| > |S|$ by definition, it follows that some vertex $x \in C'$ has a successor from C' along the cycle; let this be $y \in C'$. Then we can transform F into a Hamiltonian cycle in G by removing the edge xy and replacing it by a path through all the clique-vertices of $C \setminus C'$. \square

The remainder of the proof is aimed at proving the reverse implication. For this, we introduce some terminology. For a vertex set S^* in a graph G^* , we define a *path cover* of S^* in G^* as a set of pairwise vertex-disjoint simple paths P_1, \dots, P_ℓ in G^* , such that each vertex of S^* belongs to exactly one path P_i . For a vertex set C^* in G^* , we say the path cover *has*

C^* -endpoints if the endpoints of each path P_i belong to C^* . We will sometimes interpret a subgraph in which each connected component is a path as a path cover, in the natural way.

► **Claim 3.4.** *If there is a path cover of S in G' having C' -endpoints, then G' is Hamiltonian.*

Proof. Any path cover of S consists of at least one path (since S is nonempty by assumption) and the endpoints of the paths are all distinct. Hence a path cover consisting of $\ell \geq 1$ paths has exactly 2ℓ distinct endpoints $\{s_1, t_1, \dots, s_\ell, t_\ell\}$, which are vertices in the clique C' . Let $P_{\ell+1}$ be a simple path in G' visiting all vertices that are not touched by the path cover; such a path exists because the only vertices not touched by the path cover belong to the clique C' . Then one can obtain a Hamiltonian cycle in G' by taking the edges of $P_1, \dots, P_\ell, P_{\ell+1}$, together with edges connecting the end of path P_i to the beginning of path P_{i+1} for all relevant values of i . ◻

To prove that Hamiltonicity of G implies Hamiltonicity of G' , we will construct a path cover of S in G' having C' -endpoints, using a hypothetical Hamiltonian cycle in G . To do so we need several properties enforced by the matching M in H , which we now explore.

Let U_C be the vertices of C that are not saturated by M . Let R denote the vertices of H that are reachable from U_C by an M -alternating path in the bipartite graph H (which necessarily starts with a non-matching edge), and define $R_C := R \cap C$ and $R_{S'} := R \cap S'$.

► **Claim 3.5.** *The sets $R, R_C, R_{S'}$ satisfy the following.*

1. *Each M -alternating path in H from U_C to a vertex in $R_{S'}$ (resp. R_C) ends with a non-matching (resp. matching) edge.*
2. *Each vertex of $R_{S'}$ is matched by M to a vertex in R_C .*
3. *For each vertex $x \in R_C$ we have $N_H(x) \subseteq R_{S'}$.*
4. *For each vertex $v \in S$ we have $v_1 \in R_{S'} \Leftrightarrow v_2 \in R_{S'}$.*
5. *For each vertex $v \in S' \setminus R_{S'}$, we have $N_H(v) \cap R_C = \emptyset$ and each vertex of $N_H(v)$ is saturated by M .*

Proof. (1) An M -alternating path starting in U_C must start with a non-matching edge, since U_C consists of unsaturated vertices, and it starts from the C -partite set of H . Hence such a path moves to the S' -partite set over non-matching edges, and moves back to the C -partite set over matching edges.

(2) If a vertex $x \in R_{S'} \subseteq R$ is not saturated, then the M -alternating path from U_C witnessing $x \in R$ starts and ends with a non-matching edge (by (1)) and is in fact an M -augmenting path. This contradicts that M is a maximum matching. Hence each $x \in R_{S'}$ is matched by M to some vertex y . By (1) the M -alternating path from U_C to x that witnesses $x \in R_{S'}$ ends with a non-matching edge, so together with the matching edge $\{x, y\}$ this forms an M -alternating path witnessing $y \in R_C$.

(3) Consider a vertex $x \in R_C$ and an M -alternating path P from U_C witnessing $x \in R$. By (1) the last edge on P (if any) is a matching edge. Hence if x is saturated by M , then its matching partner y is the predecessor of x on P and a prefix of P witnesses $y \in R$ and hence $y \in R_{S'}$. For any vertex $z \in N_H(x)$ that is not the matching partner of x , we can augment P by the edge xz to obtain an M -alternating path from U_C to z witnessing $z \in R_{S'}$. Together, these two arguments show $N_H(x) \subseteq R_{S'}$.

(4) Suppose $v_1 \in R_{S'}$ and let P be an M -alternating path from U_C to v_1 . By (1) path P ends with a non-matching edge xv_1 . Since v_1 and v_2 have identical neighborhoods in H , we can replace the last edge of P by xv_2 to obtain an M -alternating path witnessing $v_2 \in R_{S'}$. The case that $v_2 \in R_{S'}$ is symmetric.

(5) Consider $v \in S' \setminus R_{S'}$. If $x \in N_H(v) \cap R_C$, then (3) implies $v \in R_{S'}$, a contradiction. Hence $N_H(v) \cap R_C = \emptyset$. An unsaturated H -neighbor x of v would imply $x \in N_H(v) \cap U_C \subseteq N_H(v) \cap R_C$, so each vertex of $N_H(v)$ is saturated by M . \square

Using these structural insights we can now prove the desired converse to Claim 3.3. Before we give the formal proof, we present the main idea. To prove that G' is Hamiltonian if G is, we take a Hamiltonian cycle F in G and turn it into a path cover of S in G' with C' -endpoints. Any Hamiltonian cycle F in G yields a path cover of S with S -endpoints, by simply taking the restriction of F onto the vertices of S . The challenge is to extend this path cover with edges into C' to give it the desired C' -endpoints: if the Hamiltonian cycle F used an edge to jump from S to C , we have to provide a similar jump in G' . If F jumps from a vertex $v \in S$ whose corresponding copies $v_1, v_2 \in S'$ do not belong to $R_{S'}$, then the C -endpoint of the jumping edge is saturated by M , belongs to C' and therefore to G' , and can be used to provide the analogous jump in G' . On the other hand, for all vertices $v \in S$ whose copies v_1, v_2 belong to $R_{S'}$, we will globally assign new jumping edges based on the matching H . The properties of a matching will ensure that these jumping edges lead to distinct targets and give a valid path cover of S in G' having C' -endpoints. We now formalize these ideas.

► **Claim 3.6.** *If G is Hamiltonian, then G' is Hamiltonian.*

Proof. Let F be a Hamiltonian cycle in G . By Claim 3.4 it suffices to build a path cover of S in G' with C' -endpoints. View F as a 2-regular subgraph of G , and let $F_1 := F[S]$ be the subgraph of F induced by S . Since F spans G and all vertices of S are present in G' , it follows that F_1 is a path cover of S in G' . However, the paths in F_1 have their endpoints in S rather than in C' . We resolve this issue by inserting edges into F_1 to turn it into an acyclic subgraph F_2 of G' in which each vertex of S has degree exactly two. This structure F_2 must be a path cover of S in G' with C' -endpoints, since the degree-two vertices S cannot be endpoints of the paths. To do the augmentation, initialize F_2 as a copy of F_1 . Define $R_S := \{v \in S \mid v_1 \in R_{S'} \vee v_2 \in R_{S'}\}$ and proceed as follows.

- For each vertex $v \in R_S$, we have $v_1, v_2 \in R_{S'}$ by Claim 3.5(4), which implies by Claim 3.5(2) that both v_1 and v_2 are matched to distinct vertices $x_1, x_2 \in R_C$. If v has degree zero in subgraph F_1 , then add the edges vx_1, vx_2 to F_2 . If v has degree one in F_2 then only add the edge vx_1 . Do not add any edges if v already has degree two in F_1 .
- For each vertex $v \in S \setminus R_S$, we claim that $N_G(v) \cap R_C = \emptyset$. This follows from the fact that $N_G(v) = N_H(v_1) = N_H(v_2)$ and Claim 3.5(5), using that $v \notin R_S$ implies $v_1, v_2 \notin R_{S'}$. Hence the (up to two) neighbors that $v \in S \setminus R_S$ has in C on the Hamiltonian cycle F do not belong to R_C , while Claim 3.5(5) ensures that all vertices of $N_G(v)$ are saturated by H and hence belong to C' . For each vertex $v \in S \setminus R_S$, for each edge from v to $C \cap C'$ incident on v in F , we insert the corresponding edge into F_2 .

It is clear that the above procedure produces a subgraph F_2 in which all vertices of S have degree exactly two. To see that F_2 is indeed a path cover, having no vertex of degree larger than two, it suffices to notice that the edges inserted for $v \in R_S$ connect to *distinct* vertices in $C' \cap R_C$, while the edges inserted for $v \in S \setminus R_S$ connect to $C' \setminus R_C$ in the same way as in the Hamiltonian cycle F . Hence F_2 forms a path cover of S in G' having C' -endpoints, which implies that G' is Hamiltonian and proves Claim 3.6. \square

Claims 3.3 and 3.6 prove the correctness of the reduction and Claim 3.2 gives the desired size bound. Since the reduction can easily be performed in polynomial time, this completes the proof of Theorem 3.1. ■

Observe that the proof of Lemma 3.3 explicitly constructs the Hamiltonian cycle in case of a “yes”-answer. The running time of the reduction is dominated by the bipartite matching step, and the process of undoing the Bondy-Chvátal augmentations (Lemma 2.3), if a cycle of the original graph is to be constructed. Both tasks can be performed in time $O(n^3)$.

4 Relaxing the degree-constraint (proof of Theorem 1.3)

The outline of the proof largely follows an earlier argument of Häggkvist [22]. We improve the $O(n^{5k})$ running time of Häggkvist's algorithm to $c^k \cdot n^{O(1)}$.

The algorithm either finds a Hamiltonian cycle or constructs a certificate of non-Hamiltonicity, in the form of a cut (S, T) of the graph, such that the vertices of T can not be covered by $|S|$ vertex-disjoint paths, and this certificate can be verified within the required running time. (Observe that a Hamiltonian cycle induces such a path-cover for an arbitrary cut; paths consisting of single vertices are allowed.)

Assume that $k < n/34$, and thus $\delta > 8n/17$. (Otherwise we revert to a standard exponential-time algorithm.) Furthermore, $\delta < n/2$ may be assumed, as otherwise G is Hamiltonian by Dirac's theorem. Also assume that G is 2-connected (otherwise it is not Hamiltonian).

Start by running the procedure from the proof of Lemma 2.4, either obtaining a Hamiltonian cycle, or an independent set of size $\delta + 1$. Assume that the latter is the case, and label the obtained independent set as A_1 .

Partition $V(G)$ into sets A_1 , A_2 , and A_3 , where A_2 denotes the set of vertices in $v \in V(G) \setminus A_1$ such that $|N(v) \cap A_1| \geq \delta/2$, and $A_3 = V(G) \setminus (A_1 \cup A_2)$. In words, A_2 contains vertices that are sufficiently highly connected to the obtained independent set, and A_3 contains the remaining vertices.

► **Lemma 4.1** ([22], Thm. 2). *Given sets A_1, A_2, A_3 as defined, we can find a set of vertices $S \subseteq V(G)$ such that $|S| \geq 3\delta - n + 2$, and $G[V(G) \setminus S]$ can be covered by $|S|$ vertex-disjoint paths if and only if G is Hamiltonian.*

We sketch the argument, referring to Häggkvist [22, p. 32-33] for the full details.

Proof sketch. Let $T = V(G) \setminus S$. The “if” direction is trivial, since every Hamiltonian cycle of G induces a cover of $G[T]$ by $|S|$ vertex-disjoint paths, for arbitrary S .

It remains to show the converse, i.e. that a cover of $G[T]$ by $|S|$ vertex-disjoint paths can be extended into a Hamiltonian cycle of G , for a suitably chosen S .

We let $S = A_2$ if $|A_2| \leq |A_1 \cup A_3|$, and $S = A_1 \cup A_3$ otherwise.

Consider the bipartite graph with sides S and T containing only those edges of G that have one endpoint in S and another endpoint in T . Add to this bipartite graph the edges of the path-cover of T by $|S|$ vertex-disjoint paths (assuming such a cover exists). Call this graph B . Let B' be the graph obtained from B by connecting all pairs of vertices in S .

Observe that B is Hamiltonian if and only if B' is Hamiltonian (furthermore, in case they are Hamiltonian, they admit the exact same Hamiltonian cycles). The “only if” case is obvious since $E(B') \supseteq E(B)$. In the other direction, $B'[T]$ contains $|S|$ disjoint components, therefore a Hamiltonian cycle of B' can not traverse any edge of $B'[S]$. (This is because for an arbitrary Hamiltonian cycle H of B' the induced graphs $H[S]$ and $H[T]$ have the same number of connected components.)

Now we form graph B'' from B' , by repeatedly applying the Bondy-Chvátal theorem (Lemma 2.3), adding edges with one endpoint in S and one endpoint in T . It can be shown that this results in adding all edges between S and T .

As B'' fully connects S and T , it is clearly Hamiltonian, as we can traverse the $|S|$ vertex-disjoint paths in $B''[T]$ and link them together through hops via arbitrary vertices of S (using each vertex in S exactly once).

The claim on the size of S and the fact that all possible edges between S and T satisfy the degree-conditions of the Bondy-Chvátal Theorem (Lemma 2.3) follow from a delicate counting argument, which crucially relies on the fact that $8n/17 < \delta < n/2$. We refer to [22] for the details. ■

It remains to verify whether $G[T]$ can be covered by $|S|$ vertex-disjoint paths.

► **Lemma 4.2.** *Given an n -vertex graph G , we can find in time $O(c^t \cdot n^3)$ a cover of G with $n - t$ vertex-disjoint paths, or report that no such cover exists, for arbitrary $c > (2e)^2$.*

Proof. Apply color-coding [3], [11, §5.2]. Call a path nontrivial if it has more than one vertex. Say that a coloring is *good* for a cover by vertex-disjoint paths, if all vertices that appear in a nontrivial path receive a different color. Clearly, if there is a cover by at most $n - t$ paths then there is a cover by exactly $n - t$ paths, and in such a cover there are at most $2t$ vertices that appear in a nontrivial path. So if there is a path cover with $n - t$ paths, a random coloring with $2t$ colors is *good* for this cover with probability e^{-2t} . (See e.g. [11, Lemma 5.4].)

On a vertex-colored graph with color set $C = \{1, \dots, 2t\}$, we solve the following problem by dynamic programming: for a set $X \subseteq C$ and $v \in V(G)$, let $T[X, v]$ be the smallest number q for which there exists a collection P_1, \dots, P_q of vertex-disjoint paths in G , such that P_q ends in vertex v and the multiset of colors used in P_1, \dots, P_q is exactly equal to X . (In particular, this implies that no two vertices in a path may have the same color, for it would appear twice in the multiset and only once in the set X .)

Let the $2t$ -coloring of G be given by $f : V(G) \rightarrow [2t]$. Then $T[X, v]$ satisfies the following recurrence:

- $T[\{c\}, v] = 1$ if $f(v) = c$,
- $T[X, v] = +\infty$ if $f(v) \notin X$,
- $T[X, v] = \min \left\{ 1 + \min_{u \in V(G) \setminus \{v\}} T[X \setminus \{f(v)\}, u], \min_{u \in N_G(v)} T[X \setminus \{f(v)\}, u] \right\}$, otherwise.

Intuitively, the interesting part of the recurrence has two cases: either we can let v be a trivial path (so we pay 1 for having a path with v , and then need a collection of paths that can end at any other vertex u that covers the remaining colors), or we take a system of paths covering the remaining colors that ends in a neighbor u of v , and add the edge uv to the end of that path.

Now, observe that for any color-subset X and vertex v , there is a cover of G with $T[X, v] + (n - |X|)$ paths: we cover $|X|$ vertices, one of each color in X , by $T[X, v]$ paths and cover the remaining $(n - |X|)$ vertices by trivial paths. So if we encounter a set X and vertex v for which $T[X, v] + (n - |X|) \leq n - t$, or equivalently, $T[X, v] \leq |X| - t$, then the answer is “yes”. On the other hand, if G has a path cover by $n - t$ paths and a coloring is good for this cover, then letting X be the set of colors of vertices that appear in a nontrivial path and v an endpoint of such a path, we obtain $T[X, v] + (n - |X|) \leq n - t$.

So by trying e^{2t} random colorings and solving the dynamic program for each one, we solve the “cover by $n - t$ disjoint paths” problem with constant success probability. With c^{2t} for $c > e$, we can boost the success probability arbitrarily close to 1. The dynamic program can be solved in time $O(2^{2t} n^3)$. The claimed running time follows.

We may de-randomize the algorithm by replacing the randomized coloring by a deterministic construction, e.g. via *splitters*. We omit the details of this, by now standard, technique [11, §5.6] ■

In our application of Lemma 4.2, we need to cover $G[T]$ by $|S|$ vertex-disjoint paths. Observe that $|S| \geq n/2 - 3k$, and consequently $|T| \leq n/2 + 3k$. The difference between the order of the graph $G[T]$ and the number of paths t with which we want to cover it, is therefore at most $6k$.

Applying Lemma 4.2, the running time of this step is thus $O(c^{6k} \cdot n^3)$, for arbitrary $c > (2e)^2$.

To construct a Hamiltonian cycle, find the set S using Lemma 2.4 and Lemma 4.1, find an appropriate path-cover using Lemma 4.2, and recover the Hamiltonian cycle of G by undoing the Bondy-Chvátal steps in Lemma 4.1. The claimed running time of Theorem 1.3 follows by adding up the corresponding terms and by using straightforward data structuring.

5 Remarks and open questions

We described two algorithms that solve the Hamiltonian cycle problem, with running time that depends polynomially on the graph size and single-exponentially on the distance from Dirac's bound, a condition that guarantees the Hamiltonicity of a graph. We have considered two different ways of measuring this distance. It would be interesting to improve the bases of the exponentials in our running times, and to obtain a polynomial kernel for the second parameterization.

A natural question left open by our work is whether the two parameterizations can be combined, to obtain a generalization of both. We suspect but have not been able to prove that the following holds.

► **Conjecture 5.1.** *If at least $n - k$ vertices of G have degree at least $n/2 - k$, then HAMILTONIAN CYCLE with input G can be solved in time $c^k \cdot n^{O(1)}$ for some constant c .*

The results of this paper can be extended with minimal changes to similar parameterizations of Ore's theorem (Lemma 2.2). Extending the results to generalizations of Dirac's and Ore's theorems to *digraphs* would be interesting. More generally, finding new algorithms by parameterizing structural results of graph theory (whether related to Hamiltonicity or not) is a promising direction.

6 Acknowledgement

We thank Naomi Nishimura, Ian Goulden, and Wendy Rush for obtaining a copy of Bondy's 1980 research report [8].

References

- 1 Faisal N. Abu-Khzam, Michael R. Fellows, Michael A. Langston, and W. Henry Suters. Crown structures for vertex cover kernelization. *Theory Comput. Syst.*, 41(3):411–430, 2007.
- 2 Noga Alon, Gregory Gutin, Eun Jung Kim, Stefan Szeider, and Anders Yeo. Solving MAX-r-SAT above a tight lower bound. *Algorithmica*, 61(3):638–655, Nov 2011.
- 3 Noga Alon, Raphael Yuster, and Uri Zwick. Color-coding. *J. ACM*, 42(4):844–856, 1995.
- 4 Richard Bellman. Dynamic programming treatment of the travelling salesman problem. *J. Assoc. Comput. Mach.*, 9:61–63, 1962.
- 5 Ivona Bezáková, Radu Curticapean, Holger Dell, and Fedor V. Fomin. Finding detours is fixed-parameter tractable. In *ICALP 2017*, pages 54:1–54:14, 2017.
- 6 Andreas Björklund. Determinant sums for undirected Hamiltonicity. *SIAM J. Comput.*, 43(1):280–299, 2014.
- 7 J. A. Bondy and V. Chvátal. A method in graph theory. *Discrete Math.*, 15(2):111–135, January 1976.
- 8 J.A. Bondy. *Longest Paths and Cycles in Graphs of High Degree*. Research report. Department of Combinatorics and Optimization, University of Waterloo, 1980.
- 9 Benny Chor, Mike Fellows, and David W. Juedes. Linear kernels in linear time, or how to save k colors in $O(n^2)$ steps. In *Proc. 30th WG*, pages 257–269, 2004.
- 10 Robert Crowston, Mark Jones, and Matthias Mnich. Max-cut parameterized above the Edwards-Erdős bound. *Algorithmica*, 72(3):734–757, 2015.
- 11 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 12 Elias Dahlhaus, Péter Hajnal, and Marek Karpinski. On the parallel complexity of Hamiltonian cycle and matching problem on dense graphs. *J. Algorithms*, 15(3):367–384, 1993.
- 13 Reinhard Diestel. *Graph Theory, 4th Edition*, volume 173 of *Graduate texts in mathematics*. Springer, 2012.
- 14 G. A. Dirac. Some theorems on abstract graphs. *Proceedings of the London Mathematical Society*, s3-2(1):69–81, 1952.
- 15 Michael R. Fellows. Blow-ups, win/win’s, and crown rules: Some new directions in FPT. In Hans L. Bodlaender, editor, *Proc. 29th WG*, pages 1–12. Springer, 2003.
- 16 Jörg Flum and Martin Grohe. *Parameterized Complexity Theory*. Texts in Theoretical Computer Science. An EATCS Series. Springer-Verlag, Berlin, 2006.
- 17 Ronald J Gould. Recent advances on the Hamiltonian problem: Survey iii. *Graphs and Combinatorics*, 30(1):1–46, 2014.
- 18 Jiong Guo, Falk Hüffner, and Rolf Niedermeier. A structural view on parameterizing problems: Distance from triviality. In *IWPEC 2004*, pages 162–173, 2004.
- 19 Gregory Z. Gutin, Eun Jung Kim, Michael Lampis, and Valia Mitsou. Vertex cover problem parameterized above and below tight bounds. *Theory Comput. Syst.*, 48(2):402–410, 2011.
- 20 Gregory Z. Gutin, Eun Jung Kim, Stefan Szeider, and Anders Yeo. A probabilistic approach to problems parameterized above or below tight bounds. *J. Comput. Syst. Sci.*, 77(2):422–429, 2011.
- 21 Gregory Z. Gutin and Viresh Patel. Parameterized traveling salesman problem: Beating the average. *SIAM J. Discrete Math.*, 30(1):220–238, 2016.

- 22 Roland Häggkvist. On the structure of non-Hamiltonian graphs i. *Combinatorics, Probability and Computing*, 1(1):27–34, 1992.
- 23 Michael Held and Richard M. Karp. A dynamic programming approach to sequencing problems. *J. Soc. Indust. Appl. Math.*, 10:196–210, 1962.
- 24 Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *J. Comput. Syst. Sci.*, 63(4):512–530, 2001.
- 25 Richard M. Karp. Reducibility among combinatorial problems. In *Complexity of computer computations*, pages 85–103. Springer, 1972.
- 26 Richard M. Karp. Dynamic programming meets the principle of inclusion and exclusion. *Operations Research Letters*, 1(2):49 – 51, 1982.
- 27 D.E. Knuth. *The Art of Computer Programming: Updates; Pre-Fascicle 8A, A draft of section 7.2.2.4: Hamiltonian paths and cycles*. Number v. 4 in Addison-Wesley series in computer science and information proceedings. Addison-Wesley, 2018.
- 28 Daniela Kühn and Deryk Osthus. Hamilton cycles in graphs and hypergraphs: an extremal perspective. *arXiv preprint arXiv:1402.4268*, 2014.
- 29 E.L. Lawler, D.B. Shmoys, A.H.G.R. Kan, and J.K. Lenstra. *The Traveling Salesman Problem*. John Wiley & Sons, 1985.
- 30 Hao Li. Generalizations of Dirac’s theorem in Hamiltonian graph theory—a survey. *Discrete Mathematics*, 313(19):2034 – 2053, 2013. Cycles and Colourings 2011.
- 31 Meena Mahajan and Venkatesh Raman. Parameterizing above guaranteed values: Maxsat and maxcut. *J. Algorithms*, 31(2):335–354, 1999.
- 32 Meena Mahajan, Venkatesh Raman, and Somnath Sikdar. Parameterizing above or below guaranteed values. *J. Comput. Syst. Sci.*, 75(2):137–153, 2009.
- 33 C.St.J.A. Nash-Williams. Edge-disjoint Hamiltonian circuits in graphs with large valency. In L. Mirksy, editor, *Studies in Pure Mathematics*, pages 157–183. Academic Press, London, 1971.
- 34 Oystein Ore. Note on Hamilton circuits. *The American Mathematical Monthly*, 67(1):55–55, 1960.