Capillary zone electrophoretic separation of beta-blockers using citrate buffer at low pH

Citation for published version (APA):

Reijenga, J. C., & Ingelse, B. A. (1997). Capillary zone electrophoretic separation of beta-blockers using citrate buffer at low pH. *Journal of Chromatography, A*, 776(2), 361-. [https://doi.org/10.1016/S0021-9673\(97\)00438-X](https://doi.org/10.1016/S0021-9673(97)00438-X)

DOI:

[10.1016/S0021-9673\(97\)00438-X](https://doi.org/10.1016/S0021-9673(97)00438-X)

Document status and date:

Published: 01/01/1997

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

ELSEVIER

Journal of Chromatography A, 776 (1997) 361

JOURNAL OF
CHROMATOGRAPHY A

Discussion

Capillary zone electrophoretic separation of β -blockers using citrate buffer at low pH

J.C. Reijenga*, B.A. Ingelse

Laboratory of Instrumental Analysis, University of Technology, P.O. Box 513, 5600 MB Eindhoven, Netherlands

Received 17 December 1996; revised 21 February 1997; accepted 7 March 1997

Keywords: Ionic strength; pH effect; β -Blockers

In a recent contribution to this journal [1], Lin et al. report on the influence of ionic strength and pH on the separation of β -blockers. Although it is not disputed that these two parameters can be of key importance in capillary electrophoresis (CE) optimization, we disagree on the interpretation of the results obtained, because the authors have underestimated the importance of ionic strength and overestimated the effect of pH.

When investigating the effect of pH, ionic strength was not kept constant (as it should have been). In fact ionic strength changed by as much as a factor 17. Not surprisingly, mobilities changed by an average of 30%. These mobility changes can be different for each component, apparently resulting in minor selectivity changes, as seen in Fig. 5.

These selectivity changes certainly cannot be attributed to different extents of ionization, as sug-

gested by the authors, because in the pH range investigated, the components must be considered fully protonated.

In fact the pK values of the β -blockers are 8.8 for pindolol, 9.45 for propranolol, 9.68 for metoprolol [2] and are likely in the same range for the others. Therefore there is no reason to hope for selectivity based on a different extent of ionization of the analytes in the pH range investigated. Summarizing: selectivity effects reported are solely due to minor ionic strength effects.

References

- [1] C.-E. Lin, C.-C. Chang, W.-C. Lin, E.C. Lin, *J. Chromatogr. A* 753 (1996) 133–138.
- [2] G.S. Avery (Editor), *Drug Treatment*, Publishing Sciences Group, Littleton, MA, 1976, pp. 890–896.

*Corresponding author.