Multiparticle dispersion in fully developed turbulence

Citation for published version (APA):

Biferale, L., Boffetta, G., Celani, A., Devenish, B. J., Lanotte, A., & Toschi, F. (2005). Multiparticle dispersion in fully developed turbulence. *Physics of Fluids*, 17(1), 111701-1/4. <https://doi.org/10.1063/1.2130751>

DOI:

[10.1063/1.2130751](https://doi.org/10.1063/1.2130751)

Document status and date:

Published: 01/01/2005

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Multiparticle dispersion in fully developed turbulence

L. Biferale

Department of Physics and INFN, University of Tor Vergata, Via della Ricerca Scientifica 1, 00133 Rome, Italy

G. Boffetta

Dipartimento di Fisica Generale and INFN, Università degli Studi di Torino, Via Pietro Giuria 1, 10125 Torino, Italy and CNR-ISAC, Sezione di Torino, Corso Fiume 4, 10133 Torino, Italy

A. Celani

CNRS, INLN, 1361 Route des Lucioles, 06560 Valbonne, France

B. J. Devenish

Department of Physics, University of Tor Vergata, Via della Ricerca Scientifica 1, 00133 Rome, Italy

A. Lanotte

CNR-ISAC, Sezione di Lecce, Strada Prov. Lecce-Monteroni km. 1.200, 73100 Lecce, Italy

F. Toschi

Istituto per le Applicazioni del Calcolo, CNR, Viale del Policlinico 137, 00161 Rome, Italy and INFN Sezione di Ferrara, via G. Saragat 1, 44100 Ferrara, Italy

(Received 24 July 2005; accepted 6 October 2005; published online 2 November 2005)

The statistical geometry of dispersing Lagrangian clusters of four particles (tetrahedra) is studied by means of high-resolution direct numerical simulations of three-dimensional homogeneous isotropic turbulence. We give evidence of a self-similar regime of shape dynamics characterized by almost two-dimensional, strongly elongated geometries. The analysis of four-point velocity-difference statistics and orientation shows that inertial-range eddies typically generate a straining field with a strong extensional component aligned with the elongation direction and weak extensional/compressional components in the orthogonal plane. © 2005 American Institute of Physics.

[DOI: [10.1063/1.2130751](https://doi.org/10.1063/1.2130751)]

One of the most characteristic attributes of turbulence is the efficient dispersion and mixing of advected Lagrangian particles.¹ Even though turbulent dispersion bears some similarities to Brownian motion, especially at very large scales and for long times, it has a much richer structure at small scales. This is already visible at the level of single particle dispersion, which is characterized by nontrivial time correlations of the velocity experienced by the particle along its trajectory (see, e.g., Refs. 2–4). The statistics of pair dispersion display interesting properties as well (see, e.g., Refs. 5–10), yet the complexity of Lagrangian turbulence is particularly evident when looking at the dispersion of three or more particles. This calls for the description of the geometrical properties of Lagrangian dispersion—the “shape” of the particles’ cloud as well as its “size.” The geometrical characterization of dispersion proved extremely important for the understanding of the problem of passive scalar advection¹¹ and provided the basis for the efficient modeling of the small-scale velocity dynamics itself.^{12–15} Previous studies dealt with two-dimensional flows,^{16,17} synthetic flows,^{18,19} or three-dimensional turbulence at moderate Reynolds numbers.^{13,20,21} In this Letter we study multiparticle Lagrangian statistics by means of high-resolution direct numerical simulations of three-dimensional Navier-Stokes turbulence. Simulations were done at resolutions of 1024^3 corresponding to a Reynolds number $R_\lambda \sim 280$ (see Ref. 22). The other parameters of the numerical simulation are as fol-

lows: energy dissipation $\varepsilon = 0.81(8)$, viscosity $\nu = 8.8 \times 10^{-4}$, Kolmogorov length scale $\eta = 5 \times 10^{-3}$, integral scale $L = 3.14$, Lagrangian velocity autocorrelation time $T_L = 1.2$, Kolmogorov time scale $\tau_\eta = 3.3 \times 10^{-2}$. With the present choice of parameters the dissipative range of length scales is well resolved. Upon having reached a statistically stationary velocity field, the Lagrangian tracers were seeded in the flow. Their trajectories were integrated according to

$$\frac{d\mathbf{x}}{dt} = \mathbf{u}(\mathbf{x}(t), t)$$

over a time lapse of the order of a few Lagrangian correlation times, T_L . The velocity field, $\mathbf{u}$, results from the time integration of the three-dimensional Navier-Stokes equations (for further details see Ref. 22).

A set of 3.84×10^5 particles were initially seeded in quadruplets forming 9.6×10^4 regular tetrahedra of the size of the Kolmogorov scale, with centers of mass uniformly distributed over the domain. The evolution of the separations between different particles in each tetrahedron provides a way to quantify the shape evolution. As particles move with the flow, the sizes of the tetrahedra grow in time and their shapes deform, generating a variety of irregular objects. A description of this process is then given in terms of the probability density functions (pdf) of sizes and shapes. Within the inertial range of scales a self-similar evolution of size ac-

FIG. 1. Evolution of the mean eigenvalues g_1 (+), g_2 (x) and g_3 (*) of the moment of inertia matrix $\mathbf{I} = \boldsymbol{\rho}\boldsymbol{\rho}^T$. The line represents the dimensional scaling t^3 . In the inset, from top to bottom: evolution at small times of $\langle \ln A(t) \rangle$ (surface), $\langle \ln R(t) \rangle$ (distance), $\langle \ln V(t) \rangle$ (volume). The linear slopes of the three curves in the range of times $\tau_\eta < t < 3\tau_\eta$ yield $\lambda_1 + \lambda_2$, λ_1 , and $\lambda_1 + \lambda_2 + \lambda_3$, respectively.

cording to Richardson's law and a stationary shape distribution are expected.

In order to characterize the shape dynamics quantitatively, it is useful to introduce the following change of coordinates:¹² $\boldsymbol{\rho}_0 = (\mathbf{x}_1 + \mathbf{x}_2 + \mathbf{x}_3 + \mathbf{x}_4)/2$, $\boldsymbol{\rho}_1 = (\mathbf{x}_2 - \mathbf{x}_1)/\sqrt{2}$, $\boldsymbol{\rho}_2 = (2\mathbf{x}_3 - \mathbf{x}_2 - \mathbf{x}_1)/\sqrt{6}$, $\boldsymbol{\rho}_3 = (3\mathbf{x}_4 - \mathbf{x}_3 - \mathbf{x}_2 - \mathbf{x}_1)/\sqrt{12}$. By virtue of the statistical homogeneity of the velocity field as well as of the initial distribution of the centers of mass, the Lagrangian statistics do not depend on $\boldsymbol{\rho}_0$. The information about the particle separations can be embodied in the square matrix $\boldsymbol{\rho}$ whose columns are the three vectors $\boldsymbol{\rho}_i$ with $i=1,2,3$. Denoted by g_i ($g_1 \geq g_2 \geq g_3$) the eigenvalues of the moment of the inertia matrix, $\mathbf{I} = \boldsymbol{\rho}\boldsymbol{\rho}^T$ (that is positive defined), we have that the size of the tetrahedron is $r \equiv \sqrt{\text{tr}(\mathbf{I})} = \sqrt{g_1 + g_2 + g_3} = \sqrt{\frac{1}{8} \sum_{i,j} |\mathbf{x}_i - \mathbf{x}_j|^2}$, whereas the volume can be expressed as $V = \frac{1}{3} \det(\boldsymbol{\rho}) = \frac{1}{3} \sqrt{g_1 g_2 g_3}$. A convenient characterization of shapes is given in terms of the dimensionless quantities $I_i = g_i/r^2$ (where obviously $I_1 + I_2 + I_3 = 1$). For a regular tetrahedron one has $I_1 = I_2 = I_3 = 1/3$. If the four points are coplanar one has $I_3 = 0$ and for a collinear configuration $I_2 = I_3 = 0$.

Figure 1 shows the temporal evolution of the mean eigenvalues of $\boldsymbol{\rho}\boldsymbol{\rho}^T$ for the smallest regular tetrahedra with $g_i(0) = \delta x^2/2$. Two very different regimes are evident; at small times $t < \tau_\eta$ the evolution of tetrahedra is governed by the dissipative range of turbulence. Because of the smoothness and incompressibility of the velocity field in this range, the volume of each tetrahedron is approximately preserved and so is its average value which is shown in Fig. 1. In the viscous range the shape dynamics are essentially characterized by the Lagrangian Lyapunov exponents;²³ as a consequence the mean square separation r^2 grows exponentially in time. From the average growth rate of the logarithms of the separations, $R(t) = |\dot{\boldsymbol{\rho}}_1|$, areas $A(t) = \sqrt{3}/2 |\boldsymbol{\rho}_1 \times \boldsymbol{\rho}_2|$ and volumes $V(t) = \frac{1}{3} |\boldsymbol{\rho}_1 \times \boldsymbol{\rho}_2 \times \boldsymbol{\rho}_3|$ at small times, we can obtain an estimation of the Lagrangian Lyapunov spectrum as shown in Fig. 1. We found two positive Lyapunov exponents, with $\lambda_1 \tau_\eta \approx 0.12$ and $\lambda_2 \approx \lambda_1/4$, in agreement with previous findings at lower R_λ .^{24,25} The sum of the three Lyapunov expo-

FIG. 2. Doubling times for the eigenvalues, g_i , of the moment of inertia matrix, $\boldsymbol{\rho}\boldsymbol{\rho}^T$. In the inset: the same data rescaled on the horizontal axis with the proportions $g_1:g_2:g_3=40:8:1$.

nents so obtained is close to zero for times up to $3\tau_\eta$.

The exponential growth brings particle separations outside the dissipative range, where the velocity field becomes rough and the inertial range sets in. According to the Kolmogorov-Richardson scaling, eigenvalues should grow as $g_i \sim t^3$. As previously reported,¹³ it is hard to extract a clear scaling regime for the shape dynamics shown in Fig. 1. The main reason for the lack of self-similarity is due to the contamination of the inertial range by the dissipative range. Indeed, because of the strong shape distortion taking place at the crossover between the dissipative and inertial ranges (as shown in Fig. 1 by the separation of the three eigenvalues), a significant fraction of tetrahedra has one side in the dissipative range even at times much larger than τ_η . In order to overcome this problem we have utilized the technique of doubling time statistics that has already been successfully used to remove contaminations in the statistics of pair dispersion.^{10,26,27} At variance with fixed-time statistics, doubling times are essentially insensitive to the choice of the initial size of the tetrahedron.¹⁰ Here, we focus on the doubling times of the eigenvalues g_i : we compute the times, $T(g_i)$, taken by a tetrahedron to increase its value of g_i by a factor a . The result is shown in Fig. 2. The presence of a scaling range $T \sim g^{1/3}$ is more clear and the self-similarity is made evident by superimposing the three curves on top of each other by a simple multiplicative factor on the g axis. The ratio of the three eigenvalues in the scaling range is $g_1:g_2:g_3=40:8:1$, corresponding to shape indices $I_2 \approx 0.16$ and $I_3 \approx 0.02$. The presence of a range where the doubling times for different eigenvalues are the same is equivalent to stating that the typical shape of the tetrahedron is preserved while its size increases according to Richardson's law.

In view of the existence of a self-similar regime for shape evolution, one would expect that the statistics of the shape indices, I_i , should reach a time-independent distribution. However, a direct inspection of the data does not support this conclusion (not shown here, the results do not present an appreciable scaling range in time in spite of the relatively high R_λ as compared with Ref. 13). Once more this lack of a scaling range in the time domain can be traced back to the contamination by the dissipative range dynamics.

This difficulty can be overcome by selecting those tetra-

FIG. 3. Probability density function of shape indices I_2 and I_3 (inset) at times $t=35\tau_\eta$ (+) and $t=63\tau_\eta$ (x). The full lines are the pdfs for independent, Gaussian distributed particle positions.

hedra with eigenvalues in the ranges $5 \times 10^2 \eta^2 < g_1 < 5 \times 10^5 \eta^2$, $5 \times 10^1 \eta^2 < g_2 < 5 \times 10^4 \eta^2$, $5 \times \eta^2 < g_3 < 5 \times 10^3 \eta^2$. The thresholds are obtained by identifying the scaling ranges in Fig. 2. This procedure removes about 60% of the initial tetrahedra, mostly because g_3 falls below its lower threshold. The probability density functions of the shape indices I_2, I_3 after the selection are shown in Fig. 3. The existence of an invariant regime appears now very clearly. In this regime, the normalized probability density functions at different times collapse, and the mean values hence display a plateau in time; for the third index, the mean value $\langle I_3 \rangle \approx 0.011 \pm 0.001$ is not too far from the Gaussian value 0.03, while the second index is concentrated on values much smaller ($\langle I_2 \rangle \approx 0.135 \pm 0.003$ as opposed to 0.22).^{13,20} Those values indicate a relative abundance of flat and elongated configurations. The tendency to form almost two-dimensional structures has mostly an “entropic” origin; indeed there are a large number of pancakelike tetrahedra (very small I_3) already for Gaussian, independent particle positions, as shown by the corresponding distribution in Fig. 3. However, it has to be remarked that the pdf of I_3 is significantly more peaked at small values than the Gaussian one. The preference for elongated structures ($I_2 \ll I_1$) has a clear dynamical origin, since it has no equivalent in the Gaussian ensemble.

An interesting issue that we do not address here is connected with the possibility of subleading, anomalous scaling in the tetrahedra distribution. In the simpler case of particles advected by a Gaussian and white-in-time velocity field, it is known that the asymptotic behavior of the multiparticle pdf, when the initial points are close, is governed by an expansion in zero modes and slow modes of a given evolution operator.¹¹ There, anomalous corrections emerge as subleading terms to the Richardson scaling. These corrections are connected to the anomalous scaling of the structure functions of a passive scalar field advected by the flow. Here, in the presence of a real turbulent flow, one can expect that similar properties may still hold.¹⁶ In order to check this, one should perform a delicate compensation between the evolution of the pdf with different initial tetrahedra shapes, in order to cancel the leading scaling terms and to highlight the subleading contributions.

FIG. 4. Evolution of the mean eigenvalues of the “turbulent diffusion tensor,” $\mathbf{K}$, as a function of the tetrahedron size, r . In the inset, the eigenvalues as a function of time.

The dynamics of the shape evolution can be elucidated by analyzing the local geometrical properties of Lagrangian velocities. In analogy with the relative coordinates $\boldsymbol{\rho}$, we introduce the relative velocity matrix $\mathbf{W}$:¹² $\mathbf{W}_1 = (\mathbf{u}_2 - \mathbf{u}_1)/\sqrt{2}$, $\mathbf{W}_2 = (2\mathbf{u}_3 - \mathbf{u}_2 - \mathbf{u}_1)/\sqrt{6}$, $\mathbf{W}_3 = (3\mathbf{u}_4 - \mathbf{u}_3 - \mathbf{u}_2 - \mathbf{u}_1)/\sqrt{12}$. Obviously, $\dot{\boldsymbol{\rho}} = \mathbf{W}$. The geometrical aspects of Lagrangian velocity evolution can be described by the tetrahedron “turbulent diffusion” tensor

$$\mathbf{K} \equiv \frac{1}{2} \frac{d}{dt} \boldsymbol{\rho} \boldsymbol{\rho}^T = \frac{1}{2} (\mathbf{W} \boldsymbol{\rho}^T + \boldsymbol{\rho} \mathbf{W}^T). \quad (1)$$

The trace $\text{tr}(\mathbf{K}) = \frac{1}{8} \sum_{i,j} (\mathbf{u}_i - \mathbf{u}_j) \cdot (\mathbf{x}_i - \mathbf{x}_j)$ is proportional to the longitudinal velocity difference multiplied by the separation averaged over all pairs within the tetrahedron. The geometrical information about the Lagrangian velocity fluctuations may be obtained from the eigenvalues $\kappa_1 \geq \kappa_2 \geq \kappa_3$ of $\mathbf{K}$, which are shown in Fig. 4. On dimensional grounds these should grow in time as t^2 or, equivalently, with the tetrahedron size, r , as $r^{4/3}$. This is satisfied to a good accuracy for all three eigenvalues, especially as a function of size. The third eigenvalue, κ_3 , is negative (notice that, strictly speaking, this conflicts with the definition of $\mathbf{K}$ as a diffusion tensor). Geometrically this means that the local velocity field experienced by the tetrahedron has two extensional components: a strong one and a weak one, $\kappa_1 \gg \kappa_2$, with the latter smaller by a factor of 10 than the former, and a weak compressional component $|\kappa_3| \approx \kappa_2$. It is also interesting to study the relative orientation of the eigenvectors of the matrix $\mathbf{I} = \boldsymbol{\rho} \boldsymbol{\rho}^T$, i.e., the principal axes of inertia, and the eigenvectors of the matrix $\mathbf{K}$. We found that the directions of the eigenvectors associated with g_1 and κ_1 are preferentially aligned. About 45% of the tetrahedra show a relative angle smaller than $\pi/6$ (for a uniform distribution on a unit sphere one would have 13%). This agrees with the intuitive idea that strongly extensional velocity differences result in intense elongations approximately in the same direction. In the plane orthogonal to the first principal axis of inertia, the eigenvectors of $\mathbf{I}$ and $\mathbf{K}$ associated with the smaller eigenvalues are also aligned albeit to a lesser degree (about 25% of relative angles below $\pi/6$).

The overall geometrical picture that emerges is the following: Tetrahedra tend to be elongated, almost coplanar objects, subject to a straining velocity field that has a strong extensional part in the direction of elongation and relatively weak compressive and extensional contributions in the orthogonal plane of approximately equal magnitude. The recent advances in experimental techniques for particle tracking should soon allow precise measurements of shape dynamics in real turbulent flows.^{2,6,28} The joint effort on the numerical and experimental side can shed further light on the geometrical statistics of Lagrangian turbulence. This, in turn, will lead to the development of new, more effective parametrizations of small-scale turbulence, a problem of paramount importance for geophysical and industrial applications.

We are grateful to A. Pumir for discussions and useful suggestions. The simulations were performed on the IBM-SP4 of the Cineca (Bologna, Italy). We are grateful to C. Cavazzoni and G. Erbacher for resource allocation and precious technical assistance.

¹A. S. Monin and A. M. Yaglom, *Statistical Fluid Mechanics* (MIT Press, Cambridge, MA, 1975), Vol. 2.

²N. Mordant, P. Metz, O. Michel, and J. F. Pinton, "Measurement of Lagrangian velocity in fully developed turbulence," *Phys. Rev. Lett.* **87**, 214501 (2001).

³L. Chevillard, S. G. Roux, E. Leveque, N. Mordant, J. F. Pinton, and A. Arneodo, "Lagrangian velocity statistics in turbulent flows: Effects of dissipation," *Phys. Rev. Lett.* **91**, 214502 (2003).

⁴L. Biferale, G. Boffetta, A. Celani, B. J. Devenish, A. Lanotte, and F. Toschi, "Multifractal statistics of Lagrangian velocity and acceleration in turbulence," *Phys. Rev. Lett.* **93**, 064502 (2004).

⁵P. K. Yeung, "Direct numerical simulation of two-particle relative diffusion in isotropic turbulence," *Phys. Fluids* **6**, 3416 (1994).

⁶S. Ott and J. Mann, "An experimental investigation of the relative diffusion of particle pairs in three-dimensional turbulent flow," *J. Fluid Mech.* **422**, 207 (2000).

⁷T. Ishihara and Y. Kaneda, "Relative diffusion of a pair of fluid particles in the inertial subrange of turbulence," *Phys. Fluids* **14**, L69 (2002).

⁸G. Boffetta and I. M. Sokolov, "Statistics of two-particle dispersion in two-dimensional turbulence," *Phys. Fluids* **14**, 3224 (2002).

⁹P. K. Yeung and M. S. Borgas, "Relative dispersion in isotropic turbulence. Part 1. Direct numerical simulations and Reynolds-number depen-

dence," *J. Fluid Mech.* **503**, 93 (2004).

¹⁰L. Biferale, G. Boffetta, A. Celani, B. J. Devenish, A. Lanotte, and F. Toschi, "Lagrangian statistics of particle pairs in homogeneous isotropic turbulence," *Phys. Fluids* (to be published).

¹¹G. Falkovich, K. Gawędzki, and M. Vergassola, "Particles and fields in fluid turbulence," *Rev. Mod. Phys.* **73**, 913 (2001).

¹²M. Chertkov, A. Pumir, and B. I. Shraiman, "Lagrangian tetrad dynamics and the phenomenology of turbulence," *Phys. Fluids* **11**, 2394 (1999).

¹³A. Pumir, B. I. Shraiman, and M. Chertkov, "Geometry of Lagrangian dispersion in turbulence," *Phys. Rev. Lett.* **85**, 5324 (2000).

¹⁴A. Pumir, B. I. Shraiman, and M. Chertkov, "The Lagrangian view of energy transfer in turbulent flow," *Europhys. Lett.* **56**, 379 (2001).

¹⁵A. Pumir and B. Shraiman, "Lagrangian particle approach to large eddy simulations of hydrodynamic turbulence," *J. Stat. Phys.* **113**, 693 (2003).

¹⁶A. Celani and M. Vergassola, "Statistical geometry in scalar turbulence," *Phys. Rev. Lett.* **86**, 424 (2001).

¹⁷P. Castiglione and A. Pumir, "Evolution of triangles in a two-dimensional turbulent flow," *Phys. Rev. E* **64**, 056303 (2001).

¹⁸U. Frisch, A. Mazzino, and M. Vergassola, "Intermittency in passive scalar advection," *Phys. Rev. Lett.* **80**, 5532 (1998).

¹⁹K. A. I. Khan, A. Pumir, and J. C. Vassilicos, "Kinematic simulation of turbulent dispersion of triangles," *Phys. Rev. E* **68**, 026313 (2003).

²⁰M. S. Borgas, "Meandering plume models in turbulent flows," in *Proceedings of the 13th Australasian Fluid Mechanics Conference*, Monash University, Melbourne, 1988.

²¹P. K. Yeung, S. Xu, M. Borgas, and B. L. Sawford, "Scaling of multi-particle Lagrangian statistics in direct numerical simulations," in *Reynolds Number Scaling in Turbulent Flow*, edited by A. J. Smits (Kluwer, Dordrecht, 2002).

²²L. Biferale, G. Boffetta, A. Celani, A. Lanotte, and F. Toschi, "Particle trapping in three-dimensional fully developed turbulence," *Phys. Fluids* **17**, 021701 (2005).

²³A. Crisanti, M. Falcioni, G. Paladin, and A. Vulpiani, "Lagrangian chaos: Transport, mixing and diffusion in fluids," *Riv. Nuovo Cimento* **14**, 1 (1991).

²⁴S. S. Girimaji and S. B. Pope, "Material-element deformation in isotropic turbulence," *J. Fluid Mech.* **220**, 427 (1990).

²⁵G. Falkovich and A. Pumir, "Intermittent distribution of heavy particles in a turbulent flow," *Phys. Fluids* **16**, L47 (2004).

²⁶G. Boffetta and I. M. Sokolov, "Relative dispersion in fully developed turbulence: The Richardson's law and intermittency corrections," *Phys. Rev. Lett.* **88**, 094501 (2002).

²⁷V. Artale, G. Boffetta, A. Celani, M. Cencini, and A. Vulpiani, "Dispersion of passive tracers in closed basins: Beyond the diffusion coefficient," *Phys. Fluids* **9**, 3162 (1997).

²⁸G. A. Voth, A. La Porta, A. M. Crawford, J. Alexander, and E. Bodenschatz, "Measurement of particle accelerations in fully developed turbulence," *J. Fluid Mech.* **469**, 121 (2002).