Solution to Problem 91-19 : A finite part integral

Citation for published version (APA):

Boersma, J., & Doelder, de, P. J. (1992). Solution to Problem 91-19 : A finite part integral. *SIAM Review*, 34(6), 656-658. <https://doi.org/10.1137/1034129>

DOI:

[10.1137/1034129](https://doi.org/10.1137/1034129)

Document status and date:

Published: 01/01/1992

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

with respect to c_2 , we obtain

$$\int_0^\pi \frac{\sin^2 \theta}{g^2} d\theta = \frac{\pi}{2c_1 c_2^3}.$$

These integrals, on substitution in the formula for I , confirm that $I = c_1/c_2$ as desired.

Also solved by C. C. GROSJEAN (State University of Ghent, Belgium).

A Finite Part Integral

Problem 91-19, by B. BERTRAM AND O. G. RUEHR (Michigan Technological University).

Evaluate, in terms of elementary functions, the integral

$$I_\nu(x) = \oint_0^1 \frac{t^\nu}{(t-x)^2} dt, \quad 0 < x < 1$$

for rational $\nu > -1$. Here, the double bar denotes that the integral is to be taken in the Hadamard or finite part sense, which can be shown to be the derivative of the Cauchy principal value integral, i.e.,

$$\oint_0^1 \frac{f(t)}{(t-x)^2} dt = \frac{d}{dx} \int_0^1 \frac{f(t)}{t-x} dt.$$

The problem arose in a numerical analysis of errors encountered in applying product integration to linear finite part integral equations. Such equations occur in fracture mechanics, gas radiation, and fluid flow.

REFERENCES

- [1] A. C. KAYA AND F. ERDOGAN, *On the solution of integral equations with strongly singular kernels*, Quart. Appl. Math., 45 (1987), pp. 105–122.
- [2] J. HADAMARD, *Lectures on Cauchy's Problem in Linear Partial Differential Equations*, Dover, New York, 1952, pp. 133–141.
- [3] B. BERTRAM, *On the Use of Product Integration for the Solution of Finite Part Singular Integral Equations*, in *Integral Methods in Science and Engineering-90*, A. Haji-Sheikh, ed., Hemisphere, New York, 1991, pp. 132–136.
- [4] P. J. DAVIS AND P. RABINOWITZ, *Methods of Numerical Integration*, Second ed., Academic Press, New York, 1984.

Solution by J. BOERSMA and P. J. DE DOELDER (Eindhoven University of Technology, Eindhoven, the Netherlands).

Introduce the function

$$F_\nu(x) = \oint_0^1 \frac{t^\nu}{t-x} dt;$$

then the derivative $f'_\nu(x) = I_\nu(x)$ is to be evaluated for $0 < x < 1$ and rational $\nu > -1$.

Using [1, form. 6.2(4)], we determine the Mellin transform

$$\int_0^\infty F_\nu(x) x^{s-1} dx = \int_0^1 t^\nu dt \int_0^\infty \frac{x^{s-1}}{t-x} dx = \frac{\pi \cot(\pi s)}{s + \nu},$$

valid for $\max(0, -\nu) < \operatorname{Re} s < 1$. By inversion we obtain the representation

$$F_\nu(x) = \frac{1}{2i} \int_{c-i\infty}^{c+i\infty} \frac{\cot(\pi s)}{s + \nu} x^{-s} ds, \quad x > 0, \quad \max(0, -\nu) < c < 1.$$

By means of [1, form. 7.2(18)] the derivative $F'_\nu(x)$ can be expressed as

$$\begin{aligned} F'_\nu(x) &= -\frac{1}{2i} \int_{c-i\infty}^{c+i\infty} \frac{s}{s+\nu} \cot(\pi s) x^{-s-1} ds \\ &= -\frac{1}{x(1-x)} + \frac{\nu}{2i} \int_{c-i\infty}^{c+i\infty} \frac{\cot(\pi s)}{s+\nu} x^{-s-1} ds, \quad x > 0, \quad \max(0, -\nu) < c < 1. \end{aligned}$$

The latter integral is evaluated by closing the contour to the left, which is permissible if $0 < x < 1$. The integrand has poles at $s = -\nu$ and $s = -k$, $k = 0, 1, 2, \dots$. Notice that all poles are simple, except for a second-order pole at $s = -\nu$ if ν is integral.

In the case of integral ν , say $\nu = n$, $n = 0, 1, 2, \dots$, we thus find

$$\begin{aligned} F'_n(x) &= -\frac{1}{x(1-x)} - nx^{n-1} \log x + n \sum_{\substack{k=0 \\ k \neq n}}^{\infty} \frac{x^{k-1}}{n-k} \\ &= -\frac{1}{x(1-x)} + nx^{n-1} [\log(1-x) - \log x] + n \sum_{k=0}^{n-1} \frac{x^{k-1}}{n-k}, \quad 0 < x < 1. \end{aligned}$$

In the case of nonintegral $\nu > -1$ we find

$$(*) \quad F'_\nu(x) = -\frac{1}{x(1-x)} - \pi\nu \cot(\pi\nu) x^{\nu-1} + \nu \sum_{k=0}^{\infty} \frac{x^{k-1}}{\nu-k}, \quad 0 < x < 1.$$

If ν is rational, we may set $\nu = n - p/q$, where $n = [\nu] + 1$, and p and q are mutually prime integers with $0 < p < q$. The final series in (*) is now rewritten as

$$\nu \sum_{k=0}^{\infty} \frac{x^{k-1}}{\nu-k} = \nu \sum_{k=0}^{n-1} \frac{x^{k-1}}{\nu-k} - \nu x^{n-1} q \sum_{k=0}^{\infty} \frac{x^k}{p+qk}.$$

To evaluate the latter series we use a technique adopted from Nielsen [2, p. 21]. Starting from the well-known identity

$$\sum_{m=0}^{q-1} \left(e^{2\pi i m/q} \right)^{l-p} = \begin{cases} q & \text{if } l = p \pmod{q}, \\ 0 & \text{if } l \neq p \pmod{q}, \end{cases}$$

we have

$$\begin{aligned} q \sum_{k=0}^{\infty} \frac{t^{qk}}{p+qk} &= \sum_{l=1}^{\infty} \sum_{m=0}^{q-1} \frac{(te^{2\pi i m/q})^{l-p}}{l} \\ &= -t^{-p} \sum_{m=0}^{q-1} e^{-2\pi i m p/q} \log(1 - te^{2\pi i m/q}), \quad |t| < 1. \end{aligned}$$

By combining the previous results we find

$$\begin{aligned} F'_\nu(x) &= -\frac{1}{x(1-x)} + \nu x^{\nu-1} \left[\sum_{m=0}^{q-1} e^{2\pi i m \nu} \log(1 - x^{1/q} e^{2\pi i m/q}) - \pi \cot(\pi\nu) \right] \\ &\quad + \nu \sum_{k=0}^{[\nu]} \frac{x^{k-1}}{\nu-k}, \quad 0 < x < 1, \end{aligned}$$

valid for nonintegral rational ν . Here, q is the smallest positive integer such that $q\nu$ is integral.

REFERENCES

- [1] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER, AND F. G. TRICOMI, *Tables of Integral Transforms*, Vol. I, McGraw-Hill, New York, 1954.
 [2] N. NIELSEN, *Die Gammafunktion*, Chelsea, New York, 1965.
Also solved by C. C. GROSJEAN (University of Ghent, Ghent, Belgium) and W. F. FORD and J. A. PENNLINE (NASA Lewis Research Center) and the proposers. The proposers' solution, with a generalization, has appeared in *Product integration for finite-part singular integral equations; numerical asymptotics and convergence acceleration*, J. Comput. Appl. Math., 41 (1992), pp. 163–173.

Approximation for the Location of the Center of a Disk

*Problem 91-20**, by M. L. GLASSER AND J. KOPLOWITZ (Clarkson University).

A digital disk [1] consists of all points on a square lattice (of lattice spacing 1) that falls within a disk. If the radius of the disk is given, how accurately can one find the location of its center from the knowledge of the digital disk? Is the closest lattice point to a random circle of radius r of $O(1/r)$? If so, what is a good estimate for the coefficient?

REFERENCE

- [1] S. FISK, *Separating point sets by circles and recognition of digital disks*, IEEE Trans. Pattern Analysis and Machine Intelligence, 8 (1986), pp. 554–556.

Solution by H. L. ABBOTT (University of Alberta).

It will be shown first that the answer to the second question raised in the proposal is no. Let $f(r)$ denote the distance from the circle $x^2 + y^2 = r^2$ to the nearest exterior lattice point.

THEOREM 1. *There exists a positive constant c such that for arbitrarily large values of r ,*

$$f(r) > \frac{c(\log r)^{1/2}}{r}.$$

Proof. Denote by $h(z)$ the number of integers not exceeding z that may be written in the form $u^2 + v^2$, u , and v integers. It is a classical result of Landau [3] that there exists a positive constant α such that, as $z \rightarrow \infty$,

$$h(z) = (\alpha + o(1)) \frac{z}{(\log z)^{1/2}}.$$

Let $\ell = h(z) - h(z/2) + 1$, and let $m = z/2\ell$. For $j = 1, 2, \dots, \ell$, let $I_j = ((z/2) + (j-1)m, (z/2) + jm]$. Then $I_1, I_2, \dots, I_\ell$ are nonoverlapping intervals whose union is $I = (z/2, z]$. The number of integers in I of the form $u^2 + v^2$ is $h(z) - h(z/2)$, and hence, by the definition of ℓ , one of the subintervals, say I_t , does not contain any such number. Set $r^2 = z/2 + (t-1)m$ and $w^2 = (z/2) + tm$. Then the nearest exterior lattice point to the circle $x^2 + y^2 = r^2$ must lie outside the circle $x^2 + y^2 = w^2$. Thus

$$f(r) > w - r = \frac{w^2 - r^2}{w + r} = \frac{m}{w + r} > \frac{m}{2w} = \frac{z}{4w\ell}.$$