Exciton diffusion and annihilation in nanophotonic purcell landscapes

Citation for published version (APA):

Raziman, T. V., Visser, C. P., Wang, S., Rivas, J. G., & Curto, A. G. (2020). Exciton diffusion and annihilation in nanophotonic purcell landscapes. In *Frontiers in Optics - Proceedings Frontiers in Optics / Laser Science, Part of Frontiers in Optics + Laser Science APS/DLS, FIO 2020* [JTh4B.10] Optical Society of America (OSA).
<https://doi.org/10.1364/FIO.2020.JTh4B.10>

Document license:

Unspecified

DOI:

[10.1364/FIO.2020.JTh4B.10](https://doi.org/10.1364/FIO.2020.JTh4B.10)

Document status and date:

Published: 14/09/2020

Document Version:

Accepted manuscript including changes made at the peer-review stage

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Exciton Diffusion and Annihilation in Nanophotonic Purcell Landscapes

T. V. Raziman,^{1,*} C. Peter Visser,¹ Shaojun Wang,^{1,2} Jaime Gómez Rivas,¹ and Alberto G. Curto¹

¹Department of Applied Physics and Institute for Photonic Integration,
Eindhoven University of Technology, Eindhoven, The Netherlands

²MOE Key Laboratory of Modern Optical Technologies and Jiangsu Key Laboratory of Advanced Optical
Manufacturing Technologies, School of Optoelectronic Science and Engineering,
Soochow University, Suzhou 215006, China

*t.v.raziman@tue.nl

Abstract: Conventional nanophotonic emission enhancement neglects excitonic phenomena of diffusion and annihilation. We go beyond the localized Purcell effect and identify the enhancement mechanisms to turn their detrimental impact into additional emission.

© 2020 The Author(s)

Excitonic emitters in semiconductors such as perovskites, transition metal dichalcogenides and organic crystals disperse through diffusion and interact through exciton-exciton annihilation. [1–5] Nanophotonic resonators can improve light emission by enhancing excitation and radiative rates, and beaming the emission. [6–10] The conventional Purcell effect treats emitters like molecules and quantum dots as immobile and non-interacting. Total enhancement thus benefits from the product of excitation and emission at a point, which guides the design of nanoresonators and metamaterials made of metals and dielectrics. However, this approach is inadequate for excitonic emitters since it neglects diffusion over lengths comparable to nanophotonic scales (10 – 500 nm) and strong annihilation, which degrades the performance of light-emitting devices at high powers. [11]

Here, we go beyond the localized Purcell effect to exploit exciton dynamics for improving emission. We analyze the interplay of diffusion, annihilation, and nanophotonic enhancement for thin films of excitonic emitters in nanostructured landscapes (Figure 1). We provide analytical results for photoluminescence enhancement in the limiting regimes of high and low diffusion and annihilation. As excitons diffuse through optical hotspots and interact with each other, the balance of excitonic properties and nanophotonic geometry leads to either enhanced or suppressed photoluminescence. We present the conditions to turn the usually detrimental impact of diffusion on emission into additional enhancement and to overcome the adverse effects of annihilation at high exciton densities. Diffusion can increase photoluminescence by taking excitons to highly radiative locations when excitation and emission are spatially decoupled. Radiative rate enhancement can ameliorate the loss of efficiency arising from annihilation, while the interplay between annihilation and diffusion can improve performance by redistributing the local exciton density. Careful design of the relative strengths and spatial overlap of excitation and emission enhancements by tailoring the resonances and illumination conditions to the excitonic properties is thus key to efficient excitonic-nanophotonic systems. Our guidelines for customizing nanophotonic structures to diffusing and annihilating excitons will aid the design of efficient light-emitting devices based on excitonic nanomaterials.

Fig. 1. Photoluminescence from a film of excitonic emitters near a nanophotonic structure. Excitons form in the near field (yellow), diffuse through the film (orange arrows), and annihilate by interacting with other excitons (red) before decaying radiatively with enhancement from the Purcell effect (green) or non-radiatively (grey).

References

1. T. V. Raziman, C. P. Visser, S. Wang, J. Gómez Rivas, and A. G. Curto, “Exciton diffusion and annihilation in nanophotonic Purcell landscapes,” arXiv:2006.06288 (2020).
2. S. Deng, E. Shi, L. Yuan, L. Jin, L. Dou, and L. Huang, “Long-range exciton transport and slow annihilation in two-dimensional hybrid perovskites,” *Nat. Commun.* **11**, 1–8 (2020).
3. L. Yuan, T. Wang, T. Zhu, M. Zhou, and L. Huang, “Exciton Dynamics, Transport, and Annihilation in Atomically Thin Two-Dimensional Semiconductors,” *J. Phys. Chem. Lett.* **8**, 3371–3379 (2017).
4. O. V. Mikhnenko, P. W. M. Blom, and T.-Q. Nguyen, “Exciton diffusion in organic semiconductors,” *Energy Environ. Sci.* **8**, 1867–1888 (2015).
5. D. Sun, Y. Rao, G. A. Reider, G. Chen, Y. You, L. Brézin, A. R. Harutyunyan, and T. F. Heinz, “Observation of Rapid Exciton–Exciton Annihilation in Monolayer Molybdenum Disulfide,” *Nano Lett.* **14**, 5625–5629 (2014).
6. S. Bidault, M. Mivelle, and N. Bonod, “Dielectric nanoantennas to manipulate solid-state light emission,” *J. Appl. Phys.* **126**, 094104 (2019).
7. T. V. Raziman, R. H. Godiksen, M. A. Müller, and A. G. Curto, “Conditions for Enhancing Chiral Nanophotonics near Achiral Nanoparticles,” *ACS Photonics* **6**, 2583–2589 (2019).
8. E. Mohammadi, A. Tavakoli, P. Dehkoda, Y. Jahani, K. L. Tsakmakidis, A. Tittl, and H. Altug, “Accessible Superchiral Near-Fields Driven by Tailored Electric and Magnetic Resonances in All-Dielectric Nanostructures,” *ACS Photonics* **6**, 1939–1946 (2019).
9. A. F. Cihan, A. G. Curto, S. Raza, P. G. Kik, and M. L. Brongersma, “Silicon Mie resonators for highly directional light emission from monolayer MoS₂,” *Nat. Photon.* **12**, 284–290 (2018).
10. G. Vecchi, V. Giannini, and J. Gómez Rivas, “Shaping the fluorescent emission by lattice resonances in plasmonic crystals of nanoantennas,” *Phys. Rev. Lett.* **102**, 146807 (2009).
11. Y. Yu, Y. Yu, C. Xu, A. Barrette, K. Gundogdu, and L. Cao, “Fundamental limits of exciton-exciton annihilation for light emission in transition metal dichalcogenide monolayers,” *Phys. Rev. B* **93**, 201111 (2016).