A skew Hadamard matrix of order 36

Citation for published version (APA):

Goethals, J. M., & Seidel, J. J. (1970). A skew Hadamard matrix of order 36. *Journal of the Australian Mathematical Society*, 11(3), 343-344. <https://doi.org/10.1017/S144678870000673X>

DOI:

[10.1017/S144678870000673X](https://doi.org/10.1017/S144678870000673X)

Document status and date:

Published: 01/01/1970

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

A SKEW HADAMARD MATRIX OF ORDER 36

J. M. GOETHALS and J. J. SEIDEL

(Received 3 April 1969)

Hadamard matrices exist for infinitely many orders $4m$, $m \geq 1$, m integer, including all $4m < 100$, cf. [3], [2]. They are conjectured to exist for all such orders. Skew Hadamard matrices have been constructed for all orders $4m < 100$ except for 36, 52, 76, 92, cf. the table in [4]. Recently Szekeres [6] found skew Hadamard matrices of the order $2(p^t + 1) \equiv 12 \pmod{16}$, p prime, thus covering the case 76. In addition, Blatt and Szekeres [1] constructed one of order 52. The present note contains a skew Hadamard matrix of order 36 (and one of order 52), thus leaving 92 as the smallest open case.

The unit matrix of any order is denoted by I . The square matrices Q and R of order m are defined by their only nonzero elements

$$q_{i,i+1} = q_{m,1} = 1, \quad i = 1, \dots, m-1; \quad r_{i,m-i+1} = 1, \quad i = 1, \dots, m$$

We have

$$Q^m = I, \quad R^2 = I, \quad RQ = Q^T R.$$

Any square matrix A of order m is symmetric if $A = A^T$, skew if $A + A^T = 0$, circulant if $AQ = QA$. Hence, for circulant A we have

$$A = \sum_{i=0}^{m-1} a_i Q^i, \quad RA = A^T R.$$

Any square matrix H of order $4m$ is *skew Hadamard* if its elements are 1 and -1 (we write $+$ and $-$) and

$$HH^T = 4mI, \quad H + H^T = 2I.$$

THEOREM 1. *If A, B, C, D are square circulant matrices of order m , if A is skew, and if*

$$AA^T + BB^T + CC^T + DD^T = (4m-1)I,$$

then

$$H = \begin{bmatrix} A+I & BR & CR & DR \\ -BR & A+I & -D^T R & C^T R \\ -CR & D^T R & A+I & -B^T R \\ -DR & -C^T R & B^T R & A+I \end{bmatrix}$$

satisfies $HH^T = 4mI, H + H^T = 2I$.

PROOF. By straightforward verification.

REMARK. If, in addition, B , C , and D are symmetric, then H may be written in terms of the quaternion matrices K_4 , L_4 , M_4 and the Kronecker product $\otimes$ as follows:

$$H = I_4 \otimes (I + A) + K_4 \otimes BR + L_4 \otimes CR + M_4 \otimes DR,$$

hence looking much like a Williamson-type matrix, cf. [7].

THEOREM 2. *There exist skew Hadamard matrices of orders 36 and 52.*

PROOF. We apply theorem 1 with the following circulant matrices of order 9:

$$A = (0 \quad + \quad + \quad - \quad + \quad - \quad + \quad - \quad -), \quad B = (+ \quad - \quad + \quad + \quad - \quad - \quad + \quad + \quad -), \\ C = (- \quad - \quad + \quad + \quad + \quad + \quad + \quad + \quad -), \quad D = (+ \quad + \quad + \quad - \quad + \quad + \quad - \quad + \quad +).$$

By inspection the skew A and the symmetric B , C , D are seen to satisfy the hypotheses. Hence a skew Hadamard matrix of order 36 is obtained. Secondly, we consider the following circulant matrices of order 13:

$$A = (0 \quad + \quad + \quad + \quad - \quad + \quad + \quad - \quad - \quad + \quad - \quad - \quad -), \\ B = (- \quad + \quad - \quad + \quad + \quad - \quad - \quad - \quad - \quad + \quad + \quad - \quad +), \\ C = D = (- \quad - \quad + \quad - \quad + \quad + \quad + \quad + \quad + \quad - \quad + \quad + \quad +).$$

Application of theorem 1 to A , B , C , D yields a skew Hadamard matrix of order 52 since

$$AA^T = 15I - J + 2B, \quad BB^T = 12I - J - 2B, \quad CC^T = DD^T = 12I + J.$$

REMARK. The positive elements of B indicate the quadratic residues mod 13. The matrix of order 26

$$\begin{bmatrix} B+I & C \\ C^T & -B-I \end{bmatrix}$$

is an orthogonal matrix with zero diagonal, cf. [2] p. 1007. The matrix A describes the unique tournament of order 13 having no transitive subtournament of order 5, which was recently found by Reid and Parker [5].

References

- [1] D. Blatt and G. Szekeres, 'A skew Hadamard matrix of order 52', *Canadian J. Math.*, to appear.
- [2] J. M. Goethals and J. J. Seidel, 'Orthogonal matrices with zero diagonal', *Canadian J. Math.* 19 (1967), 1001–1010.
- [3] M. Hall, *Combinatorial theory* (Blaisdell 1967).
- [4] E. C. Johnsen, 'Integral solutions to the incidence equation for finite projective plane cases of orders $n \equiv 2 \pmod{4}$ ', *Pacific J. Math.* 17 (1966), 97–120.
- [5] K. B. Reid, and E. T. Parker, 'Disproof of a conjecture of Erdős and Moser on tournaments', *J. Combinatorial Theory*, to appear.
- [6] G. Szekeres, 'Tournaments and Hadamard matrices', *l'Enseign. Math.*, 15 (1969), 269–278.
- [7] J. Williamson, 'Hadamard's determinant theorem and the sum of four squares', *Duke Math. J.* 11, (1944), 65–81.