2,4-Di(butylureido)-6-methyl-s-triazine

Citation for published version (APA):

Kooijman, H., Spek, A. L., Beijer, F. H., Sijbesma, R. P., & Meijer, E. W. (2003). 2,4-Di(butylureido)-6-methyl-s-triazine. *Acta Crystallographica, Section E: Structure Reports*, E59(10), o1546-o1548.
<https://doi.org/10.1107/S160053680302018X>

DOI:

[10.1107/S160053680302018X](https://doi.org/10.1107/S160053680302018X)

Document status and date:

Published: 01/01/2003

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,4-Di(butylureido)-6-methyl-s-triazine

Huub Kooijman, Anthony L. Spek, Felix H. Beijer, Rint P. Sijbesma and E. W. Meijer

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

2,4-Di(butylureido)-6-methyl-s-triazine

Huub Kooijman,^{a*} Anthony L. Spek,^a Felix H. Beijer,^{b†} Rint P. Sijbesma^b and E. W. Meijer^b

^aBijvoet Center for Biomolecular Research, Department of Crystal and Structural Chemistry, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands, and ^bLaboratory of Macromolecular and Organic Chemistry, Eindhoven University of Technology, PO Box 513, 5600 MB Eindhoven, The Netherlands

† Current address: DSM Research, PO Box 18, 6160 MD Geleen, The Netherlands

Correspondence e-mail: h.kooijman@chem.uu.nl

Key indicators

Single-crystal X-ray study

$T = 150\text{ K}$

Mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$

Disorder in main residue

R factor = 0.077

wR factor = 0.187

Data-to-parameter ratio = 15.3

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

The formation of a quadruply hydrogen-bonded dimer of the title compound, $\text{C}_{14}\text{H}_{25}\text{N}_7\text{O}_2$, is prevented by an intramolecular hydrogen bond, which disrupts the linear *ADADA* donor–acceptor array. An infinite two-dimensional hydrogen-bonded network is formed instead.

Comment

The title compound, (I), was synthesized as part of a study on multiply hydrogen-bonded systems, focusing on the *DADA* motif in quadruply hydrogen-bonded dimers of triazines and pyrimidines (Beijer *et al.*, 1998). A review of quadruply hydrogen-bonded systems with several types of donor–acceptor arrays has recently been published by Sijbesma & Meijer (2003). When both ureido moieties adopt a particular *cis,trans* conformation, the title compound can arrange its hydrogen-bond donors and acceptors in such a way that a linear array of type *ADADA* (see Scheme) is formed. In this conformation, a quadruply hydrogen-bonded dimer can be formed. One of the peripheral acceptors is not directly involved in the formation of this dimer.

An atomic displacement ellipsoid plot of the title compound is given in Fig. 1, together with the atomic labelling scheme. Torsion angles describing the conformation of the ureido moieties are given in Table 1. One of the butylureido substituents of the triazine ring displays conformational disorder in the butyl moiety, as shown in Fig. 1. The disorder can be satisfactorily described with a two-site disorder model. The occupancy of the major component refined to a value of 0.533 (8).

Both ureido moieties adopt a *cis,trans* conformation, stabilized by intramolecular hydrogen bonds. The formation of a quadruply hydrogen-bonded dimer is, however, not observed in the crystalline state, because one of the intramolecular hydrogen bonds that stabilize the conformation of the molecule is donated to N103 (the triazine N located between the two ureido substituents) rather than towards N105 (a triazine N located between a ureido moiety and the methyl substituent). As a result, no linear array of type

Received 9 September 2003

Accepted 11 September 2003

Online 24 September 2003

Figure 1

View of the title compound, with the atom-numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level. H atoms are drawn as spheres of arbitrary size. The minor disorder component is indicated by dashed bonds.

ADADA is formed, and dimerization *via* quadruple hydrogen-bonding is impossible. Instead, an infinite hydrogen-bonded network is present (see Fig. 2). The hydrogen bond donated by amine N108 links molecules related by a crystallographic twofold screw axis into chains running in the direction [010]. The graph set (Bernstein *et al.*, 1995) of this unitary motif is $C(4)$. These hydrogen-bonded chains of molecules are linked into a two-dimensional network through hydrogen bonds donated by amine N116. At the unitary level, this hydrogen bond forms dimers about a crystallographic inversion centre. The graph set of the hydrogen-bonded dimer is $R_2^2(8)$. Base vectors of the two-dimensional network are [010] and [001]. Geometric details of the hydrogen bonds are given in Table 2.

Experimental

A suspension of 2,4-diamino-6-methyl-*s*-triazine (1.25 g, 10 mmol) and butyl isocyanate (11.3 ml, 100 mmol) in dry pyridine (50 ml) was heated under reflux for 16 h. After this period, water was added to the resultant solution to hydrolyse the excess of butyl isocyanate. After removal of the solvent, the resulting solid was crystallized from ethanol/water 2:1 *v/v*. Column chromatography (6% THF in chloroform), followed by crystallization from ethyl acetate and treatment with active carbon, gave an analytically pure sample (0.53 g, 16%), m.p. 502 K. $^1\text{H-NMR}$ (CDCl_3) δ : 10.17 (*br*), 9.05 (*br*), 3.32 (*q*, 4H), 2.42 (*s*, 3H), 1.52 (*m*, 4H), 1.34 (*m*, 4H), 0.89 (*t*, 6H). $^{13}\text{C-NMR}$ (CDCl_3) δ : 175.1, 162.8, 153.6, 39.4, 31.1, 25.2, 19.8, 13.3. IR (KBr) ν : 3407, 3266, 3203, 2956, 2930, 2918, 2871, 1704, 1684, 1593, 1526, 1479 cm^{-1} . Analysis: calcd. for $\text{C}_{14}\text{H}_{25}\text{N}_7\text{O}_2$: C, 52.0; H, 7.79; N, 30.32; found: C, 51.93; H, 7.70; N, 30.47.

Crystals suitable for X-ray structure determination were obtained by slow evaporation of a solution in ethanol.

Crystal data

$\text{C}_{14}\text{H}_{25}\text{N}_7\text{O}_2$
 $M_r = 323.41$
 Monoclinic, $C2/c$
 $a = 29.54$ (2) Å
 $b = 4.613$ (5) Å
 $c = 26.89$ (2) Å
 $\beta = 109.34$ (7)°
 $V = 3458$ (5) Å³
 $Z = 8$

$D_x = 1.242$ Mg m⁻³
 Mo $K\alpha$ radiation
 Cell parameters from 25 reflections
 $\theta = 8.2\text{--}17.1^\circ$
 $\mu = 0.09$ mm⁻¹
 $T = 150$ K
 Plate, colourless
 $0.50 \times 0.10 \times 0.04$ mm

Figure 2

Two-dimensional hydrogen-bonded network. The butyl and methyl moieties, as well as the H atoms not involved in hydrogen bonding, have been omitted for clarity. The hydrogen-bond motifs with graph set $C(4)$ are highlighted in green, the motifs with graph set $R_2^2(8)$ are highlighted in blue.

Data collection

Enraf-Nonius CAD-4 Turbo diffractometer
 ω scans
 Absorption correction: none
 10715 measured reflections
 3141 independent reflections
 1113 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.284$

$\theta_{\text{max}} = 25.3^\circ$
 $h = -34 \rightarrow 34$
 $k = 0 \rightarrow 5$
 $l = -32 \rightarrow 32$
 3 standard reflections
 frequency: 60 min
 intensity decay: 5%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.077$
 $wR(F^2) = 0.187$
 $S = 1.11$
 3141 reflections
 205 parameters

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.01P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.31$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.36$ e Å⁻³

Table 1

Selected geometric parameters (°).

C109—N108—C102—N103	−174.1 (5)	C117—N116—C104—N103	0.6 (8)
C102—N108—C109—O110	172.3 (5)	C104—N116—C117—O118	175.4 (5)
C112—N111—C109—O110	−2.5 (9)	C120—N119—C117—O118	−2.9 (8)

Table 2

Hydrogen-bonding geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N108—H104...O110 ⁱ	0.88	1.99	2.861 (6)	173
N111—H105...N101	0.88	1.96	2.653 (6)	135
N116—H115...O118 ⁱⁱ	0.88	1.95	2.818 (5)	170
N119—H116...N103	0.88	2.01	2.690 (7)	133

Symmetry codes: (i) $\frac{1}{2} - x, y - \frac{1}{2}, \frac{1}{2} - z$; (ii) $\frac{1}{2} - x, \frac{1}{2} - y, -z$.

H atoms were placed at calculated positions, riding on their carrier atoms. The H atoms of the methyl moieties were refined as rigid groups, allowing for rotation around the C—C bond. Isotropic displacement parameters of H atoms were set to 1.5 or 1.2 times the equivalent isotropic displacement parameter of the carrier atom for methyl and other H atoms, respectively. The C atoms of the major disorder component were refined with isotropic displacement parameters. The displacement parameters of the minor component atoms were equated to those of the corresponding atoms in the major component. The crystals of the title compound diffracted very poorly, displaying broad, weak reflections. The measured intensities therefore have a high standard uncertainty, as is shown by the value of R_σ [$= \Sigma \sigma(I)/\Sigma I$], which amounts to 0.1998. As a consequence, $R\text{-int}$ is also relatively high. However, upon merging 13314 measured

reflections into 3141 unique reflections, only 182 were found to be significantly inconsistent. Several batches of the compound were crystallized. A total of 6 different crystals were placed on the diffractometer. All of these showed poor diffraction. Full data were collected for three different crystals, all of which showed high $R\text{-int}$ values (up to 0.42). After solving the structure, all data sets revealed essentially the same structure. None of the data sets showed any of the classic signs of twinning. H atoms were placed at calculated positions, riding on their carrier atoms.

Data collection: locally modified *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *SET4* (de Boer & Duisenberg, 1984); data reduction: *HELENA* (Spek, 1997); program(s) used to solve structure: *SHELXS86* (Sheldrick, 1986); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *PLATON*.

This work was supported by the Council for the Chemical Sciences of the Netherlands Organization for Scientific Research (CW-NWO).

References

- Beijer, F. H., Kooijman, H., Spek, A. L., Sijbesma, R. P., & Meijer E. W. (1998). *Angew. Chem. Int. Ed. Engl.* **37**, 75–78.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Boer, J. L. de & Duisenberg, A. J. M. (1984). *Acta Cryst.* **A40**, C-410.
- Enraf–Nonius (1989). *CAD-4 Software*. Version 5. Enraf–Nonius, Delft, The Netherlands.
- Sheldrick, G. M. (1986). *SHELXS86*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Sijbesma, R. P. & Meijer, E. W. (2003). *Chem. Commun.* pp 5–16.
- Spek, A. L. (1997). *HELENA*. Utrecht University, The Netherlands.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.