Solution to Problem 79-15: An identity

Citation for published version (APA):

Lossers, O. P. (1980). Solution to Problem 79-15: An identity. *SIAM Review*, 22(3), 372-373.
<https://doi.org/10.1137/1022068>

DOI:

[10.1137/1022068](https://doi.org/10.1137/1022068)

Document status and date:

Published: 01/01/1980

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

When θ is small, M is a slowly-varying function of v , its Fourier harmonics are small, and the series for P_n converges rapidly. We expand M in a power series in θ :

$$\begin{aligned}\sin^2 \frac{1}{2}\theta \cos^2 u &= \sin^2 \frac{1}{2}\theta - \sin^2 \left(\frac{1}{2}\theta \sin v\right) \\ &= \sin\left[\frac{1}{2}\theta(1 - \sin v)\right] \sin\left[\frac{1}{2}\theta(1 + \sin v)\right].\end{aligned}$$

Let $v = 90^\circ - 2w$, $p = \theta \sin^2 w$, $q = \theta \cos^2 w$. Then,

$$M^2 = \frac{\sin^2 2w \sin^2 \frac{1}{2}\theta}{\sin p \sin q} = \frac{4pq \sin^2 \frac{1}{2}\theta}{\theta^2 \sin p \sin q}.$$

Now,

$$\begin{aligned}\frac{\sin p \sin q}{pq} &= \left(1 - \frac{\theta^2}{6} \sin^4 w + \cdots\right) \left(1 - \frac{\theta^2}{6} \cos^4 w + \cdots\right) \\ &= 1 - \frac{\theta^2}{24} (3 + \cos 4w) + \cdots,\end{aligned}$$

so

$$M = \frac{2}{\theta} \sin \frac{1}{2}\theta \left[1 + \frac{\theta^2}{48} (3 - \cos 2v) + O(\theta^4)\right].$$

Through θ^2 there are therefore no terms with $2k > 2$, while the coefficient of J_2 is

$$\frac{4}{\pi} \int_0^{\pi/2} -\frac{\theta^2}{48} \cos^2 2v \, dv = -\frac{\theta^2}{48},$$

and our approximation is

$$P_n(\cos \theta) \cong \frac{2}{\pi} K \left(\sin \frac{1}{2}\theta \right) J_0(x) - \frac{\theta^2}{48} J_2(x).$$

The functions J_0 and $-J_2$ are roughly in phase with each other, and they both have peaks that decrease with increasing x , or n . This proves the conjecture so long as θ is small enough for the approximation to be valid. But calculation shows that, at $\theta = 45^\circ$, the two Bessel terms give about 0.1% error in the first few peaks and in the peaks at $n = 80.5$ and $n = 4000.5$. The hypergeometric formula says successive peaks are $\pi/\theta = 4$ apart in n , so their amplitude ratio is $(n/(n+4))^{1/2}$. For this to be as small as even twice our 0.1%, n must be over 1000 and the "large n " proof is applicable. We conclude the conjecture is true in the range $0 < \theta < 45^\circ$, and therefore for all θ .

REFERENCE

- [1] M. ABRAMOWITZ AND I. STEGUN, *Handbook of Mathematical Functions*, U.S. Government Printing Office, National Bureau of Standards, 1965.

An Identity

Problem 79-15, by J. D. LOVE (Australian National University, Canberra, Australia).

In an analysis of the electrostatic potential for two charged dielectric spheres, the following identity arises:

$$\begin{aligned}\operatorname{csch} x &= P_n(\cosh x) Q_n(\cosh x) + Q_n(\cosh x) \sum_{m=0}^{n-1} P_m(\cosh x) e^{(n-m)x} \\ &\quad + P_n(\cosh x) \sum_{m=n+1}^{\infty} Q_m(\cosh x) e^{(n-m)x}\end{aligned}$$

where $P_n(\cosh x)$ and $Q_n(\cosh x)$ are modified Legendre functions of the first and second kinds, respectively. Prove the identity for real $x > 0$ and nonnegative integers n .

Solution by O. P. LOSSERS (Eindhoven University of Technology, Eindhoven, The Netherlands).

Let the right-hand side of the identity be denoted by S . Then by use of the integral [1, p. 154]

$$P_n(\cosh x)Q_m(\cosh x) = \frac{1}{2} \int_{-1}^1 \frac{P_n(t)P_m(t)}{\cosh x - t} dt, \quad n \leq m,$$

S can be reduced to

$$S = \frac{1}{2} e^{nx} \int_{-1}^1 \frac{P_n(t)}{\cosh x - t} \left(\sum_{m=0}^{\infty} P_m(t) e^{-mx} \right) dt.$$

The infinite series can be evaluated by means of the generating function for Legendre polynomials [1, p. 154], viz.,

$$\sum_{m=0}^{\infty} P_m(t) e^{-mx} = (1 - 2e^{-x}t + e^{-2x})^{-1/2} = 2^{-1/2} e^{1/2x} (\cosh x - t)^{-1/2},$$

thus leading to

$$S = 2^{-3/2} e^{(n+1/2)x} \int_{-1}^1 \frac{P_n(t)}{(\cosh x - t)^{3/2}} dt.$$

Because of [2, p. 822, 7.225 (4)],

$$\int_{-1}^1 \frac{P_n(t)}{(\cosh x - t)^{1/2}} dt = \frac{2\sqrt{2}}{2n+1} e^{-(2n+1)(x/2)}.$$

It follows by differentiation with respect to x that

$$\int_{-1}^1 \frac{P_n(t)}{(\cosh x - t)^{3/2}} dt = \frac{2\sqrt{2}}{\sinh x} e^{-(n+1/2)x},$$

which shows that

$$S = \operatorname{csch} x.$$

Also solved by H. E. FETTIS (Mountain View, California), who used an induction argument *and by* D. K. ROSS (La Trobe University, Australia), who used recurrence relations satisfied by the Legendre functions.

REFERENCES

- [1] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER AND F. G. TRICOMI, *Higher Transcendental Functions*, Vol. I, McGraw-Hill, New York, 1953.
- [2] I. S. GRADSTEYN AND I. M. RYZHIK, *Tables of Integrals, etc.*, Academic Press, New York, 1965.

ERRATUM: An Integral Inequality

Problem 77–6 by J. E. WILKINS, JR.

The erratum given for this problem (this Review, 22(1980) p. 102) is in error. It should have read: