Thermal rearrangements of (substituted allyl)dialkyl-2H-pyrroles

Citation for published version (APA):

Patterson, J. M., Ferry, J. D., Haan, de, J. W., & Boyd, M. R. (1975). Thermal rearrangements of (substituted allyl)dialkyl-2H-pyrroles. *Journal of the American Chemical Society*, 97(2), 360-362.
<https://doi.org/10.1021/ja00835a023>

DOI:

[10.1021/ja00835a023](https://doi.org/10.1021/ja00835a023)

Document status and date:

Published: 01/01/1975

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

- (39) 1,3-Diphenylisobenzothiophen has shown the similar photochemical behavior. The fluorescence of 1,3-diphenylisobenzothiophen was readily quenched by **2** ($K_q\tau_s = 3.3 \pm 0.3 \text{ M}^{-1}$ in ethanol), and the adduct formation was observed. The detailed results will be reported at a later date.
- (40) (a) K. N. Houk and R. B. Woodward, *J. Amer. Chem. Soc.*, **92**, 4143

- (1970); (b) K. N. Houk, L. J. Luskus, and N. S. Bhacca, *ibid.*, **92**, 6392 (1970).
- (41) S. Ito, K. Sakan, and Y. Fujise, *Tetrahedron Lett.*, 2873 (1970).
- (42) The [6 + 4] adduct is the major product in the reaction of tropone with isobenzofuran: H. Takeshita, Y. Wada, A. Mori, and T. Hatsui, *Chem. Lett.*, 335 (1973).

Thermal Rearrangements of (Substituted allyl)dialkyl-2H-pyrroles

John M. Patterson,* J. Douglas Ferry, Jan W. deHaan, and Michael R. Boyd

Contribution from the Department of Chemistry, University of Kentucky, Lexington, Kentucky 40506. Received September 7, 1974

Abstract: The thermal behavior of simple substituted 2H-pyrroles, which are contained in tetrahydrocorrin systems and are intermediates in trialkylpyrrole isomerizations, was examined to elucidate possible competitive [1,5] and [3,3] sigmatropic rearrangement paths. The major path followed in the rearrangement of (substituted allyl)dialkyl-2H-pyrroles depends upon the substitution pattern of the migrating allyl group. Crotyl substituents migrate by a [1,5] shift (without inversion) from the 2- to the 3-position of the ring. On the other hand, the α -methylallyl group migrates by competitive [3,3] shifts (with inversion): migration from the 2- to the 5-position and migration from the 2- to the 4-position. Of the two migrations, the former is the more rapid. The transition state in the more rapid migration presumably possesses the more stable 2H-pyrrole configuration as compared with the latter path which contains the less stable 3H-pyrrole configuration.

N-(Substituted allyl)pyrroles undergo thermal isomerization involving competitive [1,5] and [3,3] sigmatropic shifts of the allyl group.¹ When the 2- and 5-positions of the pyrrole ring are substituted with alkyl groups, allylic migration to the 3-position of the pyrrole ring occurs with both inversion and noninversion of the allyl substituent. The N- to 3-migration without inversion suggests that this competitive rearrangement path proceeds by consecutive [1,5] shifts involving 2H-pyrrole intermediates. Evidence for the participation of such 2H-pyrrole intermediates in the thermal isomerization of other trisubstituted pyrroles has been reported,² and it has been found that these 2H-pyrrole species (2-benzyl-2,5-dimethyl-2H-pyrrole) undergo a facile isomerization to the corresponding 3-isomer (3-benzyl-2,5-dimethylpyrrole).³ An investigation of possible competitive [1,5] and [3,3] sigmatropic shifts of allyl groups in the nickel tetrahydrocorrin system (contains a 2H-pyrrole species) by Grigg and coworkers⁴ showed that the 3-methyl-2-butenyl group migrated without inversion. However, because of low conversions (*ca.* 2%) it was not possible to exclude other competing paths. In order to elucidate the migration modes of substituted allyl groups in 2H-pyrrole systems, we have investigated the pyrolytic behavior of (substituted allyl)dialkyl-2H-pyrroles.

Results and Discussion

Initial pyrolyses with 2- α -methylallyl- (**1a**) and 2-crotyl-2,5-dimethyl-2H-pyrrole (**1b**) at 180° showed that the major (*ca.* 90%) thermolysis product in either experiment was 3-crotyl-2,5-dimethylpyrrole (**2b**). While these results

a, X = α -methylallyl; b, X = *trans*-crotyl; c, X = *cis*-crotyl

suggested that, as one possibility, the 2-methylallyl group migrated from the 2- to 3-position with inversion, such 2- to

3-migrations with inversion were considered to be unlikely in the pyrolysis of 2- α -methylallylpyrrole.¹ In addition, nmr studies of the isomerization of 2-crotyl-2,5-dimethyl-2H-pyrrole (**1b**) at 100° indicated that the compound rearranged predominately to the 3-crotyl compound (without inversion), while, on the other hand, the pyrolysis of 2- α -methylallyl-2,5-dimethyl-2H-pyrrole (**1a**) produced initially a 2-crotyl-2,5-dimethyl-2H-pyrrole (**1b**) which on further heating was converted to the 3-crotyl-2,5-dimethylpyrrole (**2b**). The formation of the 3-crotyl isomer in the latter experiment conceivably could have arisen from one or from combinations of the following paths: an across-the-ring migration of the α -methylallyl substituent from the 2- to 5-position with inversion followed by a shift of the crotyl group to the 4-position without inversion (eq 1); an inversion of the α -methylallyl substituent at the 2-position followed by a 2 to 3 shift of the crotyl group (eq 2); or an

Table I. Glpc Analysis^a of 2*H*-Pyrrole Pyrolyses Followed by Nmr

2 <i>H</i> -Pyrrole	<i>T</i> , °C	3-Crotylmethylethylpyrroles, ^b %			
		2-Methyl-5-ethyl		5-Methyl-2-ethyl	
		Trans (5b)	Cis (5c)	Trans (6b)	Cis (6c)
3a	70			91 ^c	3
4a	70	91 ^{d,e}	3	1	
3b	100	73 ^{d,f}	2	4	
4c	100		3	11	60 ^{g,h}

^a Values reported are area %. ^b Identifications of minor components are based on glpc retention times. ^c Three additional unidentified products of 1, 1, and 4% were present. ^d **5b** is not separable from **6c** by glpc; however, the infrared spectrum showed no cis absorption. ^e Two additional unidentified products of 1 and 2% were present. ^f Unisomerized 2*H*-pyrrole (13%), four unidentified products of 1% each, and one unidentified product (2%) were present. ^g **6c** is not separable from **5b** by glpc; however, the infrared spectrum showed no trans absorption. ^h Unisomerized 2*H*-pyrrole (4%) and five unidentified products of 1, 2, 1, 11, and 12% were present.

across-the-ring migration of the α -methylallyl group from the 2- to 4-position with inversion (eq 3).

The major isomerization paths followed were identified by carrying out the pyrolyses of 2- α -methylallyl-2-methyl-5-ethyl-2*H*-pyrrole (**3a**), 2- α -methylallyl-5-methyl-2-ethyl-2*H*-pyrrole (**4a**), 2-*trans*-crotyl-2-methyl-5-ethyl-2*H*-pyrrole (**3b**), and 2-*cis*-crotyl-5-methyl-2-ethyl-2*H*-pyrrole (**4c**), following the isomerizations by nmr and analyzing the final product mixture by glpc (see Table I).

The nmr spectrum of **3a**, which consists of absorptions of a pair of diastereomers,⁵ underwent substantial change after heating at 70° for 2 hr (see Figure 1). In addition to a reduction in intensities of original absorptions, a new triplet, multiplet, and singlet appeared at 0.68, 1.63, and 2.18 ppm, respectively. The two pairs of doublets in the ranges of 6.11–6.21 and 7.03–7.20 ppm became multiplets. These new absorptions correspond to those exhibited by 2-crotyl-5-methyl-2-ethyl-2*H*-pyrrole (**4b**). After 4.5 hr, the nmr spectrum consisted predominately of the spectrum of **4b**. Minor absorptions at 1.13 (t), 1.63 (m), 2.15 (s), 3.00 (m), and 5.45 (m) ppm are attributed to the presence of a small amount of 3-*trans*-crotyl-5-methyl-2-ethylpyrrole (**6b**) and after a heating period of 19.5 hr the spectrum became identical with that of **6b**. Decoupling experiments⁵ verified the structural assignment. The infrared spectrum of **6b** was identical with that obtained from an authentic sample and showed trans absorption (970 cm⁻¹) but no cis. The changes are summarized in eq 4.

Figure 1. Pyrolysis of **3a**.

Similar results were observed in the pyrolysis of **4a** at 70°. Isomerization initially to 2-*trans*-crotyl-2-methyl-5-ethyl-2*H*-pyrrole (**3b**) was observed followed by conversion to 3-*trans*-crotyl-2-methyl-5-ethylpyrrole (**5b**) on extended heating. The infrared spectrum of the pyrolyzate was identical with an authentic sample of **5b** and showed no cis absorption. The intermediate **3b** was synthesized⁵ and pyro-

lyzed at 100°. Isomerization occurred producing mainly **5b** (Table I) with no evidence (nmr) of a new 2*H*-pyrrole or of α -methylallyl absorptions.

These data exclude the conversion of **1a** to **1b** by path 2 (eq 2) and show that the competitive [1,5] and [3,3] sigmatropic migrations of substituted allyl groups in 2*H*-pyrroles are highly sensitive to structural features of the migrating groups. The crotyl substituent rearranges predominately⁶ by a [1,5] sigmatropic shift (without inversion), while the

Table II. Thermal Rearrangement of **1a** and **1b** at 201.0°

Residence time, sec	2 <i>H</i> -Pyrrole pyrolyzed	Products, %					
		1a	1b	1c	2a	2b	2c
33.2	1a	61.5	21.5	6.2	1.5	8.9	0.5
28.2	1b	1.6	92.1		0.5	5.8	

α -methylallyl group migrates by competitive [3,3] shifts (eq 1 and 3). A comparison of the rates of production of **4b** and **6b** from **3a** in the nmr pyrolyses (see Figure 1) indicates that the formation of **4b** has progressed to a considerable extent before detectable quantities of **6b** appear. This suggests that [3,3] shifts involving positions 2 and 5 are somewhat more rapid than [3,3] shifts involving positions 2 and 4. Additional support for the 2 to 5 [3,3] shift being the faster process was obtained from relative rate data derived from the pyrolysis of **1a** and **1b** at 201° (Tables II and III).

In the pyrolysis of **1a**, product **2b** arises from the parallel pathways outlined in eq 1 and 3. The formation of **2b** by path 1 (eq 1) is the result of the consecutive conversions of **1a** to **1b** and **1b** to **2b**. Approximate rate constants for each of these conversions were estimated from the data obtained from the pyrolysis of **1a** and **1b** (see Table II), and the concentration of **2b** formed by the consecutive reactions in path 1 was calculated in the usual way.⁷ The concentration of **2b** arising from path 3 (eq 3) was determined by the difference of the experimental concentration of **2b** (Table II) and the concentration calculated for path 1. Relative rate data for the competing processes in the pyrolyses were derived from these concentrations and are summarized in Table III.

The greater reactivity of **1a** in the 2 to 5 shift (eq 6) as compared with the 2 to 4 shift (eq 7) can be rationalized, if

it is assumed that bond formation has progressed to a considerable extent in the transition state (transition state

Table III. Relative Rates for the Isomerization of **1a** and **1b** at 201.0°

Reaction	Type	Rel rate
1a → 2a	1,5	1
1a → 2b	3,3	5
1a → 1b	3,3	14
1b → 2b	1,5	4
1b → 2a	3,3	0.3

structure resembles product structure). Since 2*H*-pyrroles had been found to be more stable than 3*H*-pyrroles in thermolysis reactions,⁸ the transition state involving 2- to 5-migrations is the more stable one.

The chair-like transition state previously described for the isomerization of allylpyrroles¹ appears to be valid in the 2*H*-pyrrole system also. In the pyrolysis of **1a**, *trans*-crotyl product is produced at a faster rate than is *cis*-crotyl product.

Experimental Section

The synthesis, separation, and characterization of starting materials used and pyrolysis products encountered in these experiments have been described previously.⁵

Nmr spectra were measured on degassed CCl₄ solutions (*ca.* 10%) using HMDS as the lock signal and TMS as internal standard. The samples were heated to 70 or 100° either in the spectrometer or externally, and runs were made using both Varian HA-60-IL and T-60 spectrometers. Glpc analyses were carried out on a Hewlett-Packard 5750 gas chromatograph using a 6 ft X 0.125 in. Carbowax 20M HP high-efficiency packed column at 140°.

Pyrolyses were carried out in the apparatus previously described¹ with the reactor and gas chromatograph in series. Product analyses were run on the Carbowax 20M column at 115° for low boiling products and at 160° for the higher boiling products. The rate of disappearance of 2- α -methylallyl-2,5-dimethyl-2*H*-pyrrole was followed at 182°, and a plot of $\ln C_0/C$ vs. residence time yielded a straight line.

Acknowledgments. This research was supported in part by grants from the U.S. Army Research Office-Durham, from the U.S. Atomic Energy Commission, and from the University of Kentucky Faculty Research Fund. One of us (J.d.H.) thanks Professor A. I. M. Keulemans for the leave of absence.

References and Notes

- (1) J. M. Patterson, J. W. deHaan, M. R. Boyd, and J. D. Ferry, *J. Amer. Chem. Soc.*, **94**, 2487 (1972).
- (2) J. M. Patterson and S. Soedigdo, *J. Org. Chem.*, **33**, 2057 (1968).
- (3) J. M. Patterson and L. T. Burka, *Tetrahedron Lett.*, 2215 (1969).
- (4) R. Grigg, A. W. Johnson, and K. W. Shelton, *J. Chem. Soc. C*, 1291 (1968).
- (5) J. M. Patterson, J. D. Ferry, and M. R. Boyd, *J. Amer. Chem. Soc.*, **95**, 4356 (1973).
- (6) In the migration of the *trans*-crotyl group in **3b**, inversion could not have occurred to an extent greater than 2% since this was the largest yield of an unidentified component (a possible α -methylallylpyrrole).
- (7) A. A. Frost and R. G. Pearson, "Kinetics and Mechanism," Wiley, New York, N.Y., 1953, p 153.
- (8) J. L. Wong, M. H. Ritchie, and C. M. Gladstone, *Chem. Commun.*, 1093 (1971).