Side-on versus end-on coordination of dinitrogen to titanium(II) and mixed-valence titanium(I)/titanium(II) amido complexes

Citation for published version (APA):

Duchateau, R., Gambarotta, S., Beydoun, N., & Bensimon, C. (1991). Side-on versus end-on coordination of dinitrogen to titanium(II) and mixed-valence titanium(I)/titanium(II) amido complexes. *Journal of the American Chemical Society*, 113(23), 8986-8988. <https://doi.org/10.1021/ja00023a080>

DOI:

[10.1021/ja00023a080](https://doi.org/10.1021/ja00023a080)

Document status and date:

Published: 01/01/1991

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Figure 2. A plot of $1/\phi_Y$ versus $1/[\text{pyridine}]$, see text.

attributed to ylide 3. The spectrum of 3 is similar to those of the other known dialkylpyridine ylides but is somewhat shifted to the blue (364 versus 390–410 nm) of the homocubyl⁵ and adamantyl⁴ ylides.

The optical yield of ylide 3 is reduced by the presence of oxygen, a selective trap of triplet carbenes.^{1,8} From this result two conclusions are possible. The singlet and triplet states of 2 may be of comparable energy, as per aryl- and arylalkylcarbenes, and are in rapid equilibrium.¹ Alternatively a referee has suggested that the excited state of $^1\text{I}^*$ may undergo intersystem crossing to form $^3\text{I}^*$ which extrudes nitrogen to form the triplet carbene.

The intensity of the ylide signal ($\phi_Y = \Delta A_{364\text{nm}}$) produced in a laser pulse increases steadily as the concentration of pyridine increases until $[\text{PYR}] \approx 1.5 \text{ M}$, at which point the optical yield is saturated. A double reciprocal plot of the data (Figure 2) is linear with a slope of $0.45 = k_T/k_{\text{PYR}}K$, where k_{PYR} is the second-order rate constant for reaction of dimethylcarbene with pyridine, K is the equilibrium constant ($K = [\text{2S}]/[\text{2T}] \ll 1$, assuming the triplet is the ground state as per methylene and that singlet-triplet equilibration is rapid),⁹ and k_T is the sum of all first-order and pseudo-first-order rate constants of all processes which consume the carbene in the absence of pyridine. These processes include isomerization ($k_{1,2}$) and reaction of the carbene with solvent (k_{RH}^1 and k_{RH}^3). Assuming that $k_{\text{PYR}}K \approx 10^7\text{--}10^9 \text{ s}^{-1}$ ⁸ pegs the lifetime of the carbene at 1–100 ns in pentane in the absence of pyridine. This lifetime must be controlled in part by reaction with solvent because there is a solvent deuterium isotope effect of 1.3 on the slope of plots such as that in Figure 2 in CH_3CN vs CD_3CN and CHCl_3 vs CDCl_3 .

The data convincingly demonstrate that dimethylcarbene is a true reactive intermediate, a species which exists in a potential energy minimum, and has a finite lifetime in solution.

Acknowledgment. Support of this work by the National Science Foundation (CHE-881950) is gratefully acknowledged.

(8) Sugiyama, M.; Celebi, S.; Platz, M. S. *J. Am. Chem. Soc.* Submitted for publication. The singlet and triplet states of methylene are not in rapid equilibrium in solution;¹⁰ their energy separation is too large.¹¹

(9) Scheme 1 is a conventional¹ interpretation of singlet-triplet phenomena in carbene chemistry. However, it has been pointed out (Griller, D.; Nazran, A. S.; Scaiano, J. J. *Am. Chem. Soc.* 1984, 106, 198) that the available data for aryl carbenes is consistent with the presence of a single reactive intermediate, one that can undergo reactions normally ascribed to either the triplet or the singlet state of the carbene. In this interpretation a triplet carbene can react with pyridine to form an ylide via a singlet-triplet surface crossing along the reaction coordinate. The surface crossing mechanism resolves inconsistencies in the kinetics of reaction of triplet diphenylcarbene and methanol. The data of this work is consistent with either a preequilibrium or surface crossing mechanism. The traditional mechanism is presented in Scheme 1 because there is no compelling reason to discard it with dimethylcarbene.

(10) Turro, N. J.; Cha, Y.; Gould, I. R. *J. Am. Chem. Soc.* 1987, 109, 2101.

(11) Leopold, D. G.; Murray, K. K.; Miller, A. E. S.; Lineberger, W. C. *J. Chem. Phys.* 1984, 83, 4849.

Side-On versus End-On Coordination of Dinitrogen to Titanium(II) and Mixed-Valence Titanium(I)/Titanium(II) Amido Complexes

Robbert Duchateau, Sandro Gambarotta,* Nohma Beydoun, and Corinne Bensimon

Department of Chemistry
University of Ottawa, Ottawa
Ontario K1N 6N5, Canada

Received May 29, 1991

Dinitrogen coordination¹ and reduction² are among the most challenging of the several transformations (dinitrogen activation,^{1,2} reactivity with unsaturated organic substrates,³ C–H bond activation,⁴ CO chemistry,⁵ and Ziegler–Natta catalysis⁶) promoted by divalent titanium compounds. Although the great ability of titanocene systems to give dinitrogen fixation with a variety of bonding modes² clearly underlines the strength of divalent titanium as a powerful reducing agent; however, the coordination chemistry of the oxidation state +II remains limited to only a few diverse cases.^{7,8} To date there are no clearly demonstrated examples where fixation/reduction of dinitrogen occurs on a nonmetallocenic titanium system.

Herein we report the isolation and characterization of two novel dinitrogen Ti(II) and mixed-valence Ti(I)/Ti(II) amido complexes obtained via chlorine replacement at *trans*-(TMEDA)₂TiCl₂^{7e} [TMEDA = *N,N,N',N'*-tetramethylethylenediamine].

The reaction of *trans*-(TMEDA)₂TiCl₂ with 1 equiv of (Me₃Si)₂NLi in toluene under nitrogen at –80 °C formed a purple jelly suspension which turned into a deep-brown solution upon standing overnight at room temperature. Brown crystals of [(Me₃Si)₂N]TiCl(TMEDA)₂(μ-N₂) (1) were obtained after evaporation and crystallization from ether at –30 °C (Scheme 1).⁹ Occasionally, small amounts of colorless crystals of mo-

(1) (a) Sanner, R. D.; Duggon, D. M.; McKenzie, T. C.; Marsch, R. E.; Bercaw, J. E. *J. Am. Chem. Soc.* 1976, 98, 8358. (b) Pez, G. P.; Armor, N. J. *Adv. Organomet. Chem.* 1981, 19, 1. (c) Pez, G. P.; Appag, P.; Crissey, R. K. *J. Am. Chem. Soc.* 1982, 104, 482. (d) Berry, D. H.; Procopio, L. J.; Carroll, P. J. *Organometallics* 1988, 7, 570. (e) Flamini, A.; Cole-Hamilton, D. J.; Wilkinson, G. J. *Chem. Soc., Dalton Trans.* 1978, 454. (f) Busby, R.; Klotzbucher, W.; Ozin, G. A. *Inorg. Chem.* 1977, 16, 822.

(2) (a) Van Tamelen, E. *Acc. Chem. Res.* 1970, 3, 361. (b) Volpin, M. E.; Shur, V. B.; Kudryavtsev, R. V.; Prodayko, L. A. *J. Chem. Soc., Chem. Commun.* 1968, 1038. (c) Sobota, P.; Janas, Z. *Inorg. Chim. Acta* 1981, 53, L11.

(3) (a) Cohen, S. A.; Auburn, P.; Bercaw, J. E. *J. Am. Chem. Soc.* 1983, 105, 1136. (b) Demersman, B.; Coupance, P. L.; Dixneuf, P. U. *J. Organomet. Chem.* 1985, 297, C35. (c) Cohen, S. A.; Bercaw, J. E. *Organometallics* 1985, 4, 1006. (d) Sekutowski, D. G.; Stucky, G. D. *J. Am. Chem. Soc.* 1976, 98, 1376. (e) Alt, H. G.; Engelhardt, H. E.; Rausch, M. D.; Kool, L. B. *J. Am. Chem. Soc.* 1985, 107, 3717. (f) Famili, A.; Farona, M. F.; Thanedar, S. J. *Chem. Soc., Chem. Commun.* 1983, 435. (g) Green, M. L. H.; Hazel, N. J.; Grebenik, P. D.; Mtetwa, V. S. B.; Prout, K. J. *Chem. Soc., Chem. Commun.* 1983, 356. (h) Thewalt, U.; Stollmaier, F. *J. Organomet. Chem.* 1982, 228, 149.

(4) Bercaw, J. E.; Marvich, R. H.; Bell, L. G.; Britzinger, H. H. *J. Am. Chem. Soc.* 1972, 94, 1219.

(5) (a) Kool, L. B.; Rausch, M. D.; Alt, H. G.; Herbershold, M.; Wolf, B.; Thewalt, U. *J. Organomet. Chem.* 1985, 297, 159. (b) Frerichs, S. R.; Stein, B. K.; Ellis, J. E. *J. Am. Chem. Soc.* 1987, 109, 5558.

(6) (a) Sinn, H.; Kaminsky, W. *Adv. Organomet. Chem.* 1980, 18, 99. (b) Boor, J. *Ziegler–Natta Catalysts and Polymerization*; Academic Press: New York, 1979.

(7) Gavens, P. D.; Bottrill, M.; Kelland, J. W.; McMeeking, J. *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. G. A., Abel, E. W., Eds.; Pergamon Press: New York, 1982.

(8) (a) Jensen, J. A.; Wilson, S. R.; Schultz, A. J.; Girolami, G. S. *J. Am. Chem. Soc.* 1987, 109, 8094. (b) Morris, R. J.; Girolami, G. S. *Inorg. Chem.* 1990, 29, 4169. (c) Girolami, G. S.; Wilkinson, G.; Galas, A. M. R.; Thornton-Pett, M.; Hursthouse, M. B. *J. Chem. Soc., Dalton Trans.* 1985, 1339. (d) Durfee, L. D.; Fanwick, P. E.; Rothwell, I. P.; Folting, K.; Huffman, J. C. *J. Am. Chem. Soc.* 1987, 109, 4720. (e) Durfee, L. D.; Hill, J. E.; Fanwick, P. E.; Rothwell, I. P. *Organometallics* 1990, 9, 75. (f) Edema, J. J. H.; Duchateau, R.; Gambarotta, S.; Hynes, R.; Gabe, E. *Inorg. Chem.* 1991, 30, 156. (g) Edema, J. J. H.; Duchateau, R.; Gambarotta, S.; Bensimon, C. *Inorg. Chem.*, in press.

Scheme I

nomeric $[(\text{Me}_3\text{Si})_2\text{N}]_2\text{TiCl}_2[(\text{TMEDA})_2\text{Li}]$ (**2**)¹⁰ were isolated upon concentration of the mother liquor and further freezing. Conversely, reaction with 2.5 equiv of $(\text{Me}_3\text{Si})_2\text{NLi}$ in the presence of a small excess of TMEDA led, under the same reaction conditions, to the formation of white crystals of $[(\text{Me}_3\text{Si})_2\text{N}]_2\text{TiCl}_2[(\text{TMEDA})_2\text{Li}]$ (**2**) and deep-purple/black

(9) Complex **1**: solid $(\text{Me}_3\text{Si})_2\text{NLi}$ (2.95 g, 17.7 mmol) was added to a suspension of $\text{TiCl}_2(\text{TMEDA})_2$ (6.2 g, 17.6 mmol) in toluene (75 mL) under nitrogen atmosphere at -80°C . The reaction mixture was allowed to slowly warm up to -40°C originating a thick purple jelly suspension which became a deep brown solution upon standing at room temperature. After 3 days at room temperature, the solution was evaporated to dryness, and the residual solid redissolved in ether. Deep brown crystals of **1** were obtained upon cooling at -30°C (1.4 g, 1.9 mmol, 11% yield based on titanium). IR [KBr, Nujol mull, cm^{-1}] 2920 (s), 1460 (s), 1380 (m), 1360 (w), 1295 (w), 1290 (m), 1260 (s), 1240 (s), 1195 (w), 1170 (w), 1125 (m), 1100 (w), 1070 (m), 1050 (m), 1020 (m), 1010 (m), 100 (sh), 950 (s), 840 (s), 810 (s), 770 (s), 760 (sh), 750 (sh), 695 (s), 655 (s), 620 (s), 610 (sh), 505 (m), 480 (w), 460 (w), 430 (m), 415 (m), 395 (m), 380 (w); ^1H NMR [C_6D_6 , 200 MHz, 23°C] δ 3.48, 2.80, 2.15, 1.90 (s, 16 H, TMEDA), 0.68 (s, 18 H, Me_3Si); ^{13}C NMR [C_6D_6 , 75 MHz, 25°C] δ 58.33, 57.89 (CH₂, TMEDA), 53.47, 51.73, 49.20 (CH₃, TMEDA), 7.51 (CH₃, Me_3Si). Anal. Calcd (found) for $\text{C}_{24}\text{H}_{68}\text{N}_6\text{Si}_4\text{Cl}_2\text{Ti}_2$: C 38.54 (38.33), H 9.17 (9.09), N 14.98 (14.76), Ti 12.80 (12.71). Complex **2**, method A: solid $(\text{Me}_3\text{Si})_2\text{NLi}$ (2.45 g, 14.7 mmol) was added at -80°C to a suspension of $\text{TiCl}_2(\text{TMEDA})_2$ (2.6 g, 7.4 mmol) in THF (75 mL) containing TMEDA (5 mL) under nitrogen atmosphere. The reaction mixture was kept at -40°C for 2 h and then allowed to slowly warm up to room temperature, originating a light brown solution. After evaporation to dryness and crystallization from ether, white crystals of **2** were obtained (2.0 g, 2.9 mmol, 39% yield): IR [KBr, Nujol mull, cm^{-1}] ν 2900 (s), 2800 (m), 1460 (s), 1380 (m), 1360 (m), 1290 (s), 1260 (sh), 1240 (s), 1190 (m), 1170 (m), 1140 (m), 1105 (m), 1070 (m), 1030 (s), 1015 (s), 930 (s), 890 (s), 835 (s), 785 (s), 760 (m), 720 (w), 695 (s), 670 (s), 630 (m), 615 (w), 500 (m), 450 (m), 410 (m), 390 (m). Anal. Calcd (found) for $\text{C}_{24}\text{H}_{68}\text{N}_6\text{Si}_4\text{Cl}_2\text{Ti}_2$: C 42.46 (42.38), H 10.10 (10.00), N 12.38 (12.33), Ti 7.05 (7.01). Method B: A suspension of $\text{TiCl}_3(\text{THF})_3$ (20 g, 54 mmol) in THF (200 mL) was treated with TMEDA (17 mL). The addition of $\text{Me}_3\text{Si}_2\text{NLi}$ (18.1 g, 108 mmol) to the resulting light blue solution turned the color light green. After standing 30 min at room temperature the solvent was evaporated in vacuo. The residual solid was redissolved in ether (500 mL), and insoluble material filtered out. Colorless crystal of **2** (21.9 g, 32 mmol, 59%) separated upon concentration and standing at -30°C overnight. Complex **3**: Solid $(\text{Me}_3\text{Si})_2\text{NLi}$ (3.40 g, 20.6 mmol) was added at -80°C to a suspension of $\text{TiCl}_2(\text{TMEDA})_2$ (2.4 g, 6.8 mmol) in toluene (100 mL) containing TMEDA (3 mL) under argon atmosphere. The suspension slowly originated a jelly purple suspension which turned into a brown solution upon standing overnight at room temperature. The solvent was removed in vacuo, and the residual solid redissolved in ether. The first fraction of white crystalline material, which separated upon cooling at -30°C was identified as complex **2** by comparison with the IR spectrum of an analytically pure sample (0.9 g, 1.3 mmol, 19%). Further concentration and freezing gave black crystals of **3** (2.5 g, 2.4 mmol, 70% based on titanium): IR [KBr, Nujol mull, cm^{-1}] ν 2900 (s), 2800 (m), 1460 (s), 1380 (m), 1360 (m), 1290 (s), 1240 (s), 1190 (m), 1170 (m), 1140 (m), 1105 (m), 1070 (m), 1030 (s), 1015 (s), 930 (s), 890 (s), 785 (s), 760 (m), 720 (w), 695 (s), 670 (s), 630 (m), 615 (w), 500 (m), 450 (w), 410 (m), 390 (m). Anal. Calcd (found) for $\text{C}_{36}\text{H}_{104}\text{N}_{12}\text{Si}_8\text{Ti}_2$: C 41.87 (41.78), H 10.15 (10.12), N 16.28 (16.23), Ti 9.27 (9.14).

(10) Triclinic, $P1$, $a = 13.090$ (4) Å, $b = 14.683$ (4) Å, $c = 11.591$ (1) Å, $\alpha = 102.09$ (1)°, $\beta = 91.02$ (1)°, $\gamma = 92.80$ (2)°, $V = 2174.8$ (9) Å³, $Z = 2$, $R = 0.078$ ($R_w = 0.083$), $T = 298$ K for 344 parameters and 3506 significant reflections out of 6425. All non-hydrogen atoms were located and anisotropically refined. Hydrogen atom positions were calculated but not refined. Duchateau, R.; Gambarotta, S.; Bensimon, C., manuscript in preparation.

Figure 1. ORTEP drawing of **1** showing the labeling scheme. Thermal ellipsoids are drawn at the 50% probability level. Selected values of bond distances (Å) and angles (deg) are as follows: Ti(1)–Cl(1) = 2.388 (3), Ti(1)–N(1) = 2.023 (5), Ti(1)–N(2) = 2.352 (6), Ti(1)–N(3) = 2.351 (6), Ti(1)–N(4) = 1.762 (5), N(1)–Si(1) = 1.714 (6), N(4)–N(4a) = 1.289 (9), Cl(1)–Ti(1)–N(1) = 96.6 (2), N(1)–Ti(1)–N(4) = 111.3 (2), Ti(1)–N(4)–N(4a) = 168.5 (2), Ti(1)–N(1)–Si(1) = 121.2 (3).

crystals of $\{[(\text{Me}_3\text{Si})_2\text{N}]_2\text{Ti}\}_2(\mu\text{-}\eta^2\text{-}\eta^2\text{-N}_2)_2[(\text{TMEDA})_2\text{Li}]$ (**3**). The two species were isolated in moderate yield and analytically pure form,⁹ upon solvent evaporation and fractional crystallization in ether at -30°C (Scheme I). In both cases a competitive reaction is probably responsible for the formation of **2** since complexes **1** and **3** are thermally robust and did not show any appreciable decomposition in benzene at -78°C . Furthermore, **2** was the only identifiable product when the two reactions were carried out in THF. Compound **2** could conveniently be prepared on a very large scale by direct reaction of $\text{TiCl}_3(\text{THF})_3$ with 2 equiv of R_2NLi and 2 equiv of TMEDA in ether.⁹

The structures of **1** and **3** have been determined by X-ray analysis.¹¹ Complex **1** is binuclear and consists of two identical $\{[(\text{Me}_3\text{Si})_2\text{N}]_2\text{TiCl}(\text{TMEDA})\}$ fragments each bonded to one of the two nitrogen atoms of a bridging dinitrogen moiety (Figure 1). The two titanium fragments are rotated with respect to each other [torsion angle Cl(1)–Ti(1)–Ti(1a)–Cl(1a) = 48.7 (5)°] probably as a result of minimization of steric repulsion between the two bulky amido groups. The coordination geometry of the titanium atom is square-pyramidal with the titanium atom quite elevated above the basal plane [distance from the plane 0.466 (3) Å] defined by chlorine and the nitrogen atoms of TMEDA and amido groups [N(1)–Ti(1)–N(3) = 92.0 (2)°, N(3)–Ti(1)–N(2) = 76.2 (2)°, N(2)–Ti(1)–Cl(1) = 84.9 (2)°, Cl(1)–Ti(1)–N(1) = 96.6 (2)°]. The apical position of the square-pyramid is occupied by one of the two atoms of the bridging dinitrogen molecule. The remarkably short Ti–N distance [Ti(1)–N(4) = 1.762 (5) Å] together with the almost linear arrangement of the Ti–($\mu\text{-}\eta^2\text{-}\eta^2\text{-N}_2$)–Ti moiety [Ti(1)–N(4)–N(4a) = 168.5 (2)°] and the rather long N–N distance [N(4)–N(4a) = 1.289 (9) Å] are in agreement with the presence of considerable electronic delocalization over the $\text{Ti}_2(\mu\text{-N}_2)$ moiety. The Ti–N distance with the trigonal-planar amido group [Ti(1)–N(1) = 2.023 (5) Å] is significantly shorter than those formed by the nitrogen atoms of the TMEDA groups [Ti(1)–N(2) = 2.352 (6) Å, Ti(1)–N(3) = 2.351 (6) Å] suggesting the presence of some π -bond character with both silicon and titanium atoms.

(11) Complex **1**: tetragonal, $P4_2/2_1$, $a = 11.500$ (5) Å, $c = 31.644$ (9) Å, $V = 4185$ (3) Å³, $Z = 4$, $R = 0.049$ ($R_w = 0.059$), $T = 296$ K for 182 parameters and 1672 significant reflections out of 2244. All non-hydrogen atoms were anisotropically refined. Hydrogen atoms were introduced at calculated positions but not refined. Complex **3**: orthorhombic, $I22_2$, $a = 13.240$ (5) Å, $b = 22.381$ (3) Å, $c = 11.575$ (1) Å, $V = 3430.0$ (13) Å³, $Z = 2$, $R = 0.077$ ($R_w = 0.088$), $T = 298$ K for 133 parameters and 1058 significant reflections out of 1447. All non-hydrogen atoms were anisotropically refined. Hydrogen atoms were introduced at calculated positions but not refined. One of the two atoms of the dinitrogen molecules was found disordered over two positions. Disorder was modeled by assigning an occupancy of 50% to one of the two nitrogen atoms.

Figure 2. ORTEP drawing of **3** showing the anionic moiety with the labeling scheme. Thermal ellipsoids are drawn at the 50% probability level. Selected values of bond distances (Å) and angles (deg) are as follows: Ti(1)–N(1) = 2.066 (13), Ti(1)–N(3) = 2.236 (19), Ti(1)–N(4) = 2.290 (13), N(1)–Si(1) = 1.726 (14), Ti(1)–Ti(1a) = 3.680 (8), N(3)–N(4) = 1.379 (21), N(1)–Ti(1)–Ni(1a) = 107.0 (5), Ti(1)–N(1)–Si(1) = 123.0 (8).

The chemical connectivity of **3** was also demonstrated by X-ray diffraction analysis. The structure consists of two separate $[\text{Li}(\text{TMEDA})_2]$ and $[\{(\text{Me}_3\text{Si})_2\text{N}\}_2\text{Ti}\}_2(\mu-\eta^2-\eta^2-\text{N}_2)_2]$ ionic fragments. The Ti-containing unit is dinuclear and is formed by two $[(\text{Me}_3\text{Si})_2\text{N}]_2\text{Ti}$ groups symmetrically and perpendicularly placed on the two sides of the plane defined by two parallel molecules of dinitrogen (Figure 2). Four of the six coordination sites of titanium are occupied by the four nitrogen atoms of the two coplanar dinitrogen molecules, the two remaining sites being occupied by the two amido groups. The side-on geometry of the two coordinated N_2 molecules is somewhat reminiscent of that of a recently reported zirconium dinitrogen phosphine complex.¹² Even in this case, the N–N distance [N(3)–N(4) = 1.379 (21) Å] is rather long, and it is significantly longer than that displayed by complex **1**. By contrast, the Ti–N distances [Ti(1)–N(3) = 2.236 (19) Å, Ti(1)–N(4) = 2.290 (13) Å] are definitely longer as a probable result of the different fashion of bonding. It is quite difficult to speculate on the extent of dinitrogen reduction on the exclusive basis of the N–N distances, since the structural features of **3** (with no particularly long N–N and Ti–N distances) are somehow in contradiction with those of **1** (with comparable N–N distance and extremely short Ti–N distance). The completely different bonding mode of dinitrogen in the two complexes (end-on versus side-on) is intriguing indeed and at the moment can be explained only by the different steric bulk of the two complexes.

As expected, complex **1** is diamagnetic while **2** and **3** are paramagnetic [μ_{eff} = 1.75 μ_{B} , μ_{eff} = 1.37 μ_{B} , respectively]. The low value of the magnetic moment of **3** can be ascribed to either antiferromagnetic or superexchange, since the Ti...Ti nonbonding distance is rather short [Ti(1)...Ti(1a) = 3.680 (8) Å].

Acknowledgment. This work was supported by the Natural Sciences and Engineering Research Council of Canada (operating grant) and the Petroleum Research Funds administered by the American Chemical Society.

Supplementary Material Available: Tables listing atomic positional parameters, anisotropic thermal parameters, and complete bond distances and angles for **1** and **3**, and an ORTEP plot for $\text{Li}(\text{TMEDA})_2$ (15 pages); tables of observed and calculated structure factors for **1** and **3** (21 pages). Ordering information is given on any current masthead page.

Photoluminescence of Antimony(III) and Bismuth(III) Chloride Complexes in Solution

Hans Nikol and Arnd Vogler*

Institut für Anorganische Chemie
Universität Regensburg, Universitätsstrasse 31
D-8400 Regensburg, Federal Republic of Germany

Received May 28, 1991

Revised Manuscript Received September 17, 1991

Metal-centered (MC) excited states play an important role in the photophysics and photochemistry of coordination compounds.¹ However, the study of MC states has been essentially limited to d and f block elements, while the main group metals were largely ignored.² This lack of knowledge is quite surprising since MC excited states of many main group metal complexes can be easily studied by emission spectroscopy under ambient conditions. Although simple halide complexes of s^2 metals such as Ti^+ , Sn^{2+} , Pb^{2+} , and Sb^{3+} are known to be luminescent in solution,² this observation has been occasionally used only for analytical applications but hardly for the characterization of the emitting excited state. On the contrary, the emission of s^2 ions which are doped into host lattices has been studied in detail by solid-state physicists.^{3,4} We report here our observations on the photoluminescence of SbCl_4^- , SbCl_6^{3-} , BiCl_4^- , and BiCl_6^{3-} in solution and discuss the nature of the emitting excited state. While an emission of SbCl_6^{3-} in CHCl_3 has been reported before,^{5,6} the other three complex ions are not yet known to show photoluminescence in solution at room temperature.

The compounds $[\text{NET}_4]\text{SbCl}_4$ and $[\text{NET}_4]\text{BiCl}_4$ were prepared according to a published procedure.⁷ The complex ions SbCl_6^{3-} and BiCl_6^{3-} in acetonitrile were formed by the addition of an excess of $[\text{NET}_4]\text{Cl}$ to the MCl_4^- ions. The absorption spectrum of SbCl_4^- in acetonitrile (Figure 1) displays three bands (Table I). Light absorption by SbCl_4^- was accompanied by a red emission (Figure 1, Table I). The excitation spectrum agreed well with the absorption spectrum. Upon addition of chloride SbCl_4^- was converted to SbCl_6^{3-} which has an absorption spectrum (Figure 1, Table I) similar to that of SbCl_4^- . Light absorption by SbCl_6^{3-} led to a green emission (Figure 1, Table I). Again, the excitation spectrum matched the absorption spectrum. If light absorption was complete ($A > 2$), the progressive conversion of SbCl_4^- to SbCl_6^{3-} was accompanied by the appearance of an isosmestic point at $\lambda = 613$ nm. The absorption and emission spectra of BiCl_4^- and BiCl_6^{3-} (Figure 2) showed features (Table I) which are analogous to those of SbCl_4^- and SbCl_6^{3-} . In contrast to the antimony complexes, the B band of the bismuth compounds was not observed in accordance with results on $\text{Cs}_2\text{NaYCl}_6$ doped with Bi^{3+} .⁸

According to the VSEPR model⁹ the structures of complexes of s^2 metal ions should be determined by the presence of a stereochemically active lone pair. However, s^2 complexes with a coordination number of six are generally octahedral, and thus they are exceptions from the rule.^{10,11} This applies also to the structures

(1) (a) Balzani, V.; Carassiti, V. *Photochemistry of Coordination Compounds*; Academic Press: New York, 1970. (b) *Concepts of Inorganic Photochemistry*; Adamson, A. W., Fleischauer, P. D., Eds.; Wiley-Interscience: New York, 1975. (c) Ferraudi, G. J. *Elements of Inorganic Photochemistry*; Wiley-Interscience: New York, 1988.

(2) Vogler, A.; Paukner, A.; Kunkely, H. *Coord. Chem. Rev.* **1990**, *97*, 285.

(3) (a) Blasse, G. *Prog. Solid St. Chem.* **1988**, *18*, 79. (b) Blasse, G. *Rev. Inorg. Chem.* **1983**, *5*, 319.

(4) Ranfagni, A.; Mugnai, D.; Bacci, M.; Viliiani, G.; Fontana, M. P. *Adv. Phys.* **1983**, *32*, 823.

(5) Vogler, A.; Paukner, A. *Inorg. Chim. Acta* **1989**, *163*, 207.

(6) Blasse, G.; Vogler, A. *Inorg. Chim. Acta* **1990**, *170*, 149.

(7) Ahliah, G. Y.; Goldstein, M. J. *Chem. Soc. (A)* **1970**, 326.

(8) Van der Steen, A. C. *Phys. Stat. Sol. (B)* **1980**, *100*, 603.

(9) (a) Gillespie, R. J.; Nyholm, R. S. *Q. Rev. Chem. Soc.* **1957**, *11*, 339. (b) Gillespie, R. J. *Molecular Geometry*; Van Nostrand Reinhold: London, 1972.

(10) Du Bois, A.; Abriel, W. Z. *Naturforsch.* **1990**, *45B*, 573 and references therein.

(12) Fryzuk, M. D.; Haddad, T. S.; Rettig, S. J. *J. Am. Chem. Soc.* **1990**, *112*, 8185.