Solution to Problem 83-8: Gamma function expansion

Citation for published version (APA):

Lossers, O. P. (1984). Solution to Problem 83-8: Gamma function expansion. *SIAM Review*, 26(2), 278-279.
<https://doi.org/10.1137/1026049>

DOI:

[10.1137/1026049](https://doi.org/10.1137/1026049)

Document status and date:

Published: 01/01/1984

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

where

$$\eta = \tan^{-1} [ab / \{(\sqrt{a^2 + 1} - a + 1)a^2 + (\sqrt{a^2 + 1} - a - 1)b^2\}],$$

and

$$\text{Cl}_2(t) = \text{Im Li}_2(e^{it}) = \sum_{n=1}^{\infty} n^{-2} \sin(nt)$$

is Clausen's function.

Also solved by J. H. DAVENPORT (Emanuel College, Cambridge, England), W. B. JORDAN (Scotia, NY), O. P. LOSSERS (Eindhoven University of Technology, the Netherlands) and AVRAM SIDI (Technion, Haifa, Israel). Davenport showed that the integral cannot be expressed as an elementary closed form. Sidi expressed $I(a, b)$ as a rapidly convergent series of functions related to Chebyshev polynomials. Jordan found an effective series for $g(a, b)$.

Gamma Function Expansions

Problem 83-8, by W. B. JORDAN (Scotia, NY).

Prove that

$$(a) \quad \log \frac{\Gamma(z + 1/2)}{\sqrt{z}\Gamma(z)} = -\sum_{r=1}^k \frac{(1 - 2^{-2r})B_{2r}}{r(2r-1)z^{2r-1}} + O(z^{-2k+1/2}),$$

$$(b) \quad \log \frac{\Gamma(z + 3/4)}{\sqrt{z}\Gamma(z + 1/4)} = -\sum_{r=1}^k \frac{E_{2r}}{4r(4z)^{2r}} + O(z^{-2k-1/2}),$$

$$(c)^* \quad \frac{1}{z} \left[\frac{\Gamma(z + 3/4)}{\Gamma(z + 1/4)} \right]^2 = 1 + \frac{2u}{1+} \frac{9u}{1+} \frac{25u}{1+} \frac{49u}{1+} \cdots,$$

where B_{2r} and E_{2r} are Bernoulli and Euler numbers, and $u = 1/64z^2$.

Solution by O. P. LOSSERS (Eindhoven University of Technology, Eindhoven, the Netherlands).

(a), (b) The results immediately follow from Barnes' asymptotic expansion [1, form. 1.18(12)]

$$\begin{aligned} \log \Gamma(z + \alpha) &\sim (z + \alpha - 1/2) \log z - z + 1/2 \log(2\pi) \\ &\quad + \sum_{n=1}^{\infty} \frac{(-1)^{n+1} B_{n+1}(\alpha)}{n(n+1)} z^{-n}, \quad z \rightarrow \infty, \quad |\arg z| < \pi, \end{aligned}$$

in which $B_{n+1}(\alpha)$ is the Bernoulli polynomial, and the known values [1, form. 1.14(7)]

$$B_{n+1}(1/2) - B_{n+1}(0) = -2^{-n}(2^{n+1} - 1)B_{n+1},$$

$$B_{n+1}(3/4) - B_{n+1}(1/4) = 2^{-2n-1}(n+1)E_n;$$

furthermore, recall that B_{n+1} vanishes for even n and E_n vanishes for odd n . The remainder terms may in fact be replaced by $O(z^{-2k-1})$ in case (a), and by $O(z^{-2k-2})$ in case (b).

In the same manner it can be shown that

$$\log \frac{\Gamma(z + \alpha + 1/2)}{\sqrt{z}\Gamma(z + \alpha)} \sim \sum_{n=1}^{\infty} \frac{(-1)^{n+1} E_n(2\alpha)}{2^{n+1}n} z^{-n}, \quad z \rightarrow \infty, \quad |\arg z| < \pi,$$

where $E_n(2\alpha)$ is the Euler polynomial. This expansion includes the results (a) and (b) as special cases.

(c)* The continued fraction is known from Perron [2, p. 36, eq. (24)].

Also solved by V. BELEVITCH (Philips Research Laboratory, Brussels), BRUCE BERNDT (University of Illinois at Urbana-Champaign), M. L. GLASSER (Clarkson College), OTTO G. RUEHR (Michigan Technological University), ROBERT E. SHAFER (Berkeley, CA), JAMES A. WILSON (Iowa State University) and the proposer.

In their discussions of part (c), Belevitch mentioned Perron [2]; Berndt referred to Bauer [3], Ramanujan [4], and Stieltjes [5]; Ruehr appealed to Wall [7]; and Glasser referred to Belevitch [6].

Wilson employed the theory of orthogonal polynomials to prove (c) and obtained the generalization

$$\frac{1}{z} \frac{\Gamma(3/4 + p + z)}{\Gamma(1/4 + p + z)} \frac{\Gamma(3/4 - p + z)}{\Gamma(1/4 - p + z)} = 1 + \frac{(1 - 16p^2)/32z^2}{1} + \frac{c_1/z^2}{1} + \frac{c_2/z^2}{1} + \dots$$

for $\operatorname{Re} z > 0$, with $c_n = [(2n + 1)^2 - 16p^2]/64$.

Shafer also referred to Ramanujan [4] and derived the following generalization:

$$\frac{\Gamma(z + \alpha + \beta + 3/4)\Gamma(z + \alpha - \beta + 3/4)\Gamma(z - \alpha + \beta + 3/4)\Gamma(z - \alpha - \beta + 3/4)}{\Gamma(z + \alpha + \beta + 1/4)\Gamma(z + \alpha - \beta + 1/4)\Gamma(z - \alpha + \beta + 1/4)\Gamma(z - \alpha - \beta + 1/4)} \\ = z^2 + 1/16 - \alpha^2 - \beta^2 - \frac{2(1/16 - \alpha^2)(1/16 - \beta^2)}{z^2 + 1/16 + 1^2 - \alpha^2 - \beta^2 - \frac{(9/16 - \alpha^2)(9/16 - \beta^2)}{z^2 + 1/16 + 2^2/2 - \alpha^2 - \beta^2 - \dots}}$$

REFERENCES

- [1] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER AND F. G. TRICOMI, *Higher Transcendental Functions*, Vol. I, McGraw-Hill, New York, 1953.
- [2] O. PERRON, *Die Lehre von den Kettenbrüchen*, Band II, Teubner, Stuttgart, 1957.
- [3] F. BAUER, *Von einem Kettenbruche Euler's und einem Theorem von Wallis*, Abh. Bayer. Akad. Wiss., 11 (1872), pp. 96–116.
- [4] S. RAMANUJAN, *Collected Papers*, Chelsea, New York, 1962, p. xxvii.
- [5] T. J. STIELTJES, *Note sur quelques fractions continues*, Quart. J. Math., 25 (1891), pp. 198–200.
- [6] V. BELEVITCH, *The Gauss hypergeometric ratio as a positive real function*, SIAM J. Math. Anal., 13 (1982), pp. 1024–1040.
- [7] H. S. WALL, *Analytic Theory of Continued Fractions*, Van Nostrand, New York, 1948, p. 371, l. 94.9.

An Iteration Problem

Problem 83-9, by P. C. LIU (Great Lakes Environmental Research Laboratory, Ann Arbor, MI).

Given the integral

$$I = \int_0^\infty F^{-b} \exp\{x(1 - F^{-b/x})\} dF,$$

where $x > 0$, $0 < I \leq 1$, $2 < b < 30$, determine an efficient iteration scheme to solve for x given I and b .

Solution by L. W. FULLERTON (IMSL, Inc., Houston).

The integral may be reduced to simpler form by letting $y = x F^{-b/x}$. We find

$$(1) \quad I(x, b) = b^{-1} e^x x^{1 - ((b-1)/b)x} \Gamma\left(\frac{b-1}{b} x\right).$$

Obtaining an efficient iteration scheme depends on finding good initial approxima-