Shear effects in nonhomogeneous turbulence

Citation for published version (APA):

Toschi, F., Lvque, E., & Ruiz-Chavarria, G. (2000). Shear effects in nonhomogeneous turbulence. *Physical Review Letters*, 85(7), 1436-1439. <https://doi.org/10.1103/PhysRevLett.85.1436>

DOI:

[10.1103/PhysRevLett.85.1436](https://doi.org/10.1103/PhysRevLett.85.1436)

Document status and date:

Published: 01/01/2000

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Shear Effects in Nonhomogeneous Turbulence

F. Toschi,¹ E. Lvque,² and G. Ruiz-Chavarria³

¹*Department of Applied Physics, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands*
INFM, Unit di Tor Vergata, Roma, Italy

²*Laboratoire de Physique CNRS, ENS de Lyon, 69364 Lyon Cedex 07, France*

³*Departamento de Fsica, Facultad de Ciencias, UNAM, 04510 Mexico D.F., Mexico*
 (Received 18 January 2000)

Motivated by recent experimental and numerical results, a simple unifying picture of intermittency in turbulent shear flows is suggested. Integral structure functions (ISF), taking into account explicitly the shear intensity, are introduced on phenomenological grounds. ISF can exhibit a universal scaling behavior, independent of the shear intensity. This picture is in satisfactory agreement with both experimental and numerical data. Possible extension to convective turbulence and implication on closure conditions for large-eddy simulation of nonhomogeneous flows are briefly discussed.

PACS numbers: 47.27.-i, 47.27.Ak, 47.27.Nz

Statistical properties of turbulent flows are usually characterized in terms of the scaling behavior of velocity structure functions (SF). These quantities are defined as the statistical moments of longitudinal velocity increments across a separation $\mathbf{r}$ at the location $\mathbf{x}$: $D_p(\mathbf{x}, \mathbf{r}) = \langle \delta v(\mathbf{x}, \mathbf{r})^p \rangle$. In homogeneous and isotropic turbulence, $D_p(\mathbf{x}, \mathbf{r})$ depends only on the distance (or scale) r . Experimental and numerical observations support the idea that $D_p(r)$ display universal power-law dependence on r in the so-called inertial range, i.e., $D_p(r) \sim r^{\zeta_p}$. Universality refers here to the scaling exponents ζ_p being independent of the stirring process of turbulence. The ζ_p values are found to be in disagreement with Kolmogorov's linear prediction (K41) $\zeta_p = p/3$ [1]. The understanding of this correction to K41, usually referred to as intermittency, has stimulated many phenomenological and theoretical works during the last 40 years (see [2] for a recent review).

Only recently, interests in understanding intermittency in nonhomogeneous turbulent flows have started to emerge (see [3–10]). The major point is to understand how the phenomenology of intermittency is modified, or can be extended, in case of nonhomogeneous flows. A common characteristic of such flows, e.g., wall-bounded flows, is the presence of a nonzero mean velocity gradient (usually called shear). Note that shear does not necessarily imply inhomogeneity, e.g., in homogeneous shear, straining, and rotational flows.

Our investigation starts from the following key observations: (i) In the presence of a strong shear, intermittency, defined as the deviation of scaling exponents ζ_p from the linear law, is larger than in homogeneous and isotropic turbulence. (ii) Relative scaling exponents, measured in very different flows but in positions where the shear is strong enough, seem to be very similar (universal).

Data in Fig. 1 fully confirm our two key observations. They come from very different situations: near the wall in a channel flow numerical simulation [4,5] and experiment [6], in the logarithmic sublayer of a boundary layer flow

[7], near a strong vortex [8], in the wake of a cylinder [9], and in a Kolmogorov flow [10].

In order to provide a theoretical unifying framework for nonhomogeneous turbulent shear flows, we start from the Navier-Stokes equations. The velocity field can be decomposed into a mean value (average is meant on time) plus a fluctuating part: $\mathbf{v}(\mathbf{x}; t) = \bar{\mathbf{v}}(\mathbf{x}) + \mathbf{v}'(\mathbf{x}; t)$. It yields the usual Reynolds decomposition

$$D_t v'_i + S_{ij}(\mathbf{x}) v'_j + v'_j \partial_j v'_i - \overline{v'_j \partial_j v'_i} = -\partial_i p' + \nu \Delta v'_i \quad (1)$$

with $D_t = (\partial_t + \bar{v}_j \partial_j)$. The shear is defined as $S_{ij}(\mathbf{x}) = \partial_j \bar{v}_i(\mathbf{x})$ and will depend on the mean flow geometry. In regions where $S_{ij} = 0$, e.g., very far from the boundaries, turbulence can be considered as homogeneous. Otherwise, the shear term $S_{ij}(\mathbf{x}) v'_j$ must be taken into account. The

FIG. 1. Intermittency corrections to scaling exponents, $\zeta_p/\zeta_3 - p/3$, in homogeneous and isotropic turbulence ($\square$), and several turbulent shear flows ($\triangle$). Intermittency corrections are significantly larger in the presence of shear and display universality.

results displayed in Fig. 1 show that the presence of this term modifies significantly the statistical properties of turbulence. In order to better highlight the physical implication of the shear, we consider together the second and third terms of the left-hand side of (1), defining the following integral structure functions (ISF):

$$\tilde{D}_p(z, r) = \langle [\delta v(z, r)^3 + \alpha r \cdot S(z) \cdot \delta v(z, r)^2]^{p/3} \rangle, \quad (2)$$

where α is an empirical prefactor of order one. These ISF are expected to take into account shear effects, particularly at large scales (see next paragraph) and to display a universal behavior. We consider here *generic* situations in which the shear reduces to $S(z) = \partial_z \bar{v}_x$. Such situations occur near a rigid wall, where the principal mean-velocity component aligns in the x direction, parallel to the boundary [11]. The shear $S(z)$ characterizes the variation of v_x along the z direction, i.e., as one moves off the wall. Finally, we consider increments in the direction of the mean flow, i.e., orthogonal to the shear direction.

ISF reduce on two different SF when either the first or the second term dominates. These two terms will exactly balance at scale L_S such that $\delta v(L_S)/L_S = \alpha S$. At scales $r \ll L_S$, shear effects become negligible and homogeneous and isotropic scalings are expected. Kolmogorov's scaling then yields $\delta v(r) \sim \epsilon^{1/3} r^{1/3}$, where ϵ denotes the mean energy dissipation rate. By extending this similarity relation to the scale $r = L_S$, one obtains the usual dimensional estimate for the shear length scale $L_S \sim (\epsilon/S^3)^{1/2}$ [12]. In the logarithmic layer of a plane near-wall flow it yields $L_S(z) \sim z$ [7]. Roughly speaking, L_S can be viewed as the size of small-scale eddies, whose turnover time equals the shear time scale $1/S$, imposed by the flow geometry and the stirring process of turbulence at large scales. Note that this estimate of the shear length scale stems out from dimensional analysis. In practice, there may be a prefactor in the expression of L_S : this is taken into account by the coefficient α . From previous reasoning it follows that $\tilde{D}_p(r) \sim D_p(r)$ for $r \ll L_S$ and $\tilde{D}_p(r) \sim (rS)^{p/3} D_{2p/3}(r)$ for $r \gg L_S$.

Another way to see that the central objects, in the presence of shear, are the ISF comes from the generalization of Yaglom's equation to homogeneous-shear flows, i.e., with $S(z) = S$ [12,13]:

$$-\frac{4}{5} \epsilon r = D_3(r) - 6\nu \frac{d D_2(r)}{dr} + \frac{2S}{r^4} \int_0^r dx x^4 \overline{v'_x(x_0) v'_z(x_0 + x)}. \quad (3)$$

Supposing that this relation can be generalized, along the same line of idea of the Kolmogorov's refined similarity hypothesis [14], one obtains

$$\delta v^3(r) + \alpha r \cdot S \cdot \overline{v'_x(x) v'_z(x+r)} \sim \epsilon(r) r, \quad (4)$$

where $\epsilon(r)$ denotes the coarse graining of the energy dissipation field $\epsilon = \frac{\nu}{2} \sum_{i,j} (\partial_i v_j)^2$, at scale r .

Simplifying, on pure dimensional grounds, the velocity cross correlation $\overline{v'_x v'_z}$, with $\delta v(r)^2$ one ends up again with $\tilde{D}_p(r)$. Furthermore, we propose the following refined similarity hypothesis:

$$\tilde{D}_p(r) = \langle [\delta v(r)^3 + \alpha r \cdot S \cdot \delta v(r)^2]^{p/3} \rangle \sim \langle \epsilon(r)^{p/3} \rangle r^{p/3}.$$

This formulation is consistent, in the limiting cases of strong and negligible shear, with some recent findings (see [4,5]). In addition to that, the ISF should be able to abridge smoothly between these two limiting regimes, i.e.,

$$r \ll L_S: D_p(r) \sim \langle \epsilon(r)^{p/3} \rangle r^{p/3}, \quad (5)$$

and

$$r \gg L_S: D_p(r) \sim \langle \epsilon(r)^{p/2} \rangle. \quad (6)$$

Equation (5) is in agreement with the restoration of homogeneity and isotropy at small scales. For $r \gg L_S$, Eq. (6) gives $D_2(r) \sim \text{const}$ (modulo possible logarithmic corrections), yielding for the energy spectrum $E(k) \sim k^{-1}$ (a relation suggested a long time ago in [12]).

In the previous picture, it is assumed that the shear length scale L_S remains larger than the dissipation length scale η . The dissipation field $\epsilon(r)$ is then expected to display the same scaling properties as in homogeneous and isotropic turbulence. However, in regions where $L_S \lesssim \eta$ (very close to the wall) it is expected that shear effects act down to the dissipative scale and therefore can modify the scaling behavior of $\epsilon(r)$. The scalings of $D_p(r)$ should then also change according to [6].

The relevance of integral structure functions for describing the scaling properties of nonhomogeneous shear flows is now tested on both experimental and numerical data. The experiment, performed in the recirculating wind tunnel of ENS de Lyon, consists of a turbulent boundary-layer flow over a smooth horizontal plate (see [7] for details about the experimental apparatus). Velocity measurements are carried out at various elevations from the plate in the logarithmic turbulent sublayer [11]. Numerical results are obtained from a direct simulation of the Navier-Stokes equations in a rectangular channel flow (see [4] for details).

In Fig. 2, D_3 and D_6 , measured in the logarithmic boundary sublayer, are compared with the corresponding $\tilde{D}_3$ and $\tilde{D}_6$. The shear $S(z) = \partial_z \bar{v}_x$ has been estimated from the mean-velocity profile. In standard nondimensional variables [15], namely, $z^+ = v_* z / \nu$ and $v^+ = \bar{v}_x / v_*$, where v_* is the characteristic velocity of the viscous sublayer, our data are well fitted by the logarithmic profile $v^+(z^+) = (1/\kappa) \log(z^+) + B$. We obtain $\kappa \approx 0.4$ and $B \approx 5.26$ in agreement with previously reported results [7]. For what concerns the coefficient α , all our results have been obtained with the fixed value

FIG. 2. In the turbulent boundary layer at the distance $z^+ = 102$, D_3 and D_6 (Δ) are compared with $\tilde{D}_3$ and $\tilde{D}_6$ ($\star$). The scale r has been renormalized by the characteristic shear length scale L_s . The solid lines passing through $\tilde{D}_3$ and $\tilde{D}_6$ indicate the expected *homogeneous* and *isotropic* power laws, respectively, $\zeta_3 = 1$ and $\zeta_6 = 1.78$. For comparison, the dashed line has slope 2.

$\alpha \approx 0.2$. Note that the constant α is not (*a priori*) intended to be universal but may strongly depend on the geometry and stirring process of the flow. However, it is expected to remain of order unity. In practice, the value of α has been extracted from data by requiring that one third-order ISF scale as r (see next paragraph). Corresponding estimates of L_s are indicated in all figures. Finally, one must point out that velocity increments have been estimated in the direction of the mean flow by use of the Taylor hypothesis. Both D_3 and D_6 exhibit a power-law dependence on r , but the scaling exponents are clearly different from those observed in homogeneous and isotropic (h-i) turbulence, respectively, $\zeta_3 \approx 1$ and $\zeta_6 \approx 1.78$. On the other hand, the corresponding ISF exhibit power laws in good agreement with h-i scalings (up to very large scales).

In Fig. 3, third-order SF and ISF are displayed for various distances from the wall. We notice that scaling behavior of D_3 changes with z^+ . On the contrary, all the corresponding $\tilde{D}_3$ display the same power-law scaling with exponent 1. We recall that the coefficient α is kept constant and the shear is estimated from the mean-velocity profile; there is no adjustable parameter.

We now report a sharper test: SF and ISF, compensated by h-i power-law scalings, are displayed in Fig. 4 for $p = 1, \dots, 6$ at distance $z^+ = 102$. A departure from h-i scalings is clearly observed for SF.

On the contrary, $\tilde{D}_p(r)/r^{\zeta_p}$ exhibits a plateau up to very large scales, indicating that ISF roughly behave as r^{ζ_p} . In other terms, ISF compensate shear effects and restore, via the extra term $\alpha r \cdot S \cdot \delta v(r)^2$, the h-i scalings. Finally, the same test, made on numerical data, is reported in Fig. 5 for the sake of comparison. Data are obtained

FIG. 3. From the boundary layer experiment, third-order structure function at various distances from the wall: $z^+ = 37$ ($\circ$), $z^+ = 124$ (Δ), and $z^+ = 233$ ($\square$). The scaling properties of D_3 (SF) do depend on the distance z^+ . On the contrary, $\tilde{D}_3$ (ISF) displays the same scaling behavior for all z^+ . The dashed line has slope 1. The curves have been shifted vertically for convenience.

at distance $z^+ = 25$ from the wall, i.e., where the shear is strong. Results are in reasonable agreement with those of Fig. 4, despite the lower resolution of the numerical simulation. We emphasize the major points of our study. Our description relates the scaling properties of velocity fluctuations in nonhomogeneous shear flows to those of the coarse-grained dissipation rate $\varepsilon(r)$. Provided that the shear length scale remains much larger than the dissipative length scale, we are inclined to believe that the scaling properties of $\varepsilon(r)$ remain flow independent; the dissipation process, operating on very small scales, is mainly insensitive to the presence of the shear. We then claim that the

FIG. 4. In the experimental boundary layer at $z^+ = 102$, SF and ISF, compensated by homogeneous and isotropic scalings, are displayed for $p = 1, \dots, 6$.

FIG. 5. As in Fig. 4 but from numerical data at $z^+ = 25$.

scalings of velocity fluctuations are different in sheared regions only because the similarity between $\delta v(r)$ and $\varepsilon(r)$ changes form [see (5) and (6)]. When $L_S \lesssim \eta$, shear effects acting down to dissipative scales are expected to modify the scaling behavior of $\varepsilon(r)$. Nonetheless, we observe in Fig. 1 that relative scalings exponents of velocity structure functions remain universal.

The introduction of ISF relies on quite simple dimensional arguments. However, they have proved to be valuable (preliminary) tools in order to capture (at leading order) shear effects on the scaling behavior of velocity structure functions. We believe that some refinements in the definition of ISF may be possible, e.g., considering the cross correlation $\overline{v_x^i v_z^j}$ instead of $\delta v(r)^2$: however, ISF are of practical interest as they are easily accessible experimentally.

A possible extension of ISF applies to thermal turbulent convection, the buoyancy term $ag\delta T(r)$ playing a similar role of $S \cdot \delta v(r)$. In that case, the ISF would read

$$\tilde{D}_p^{\text{RB}}(r) = \langle [\delta v(r)^3 + \alpha r ag\delta T(r)\delta v(r)]^p \rangle$$

and are expected to take into account buoyancy effects at scales larger than the Bolgiano length scale.

An important application of our findings concerns large eddy simulations (LES). Usual eddy viscosity models are known to fail close to the boundary. As one moves near the boundary, the shear length scale L_S becomes smaller and smaller. While L_S remains larger than the cutoff scale Δ of the LES, usual closure conditions, based on homogeneous and isotropic turbulent dynamics, remain acceptable. However, when L_S becomes comparable or smaller than Δ , shear effects must be taken into account and the closure condition should be modified. An alternative consists of

decreasing the mesh size near the wall so that L_S always remains larger than Δ . More simply, our study suggests considering $[\delta v(r)^3 + \alpha r S \delta v(r)^2]^{1/3}$ instead of $\delta v(r)$ in closure relations. Along this line of idea, the Smagorinski's closure condition [16] can be generalized in order to uniformly take into account shear effects. The eddy viscosity ν_{eddy} then reads

$$\nu_{\text{eddy}} = C_s \Delta^2 (\bar{S} + \alpha S),$$

where the extra term αS takes care of shear effects. C_s is the empirical constant of the Smagorinski's closure and $\bar{S}$ denotes the rate of strain on scale Δ .

F.T. thanks S. Ciliberto for his kind hospitality at ENS de Lyon. G.R.-C. and E.L. acknowledge the ECOS committee and CONACYT for their financial support under Project No. M96-E03. G.R.-C. also acknowledges DGAPA-UNAM for partial support under Project No. IN-107197.

-
- [1] A. N. Kolmogorov, C.R. Acad. Sci. USSR **30**, 301 (1941); **31**, 538 (1941); **32**, 16 (1941).
 - [2] U. Frisch, *Turbulence: The Legacy of A. N. Kolmogorov* (Cambridge University Press, Cambridge, England, 1995).
 - [3] L. Danaila, F. Anselmetti, T. Zhou, and R. A. Antonia, "A Generalization of Yaglom's Equation which Accounts for the Large-Scale Forcing in Heated Grid Turbulence" (to be published).
 - [4] F. Toschi, G. Amati, S. Succi, R. Benzi, and R. Piva, Phys. Rev. Lett. **82**, 5044 (1999).
 - [5] R. Benzi, G. Amati, C. M. Casciola, F. Toschi, and R. Piva, Phys. Fluids **11**, 1284 (1999).
 - [6] M. Onorato, R. Camussi, and G. Iuso, Phys. Rev. E **61**, 1447–1454 (2000).
 - [7] G. Ruiz-Chavarria, S. Ciliberto, C. Baudet, and E. Lvque, "Scaling Properties of the Streamwise Component of Velocity in a Turbulent Boundary Layer" (to be published).
 - [8] F. Chill and J.-F. Pinton, *Advances in Turbulence VII*, edited by U. Frisch (Kluwer Academic Publishers, Dordrecht, The Netherlands, 1998), pp. 211–214.
 - [9] E. Gaudin, B. Protas, S. Goujon-Durand, J. Wojciechowski, and J. E. Wesfreid, Phys. Rev. E **57**, R9 (1998).
 - [10] R. Benzi, M. V. Struglia, and R. Tripiccion, Phys. Rev. E **53**, 5565 (1996).
 - [11] H. Tennekes and J. L. Lumley, *A First Course in Turbulence* (MIT Press, Cambridge, MA, 1972).
 - [12] O. J. Hinze, *Turbulence* (McGraw-Hill, New York, 1959).
 - [13] M. V. Struglia, Ph.D. thesis, Universit di Roma, Tor Vergata, Roma, Italy, 1996.
 - [14] A. N. Kolmogorov, J. Fluid Mech. **62**, 82 (1962).
 - [15] H. Schlichting, *Boundary Layer Theory* (McGraw-Hill, New York, 1968).
 - [16] J. Smagorinski, Mon. Weather Rev. **91**, 99–164 (1963).