Solution to Problem 74-20*: Gravitational attraction

Citation for published version (APA):

Bouwkamp, C. J. (1976). Solution to Problem 74-20*: Gravitational attraction. *SIAM Review*, 18(1), 123-126.
<https://doi.org/10.1137/1018018>

DOI:

[10.1137/1018018](https://doi.org/10.1137/1018018)

Document status and date:

Published: 01/01/1976

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

*Problem 74-20**, *Gravitational Attraction*, by H. R. AGGARWAL (NASA, Ames Research Center).

Determine explicitly the mutual force of gravitational attraction between two congruent spherical segments forming a “dumb-bell” shaped body whose central cross section is given by the figure.

Solution by C. J. BOUWKAMP, Technological University, Eindhoven, the Netherlands.

Let units be so chosen that the force between two unit point masses with interdistance r equals $1/r^2$, and assume that the body has unit mass density. Further, set $p = a/R$ and let F be the mutual force of gravitational attraction between the two spherical segments. Then

$$(1) \quad F/(4\pi^2 R^4) = G(p) = \int_0^\infty (f(t, p))^2 t^{-1} dt,$$

in which

$$(2) \quad f(t, p) = \int_0^{1+p} e^{-tu} J_1(t(1 - (u - p)^2)^{1/2})(1 - (u - p)^2)^{1/2} du,$$

where J_1 denotes the Bessel function of order one.

To prove this, I first determine the mutual potential energy $V dz_1 dz_2$ between two thin coaxial disks. The first disk lies in the left segment, between z_1 and $z_1 + dz_1$, with distance z_1 to the midplane of the body; the second disk lies in the right segment, between z_2 and $z_2 + dz_2$, with distance z_2 to the same plane. Their axial interdistance is $z = z_1 + z_2$, while their radii are $R_i = (R^2 - (z_i - a)^2)^{1/2}$, $i = 1, 2$. In each disk, polar coordinates are introduced; one of the two angular integrations can be carried out because of rotational symmetry, and we are left with

$$(3) \quad V = 2\pi \int_0^{R_1} r_1 dr_1 \int_0^{R_2} r_2 dr_2 \int_0^{2\pi} d\theta (z^2 + r_1^2 + r_2^2 - 2r_1 r_2 \cos \theta)^{-1/2}.$$

This integral is easy to transform by means of Bessel-function techniques. With

$$(z^2 + r^2)^{-1/2} = \int_0^\infty e^{-zt} J_0(rt) dt,$$

the addition theorem of Bessel functions, and

$$\int_0^x t J_0(t) dt = x J_1(x),$$

it is found that

$$(4) \quad V = 4\pi^2 R_1 R_2 \int_0^\infty e^{-zt} J_1(R_1 t) J_1(R_2 t) t^{-2} dt.$$

The force between the two disks is $-(\partial V / \partial z) dz_1 dz_2$, in which the dependency of R_i on z_i is irrelevant. Since this force is in the axial direction, all forces can be added in scalar fashion. Thus

$$(5) \quad F = 4\pi^2 \int_0^\infty \frac{dt}{t} \left[\int_0^{R+a} R_1 J_1(R_1 t) e^{-tz_1} dz_1 \right] \left[\int_0^{R+a} R_2 J_1(R_2 t) e^{-tz_2} dz_2 \right].$$

Upon substituting $z_i = Ru$, the two inner integrals are seen to be identical and equal to $R^2 f(Rt, p)$, where f is defined in (2). The substitution $Rt \rightarrow t$ finally gives (1).

Let me first remark that for the “dumb-bell” shaped body the parameter p lies in the closed interval $0 \leq p \leq 1$, the endpoints corresponding to the cases of two touching half-spheres and two touching full spheres, respectively.

For two spheres in contact the force of attraction is known by elementary methods. This comes down to $G(1) = \frac{1}{9}$. I do not know whether the other limiting case was ever treated before, but I am able to prove that $G(0) = \frac{1}{12}$. Secondly, (1) and (2) are meaningful for $-1 \leq p < 0$, and then they describe the force between two segments each smaller than a half-sphere. If $G(p)$ denotes the force between the two spherical segments of the figure, then $G(-p)$ is precisely the force between their complements, the two “flat” segments that form the cut of the two full spheres. Of course, $G(-1) = 0$.

To further discuss (2), I substitute $u = p + \cos \theta$, so that

$$(6) \quad \begin{aligned} f(t, p) &= e^{-pt} \int_0^\alpha e^{-t \cos \theta} J_1(t \sin \theta) \sin^2 \theta d\theta, \\ \alpha &= \cos^{-1}(-p) = \pi - \cos^{-1}(p), \end{aligned}$$

where the principal value of the $\cos^{-1}$ function is implied. One has $\alpha = 0, \pi/2, \pi$ as $p = -1, 0, 1$, in that order. It is not too difficult to show that

$$(7) \quad \int_0^\pi e^{-t \cos \theta} J_1(t \sin \theta) \sin^2 \theta d\theta = \frac{2}{3} t,$$

but this is left to the reader. Thus $f(t, 1) = \frac{2}{3} t \exp(-t)$, and then $G(1) = \frac{1}{9}$ follows from (1).

The case $p = 0$ is not as nice, since

$$(8) \quad \begin{aligned} f(t, 0) &= \int_0^{\pi/2} e^{-t \cos \theta} J_1(t \sin \theta) \sin^2 \theta d\theta \\ &= t \left\{ \frac{1}{3} - \int_0^t x^{-2} J_2(x) dx \right\} = t \int_t^\infty x^{-2} J_2(x) dx, \end{aligned}$$

which may be left unproved here. By using the last expression of (8) and manipulating with integrals in (1), I obtain $G(0) = \frac{1}{12}$ as announced.

The two special cases above are included in the general equation

$$(9) \quad f(t, p) = t e^{-pt} \left\{ \frac{1}{3} + \frac{1}{2}p - \frac{1}{6}p^3 - (1 - p^2) \int_0^t e^{px} J_2(x\sqrt{1-p^2}) x^{-2} dx \right\},$$

first proved by my coworker D. L. A. Tjaden of the Philips Research Laboratories. If p is nonpositive, an alternative equation is

$$(10) \quad f(t, p) = t e^{-pt} (1 - p^2) \int_t^\infty e^{px} J_2(x\sqrt{1-p^2}) x^{-2} dx.$$

Several trials to use (9) or (10) in (1) have only led to unwieldy expressions, integrals of elliptic integrals, as might have been expected from (4), which is known to be expressible in terms of elliptic integrals. It does not seem possible to express F in terms of complete elliptic integrals, as I originally hoped in view of various similar problems encountered before.

A last resort to an "explicit" solution is numerical integration. With the exception of $p = 1$, the integral (1) is badly convergent; the integrand is oscillating

TABLE 1

p	$G(p)$	$G(-p)$	p	$G(p)$	$G(-p)$
0.00	0.0833333	0.0833333	0.50	0.1453554	0.0178390
0.02	0.0864331	0.0802335	0.52	0.1464798	0.0161248
0.04	0.0895273	0.0771391	0.54	0.1474414	0.0145013
0.06	0.0926106	0.0740553	0.56	0.1482337	0.0129693
0.08	0.0956770	0.0709873	0.58	0.1488503	0.0115294
0.10	0.0987209	0.0679402	0.60	0.1492849	0.0101817
0.12	0.1017364	0.0649187	0.62	0.1495313	0.0089263
0.14	0.1047175	0.0619278	0.64	0.1495831	0.0077628
0.16	0.1076582	0.0589720	0.66	0.1494345	0.0066906
0.18	0.1105525	0.0560558	0.68	0.1490793	0.0057088
0.20	0.1133941	0.0531836	0.70	0.1485116	0.0048161
0.22	0.1161769	0.0503596	0.72	0.1477259	0.0040109
0.24	0.1188944	0.0475878	0.74	0.1467164	0.0032910
0.26	0.1215405	0.0448722	0.76	0.1454778	0.0026543
0.28	0.1241086	0.0422165	0.78	0.1440047	0.0020980
0.30	0.1265924	0.0396242	0.80	0.1422923	0.0016188
0.32	0.1289854	0.0370987	0.82	0.1403356	0.0012131
0.34	0.1312811	0.0346432	0.84	0.1381302	0.0008769
0.36	0.1334728	0.0322607	0.86	0.1356718	0.0006055
0.38	0.1355542	0.0299540	0.88	0.1329565	0.0003937
0.40	0.1375185	0.0277259	0.90	0.1299809	0.0002358
0.42	0.1393593	0.0255786	0.92	0.1267417	0.0001253
0.44	0.1410700	0.0235144	0.94	0.1232366	0.0000551
0.46	0.1426439	0.0215352	0.96	0.1194636	0.0000172
0.48	0.1440746	0.0196430	0.98	0.1154217	0.0000023
0.50	0.1453554	0.0178390	1.00	0.1111111	0.0000000