Solution to Problem 85-16* : A conjectured definite integral

Citation for published version (APA):

Bouwkamp, C. J. (1986). Solution to Problem 85-16* : A conjectured definite integral. *SIAM Review*, 28(4), 568-569. <https://doi.org/10.1137/1028165>

DOI:

[10.1137/1028165](https://doi.org/10.1137/1028165)

Document status and date:

Published: 01/01/1986

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

An OC Curve Inequality

Problem 86-20*, by P. A. ROEDIGER and J. G. MARDO (U.S. Army Armament, Munitions and Chemical Command, Dover, NJ).

Let

$$\text{OC}(n, c, q) = \sum_{i=0}^c \binom{n}{i} q^{n-i} (1-q)^i$$

where $n > c > 0$ and $0 < q < 1$. Prove or disprove that

$$\text{OC}(n, c, q) < [\text{OC}(n, c, q^{1/m})]^m \quad \text{for all } m > 1.$$

In the terminology of lot-by-lot sampling inspection by attributes, e.g., per MIL-STD-105D, the Operating Characteristic (OC) curve defines the probability of accepting a lot whose true fraction effective is q , when the criterion is to accept if and only if $(n-c)$ or more effectives are found in a random n -sample. When $m > 1$ quality characteristics are distinguished, having effect rates q_i , lot quality is described by the profile $\tilde{Q} = (q_1, q_2, \dots, q_m)$ and, generally, the accept/reject criteria are such that probability of acceptance has the form

$$\text{PA}(\tilde{Q}) = \prod_{i=1}^m \text{OC}(n, c, q_i).$$

Since total lot quality q is the product of the q_i 's, one is naturally interested in $\text{PA}(\tilde{Q} | q)$, for a given q . The two sides of the proposed inequality can be shown to be optimal PA values, under this constraint. The difficulty is deciding which is the max and which is the min.

SOLUTIONS

A Conjectured Definite Integral

Problem 85-16*, by A. H. NUTTALL (Naval Underwater Systems Center, New London, CT).

It is conjectured that

$$\int_0^\pi \left\{ \frac{\sin x}{x} \exp(x \cot x) \right\}^\nu dx = \frac{\pi \nu^\nu}{\Gamma(1+\nu)} \quad \text{for } \nu \geq 0.$$

Prove or disprove.

The integral arose in a study of cross correlators. The conjectured result was discovered numerically first from the result for $\nu = \frac{1}{2}$ for which the computer output for the integral was recognized as $\sqrt{2\pi}$. The above result has been confirmed numerically to 15 decimal places for numerous values of ν in the range $[0, 150]$.

Solution by C. J. BOUWKAMP (Technische Hogeschool Eindhoven, Eindhoven, the Netherlands).

Nuttall's conjecture is true. Further, the integral is a special case of a general class of integral representations. Let $r(\beta) > 0$ be continuously differentiable on $[0, \pi)$ with $r(\beta) \rightarrow \infty$ as $\beta \rightarrow \pi$; define

$$f(\nu, \beta) := \cos\{\nu(r \sin \beta - \beta)\} + (r'/r) \sin\{\nu(r \sin \beta - \beta)\}.$$

Then

$$(1) \quad \int_0^\pi r^{-\nu} \exp(\nu r \cos \beta) f(\nu, \beta) d\beta = \frac{\pi \nu^\nu}{\Gamma(\nu+1)}.$$

The special (and most simple) choice for r is $r(\beta) = \beta / \sin \beta$, which makes $f=1$. Then replacing β by x gives the required formula.

The proof of (1) goes via a Hankel-type integral,

$$\frac{i}{2} \int_\infty^{(0+)} (-s)^{-\nu-1} e^{-\nu s} ds = \frac{\pi \nu^\nu}{\Gamma(\nu+1)},$$

the integration-path being parametrized through polar coordinates:

$$s = -r(\beta) \exp(i\beta), \quad ds = -ir(1 - ir'/r) \exp(i\beta) d\beta,$$

and assuming that the path is symmetric with respect to the real axis.

Also solved by D. J. BORDELON (Naval Underwater Systems Center, New London, CT), N. G. DE BRUIJN (Eindhoven University of Technology, the Netherlands), C. COSGROVE AND M. L. GLASSER (Clarkson University), W. E. HORNOR AND C. C. ROUSSEAU (Memphis State University), A. A. JAGERS (Technische Hogeschool Twente, Enschede, the Netherlands), D. S. JONES (University of Dundee, Scotland, U. K.), W. A. J. LUXEMBURG (California Institute of Technology), C. L. MALLOWS (AT&T Bell Laboratories, Murray Hill, NJ), O. G. RUEHR (Michigan Technological University), N. M. TEMME (Centrum voor Wiskunde en Informatica, Amsterdam, the Netherlands) and P. WAGNER (University of Innsbruck, Austria).

Most solvers employed contour integration in some fashion using a Hankel-type integral as above. Jones wrote

$$\lim_{x \rightarrow 0} \pi \left(\frac{2}{x} \right)^\nu J_\nu(\nu x) = \frac{\pi \nu^\nu}{\nu!}$$

where

$$J_\nu(\nu x) = \frac{1}{\pi} \int_0^\pi \exp\{-\nu F(\theta, x)\} d\theta,$$

and

$$F(\theta, x) = -\cot \theta (\theta^2 - x^2 \sin^2 \theta)^{1/2} + \ln \{ \theta + (\theta^2 - x^2 \sin^2 \theta)^{1/2} \} / x \sin \theta,$$

Mallows specialized his previous result [1] which was proved independently [2]. Cosgrove and Glasser also used Mallows's result which they generalized and will describe elsewhere.

Hornor and Rousseau point out the connection between this problem and the analytic continuation of the series $\sum_{n=1}^\infty (n^{n-1}/n!) w^n$ which is a well-known Lagrange-Bürman expansion for the solution of $w = ze^{-z}$.

REFERENCES

- [1] C. L. MALLOWS, *Problem 6245*, posed December, 1978, solution in *Amer. Math. Monthly*, 83 (1980), p. 584.
- [2] R. EVANS, M. E. H. ISMAIL AND D. STANTON, *Coefficients in expansions of certain rational functions*, *Canad. J. Math.*, 34 (1982), pp. 1011-1024.