Response to "Comment on: `Thermodynamics of viscoelastic fluids: the temperature equation`"

Citation for published version (APA):

Wapperom, P., & Hulsen, M. A. (1998). Response to "Comment on: `Thermodynamics of viscoelastic fluids: the temperature equation`". *Journal of Rheology*, 42(6), 1569-1570. <https://doi.org/10.1122/1.550971>

DOI:

[10.1122/1.550971](https://doi.org/10.1122/1.550971)

Document status and date:

Published: 01/01/1998

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

**Response to “Comment on: ‘Thermodynamics
of viscoelastic fluids: The temperature equation’ ”
[J. Rheol. 42, 1565–1567 (1998)]**

[S0148-6055(98)01906-3]

The thermodynamics of irreversible processes is based on the principle of local action, i.e., the principle of local and instantaneous equilibrium. This assumption means that, although the system as a whole is not in equilibrium, the material can be divided into small physical subelements, where each subelement forms a small homogeneous equilibrium system. All basic assumptions, including this one, in the theory of internal variables are discussed in detail by Kuiken (1994, Chap. 3). We will only summarize his comments on the principle of local action in relation to the temperature here. Measurement of the temperature requires that an equilibrium situation between thermometer and fluid exists. In nonequilibrium situations, measurement of the temperature is only meaningful if the thermometer is at least surrounded by a small amount of fluid that is instantaneously in equilibrium with the thermometer. Here, small means small compared to dimensions over which the temperature varies measurably, and large compared to the thermometer. Instantaneously means that the time scale of the temperature change in the fluid is large compared to the time scale of the thermometer. The principle of local action ensures that equilibrium quantities like temperature, pressure, and entropy are well defined in nonequilibrium situations. On the other hand, this principle, of course, restricts the area of application of the thermodynamic theory somewhat. If this is a serious disadvantage needs to be established by experiments. The principle of local equilibrium is not only needed for the temperature equation, but is a main assumption in the thermodynamic theory. It is, for example, also used by Jongschaap *et al.* (1994) and Leonov (1987) to derive the (isothermal) viscoelastic constitutive equations from the thermodynamics. As we describe the influences of the viscoelasticity on the temperature equation, it seems plausible that our temperature equation is valid whenever the stress constitutive equations are. For this, the discussion on the temperature and its measurement is only meaningful for “real nonequilibrium” situations, i.e., when there is no local and instantaneous equilibrium, and not for the theory in our paper.

In the penultimate paragraph Professor Grmela states that we suggest another setting for nonisothermal rheology. This is not exactly the case: in our article we have only included one type of internal variable, the configuration tensor. This quantity describes the viscoelasticity, and all well-known differential models can be obtained with it. Our purpose was to demonstrate which effects the viscoelasticity (by means of the configuration tensor) has on the temperature equation, i.e., how does the temperature equation change when the configuration tensor is introduced, and to relate the obtained effects to existing experimental data. Our purpose was not to give some general setting for nonisothermal flow. For this, we have excluded possible scalar and vectorial relaxation processes in the section “Thermodynamics of viscoelastic fluids: General.” Including those ones would not be very relevant, because there are no experimental data for validation. Until experimental results become available, it thus remains an open question whether other relaxation processes play a role, and whether our temperature equation (and possi-

bly also the stress equation) thus has to be extended with other internal variables. Possible options are a vector internal variable for the heat flux in case of large temperature gradients and a scalar internal variable for volume relaxation in case of high-pressure gradients [see Kuiken (1994) for details]. However, as we do not intend to give a more general setting than the configuration tensor, comparing the settings of our paper and of Grmela is, therefore, not very meaningful.

The fact that we do not have relaxation of the heat flux due to the temperature gradient (the Cattaneo equation) is directly caused by not taking into account vectorial internal variables, describing the relaxation of the heat flux. For the details of the derivation see Kuiken (1994). Note that by relating the anisotropic heat conduction to the configuration tensor describing the polymer chains, we also have relaxation of the heat flux, but due to the dependence of the heat conduction tensor on the configuration tensor (in a similar manner as the elastic stress has relaxation). As far as we know, no experimental data are available that indicate that the Cattaneo equation is important for viscoelastic fluids, or that the effect is relevant compared to the relaxation caused by the anisotropic heat conduction in the polymer chains.

Summarizing, we think that Professor Grmela's Letter to the Editor is not very relevant in relation to our paper. The temperature seems a problem for his thermodynamics and not for ours. Furthermore, our settings were restricted by the fact that we only wanted to show the effect of viscoelasticity on the temperature equation, not to include possible internal variables for which no experimental evidence exists. Contrarily to Professor Grmela, we would turn things around: a temperature equation for viscoelastic fluids should minimally contain the nonisothermal effects described in our temperature equation. If necessary it can be extended.

Peter Wapperom

Centre for Systems Engineering and Applied Mechanics
Université catholique de Louvain
Bâtiment Euler, Avenue Georges Lemaître, 4
B-1348 Louvain-la-Neuve, Belgium

Martien A. Hulsen

Delft University of Technology
Faculty of Mechanical Engineering and Marine Technology
Laboratory for Aero and Hydrodynamics
Rotterdamseweg 145
2628 AL Delft, The Netherlands

References

- Jongschaap, R. J. J., K. H. de Haas, and C. A. J. Damen, "A generic matrix representation of configuration tensor rheological models," *J. Rheol.* **38**, 769–796 (1994).
Kuiken, G. D. C., *Thermodynamics of Irreversible Processes* (Wiley, Chichester, U.K., 1994).
Leonov, A. I., "On a class of constitutive equations for viscoelastic fluids," *J. Non-Newtonian Fluid Mech.* **25**, 1–59 (1987).