Classification of phase transitions of finite Bose-Einstein condensates in power law traps by Fisher zeros

Citation for published version (APA):

Mülken, O., Borrmann, P., Harting, J. D. R., & Stamerjohanns, H. (2001). Classification of phase transitions of finite Bose-Einstein condensates in power law traps by Fisher zeros. *Physical Review A : Atomic, Molecular and Optical Physics*, 64(1), 013611-1/6. [013611]. <https://doi.org/10.1103/PhysRevA.64.013611>

DOI:

[10.1103/PhysRevA.64.013611](https://doi.org/10.1103/PhysRevA.64.013611)

Document status and date:

Published: 01/01/2001

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Classification of phase transitions of finite Bose-Einstein condensates in power-law traps by Fisher zeros

Oliver Mülken, Peter Borrmann, Jens Harting, and Heinrich Stamerjohanns

Department of Physics, Carl von Ossietzky University Oldenburg, D-26111 Oldenburg, Germany

(Received 20 June 2000; revised manuscript received 27 November 2000; published 5 June 2001)

We present a detailed description of a classification scheme for phase transitions in finite systems based on the distribution of Fisher zeros of the canonical partition function in the complex temperature plane. We apply this scheme to finite Bose systems in power-law traps within a semi-analytic approach with a continuous one-particle density of states $\Omega(E) \sim E^{d-1}$ for different values of d and to a three-dimensional harmonically confined ideal Bose gas with discrete energy levels. Our results indicate that the order of the Bose-Einstein condensation phase transition sensitively depends on the confining potential.

DOI: 10.1103/PhysRevA.64.013611

PACS number(s): 03.75.Fi, 05.30.Jp, 05.20.-y, 64.60.-i

I. INTRODUCTION

In 1924, Bose and Einstein predicted that in a system of bosons at temperatures below a certain critical temperature T_C , the single-particle ground state is macroscopically occupied [1]. This effect is commonly referred to as Bose-Einstein condensation, and a large number of phenomena, such as the condensation phenomena in alkali-metal atoms, the superfluidity of ^4He , and the superconductivity, are identified as signatures of this effect. However, the physical situation is very intricate in most experiments.

Recent experiments with dilute gases of alkali-metal atoms in magnetic [2] and optical [3] traps are in some sense the best experimental approximation up to now of the ideal noninteracting Bose-Einstein system in an external power-law potential. The achievement of ultralow temperatures by laser cooling and evaporative cooling provides the opportunity to study Bose-Einstein condensation under systematic variation of adjustable external parameters, e.g., the trap geometry, the number of trapped atoms, the temperature, and by the choice of the alkali-metal atoms the effective interparticle interactions. Even in the approximation of noninteracting particles, an explanation of these experiments requires some care, because the number of bosons in these novel traps is finite and fixed and the standard grand-canonical treatment is not appropriate. The effect of the finite particle numbers on the second moments of the distribution function, e.g., the specific heat and the fluctuation of the ground-state occupation number, has been addressed in a number of publications [4,5]. In [4,6], we have presented a recursion method to calculate the canonical partition function for non-interacting bosons, and we investigated the dependency of the thermodynamic properties of the condensate on the trap geometry.

The order of the phase transition in small systems sensitively depends on finite-size effects. Compared to the macroscopic system, even for systems as simple as the three-dimensional ideal gas, the order of the phase transition might change for mesoscopic systems where the number of particles is finite or for trapped gases with different trap geometries.

In this paper, we address the classification of the phase transition of a finite number of noninteracting bosons in a

power-law trap with an effective one-particle density of states $\Omega(E) = E^{d-1}$ being formally equivalent to a d -dimensional harmonic oscillator or a $2d$ -dimensional ideal gas. We use a classification scheme based on the distribution of zeros of the canonical partition function initially developed by Grossman *et al.* [7] and Fisher *et al.* [8], which has been extended by us [9] as a classification scheme for finite systems. On the basis of this classification scheme, we are able to extract a qualitative difference between the order of the phase transition occurring in Bose-Einstein condensates in three-dimensional traps [10,11] and in two-dimensional traps that was recently discovered by Safonov *et al.* in a gas of hydrogen atoms absorbed on the surface of liquid helium [12]. Since we do not consider particle interactions, this difference is only due to the difference in the confining potential.

We give a detailed review of the classification scheme in Sec. II. In Sec. III, we present a method for the calculation of the canonical partition function in the complex plane and describe details of the numerical implementation. Our results for $d=1-6$ and particle numbers varying from 10 to 300 are presented in Sec. III as well as calculations for a three-dimensional parabolically confined Bose gas.

II. CLASSIFICATION SCHEME

In 1952, Yang and Lee showed that the grand-canonical partition function can be written as a function of its zeros in the complex fugacity plane, which, for systems with hard-core interactions and for the Ising model, lie on a unit circle [13].

Grossmann *et al.* [7] and Fisher [8] extended this approach to the canonical ensemble by analytic continuation of the inverse temperature to the complex plane $\beta \rightarrow \beta = \beta + i\tau$. Within this treatment, all phenomenologically known types of phase transitions in macroscopic systems can be identified from the properties of the distribution of zeros of the canonical partition function.

In [9], we presented a classification scheme for finite systems that has its macroscopic equivalent in the scheme given by Grossmann. As usual, the canonical partition function reads

$$Z(\mathcal{B}) = \int dE \Omega(E) \exp(-\mathcal{B}E), \quad (2.1)$$

which we write as a product $Z(\mathcal{B}) = Z_{\text{lim}}(\mathcal{B})Z_{\text{int}}(\mathcal{B})$, where $Z_{\text{lim}}(\mathcal{B})$ describes the limiting behavior of $Z(\mathcal{B})$ for $T \rightarrow \infty$, imposing that $\lim_{T \rightarrow \infty} Z_{\text{int}}(\mathcal{B}) = 1$. This limiting partition function will only depend on the external potential applied to the system, whereas $Z_{\text{int}}(\mathcal{B})$ will depend on the specific interaction between the system particles. For example, for an N -particle system in a d -dimensional harmonic trap, $Z_{\text{lim}}(\mathcal{B}) = \mathcal{B}^{-dN}$ and thus the zeros of $Z(\mathcal{B})$ are the same as the zeros of $Z_{\text{int}}(\mathcal{B})$. Since the partition function is an integral function, the zeros $\mathcal{B}_k = \mathcal{B}_{-k}^* = \beta_k + i\tau_k$ ($k \in \mathbb{N}$) are complex conjugate and the partition function reads

$$Z(\mathcal{B}) = Z_{\text{lim}}(\mathcal{B})Z_{\text{int}}(0)\exp(\mathcal{B}\partial_{\mathcal{B}}\ln Z_{\text{int}}(0)) \times \prod_{k \in \mathbb{N}} \left(1 - \frac{\mathcal{B}}{\mathcal{B}_k}\right) \left(1 - \frac{\mathcal{B}}{\mathcal{B}_k^*}\right) \exp\left(\frac{\mathcal{B}}{\mathcal{B}_k} + \frac{\mathcal{B}}{\mathcal{B}_k^*}\right). \quad (2.2)$$

The zeros of $Z(\mathcal{B})$ are the poles of the Helmholtz free energy $F(\mathcal{B}) = -(1/\mathcal{B})\ln Z(\mathcal{B})$, i.e. The free energy is analytic everywhere in the complex temperature plane except at the zeros of $Z(\mathcal{B})$.

Different phases are represented by regions of holomorphy that are separated by zeros lying dense on lines in the complex temperature plane. In finite systems, the zeros do not squeeze on lines, which leads to a more blurred separation of different phases. We interpret the zeros as boundary posts between two phases. The distribution of zeros contains the complete thermodynamic information about the system, and all thermodynamic properties are derivable from it. Within this picture, the interaction part of the specific heat is given by

$$C_{V,\text{int}}(\mathcal{B}) = -k_B \mathcal{B}^2 \sum_{k \in \mathbb{N}} \left[\frac{1}{(\mathcal{B}_k - \mathcal{B})^2} + \frac{1}{(\mathcal{B}_k^* - \mathcal{B})^2} \right]. \quad (2.3)$$

The zeros of the partition function are poles of $C_V(\mathcal{B})$. As can be seen from Eq. (2.3), a zero approaching the real axis infinitely close causes a divergence at real temperature. The contribution of a zero $\mathcal{B}_k$ to the specific heat decreases with increasing imaginary part τ_k . Thus, the thermodynamic properties of a system are governed by the zeros of Z close to the real axis.

The basic idea of the classification scheme for phase transitions in small systems presented in [9] is that the distribution of zeros close to the real axis can be described approximately by three parameters, where two of them reflect the order of the phase transition and the third merely the size of the system.

We assume that the zeros lie on straight lines (see Fig. 1) with a discrete density of zeros given by

$$\phi(\tau_k) = \frac{1}{2} \left(\frac{1}{|\mathcal{B}_k - \mathcal{B}_{k-1}|} + \frac{1}{|\mathcal{B}_{k+1} - \mathcal{B}_k|} \right), \quad (2.4)$$

FIG. 1. Schematic illustration of the zeros in the complex temperature plane.

with $k=2,3,4,\dots$, and we approximate for small τ the density of zeros, by a simple power law $\phi(\tau) \sim \tau^\alpha$. Considering only the first three zeros the exponent α can be estimated as

$$\alpha = \frac{\ln \phi(\tau_3) - \ln \phi(\tau_2)}{\ln \tau_3 - \ln \tau_2}. \quad (2.5)$$

The second parameter to describe the distribution of zeros is given by $\gamma = \tan \nu \sim (\beta_2 - \beta_1)/(\tau_2 - \tau_1)$, where ν is the crossing angle of the line of zeros with the real axis (see Fig. 1). The *discreteness* of the system is reflected in the imaginary part τ_1 of the zero closest to the real axis.

In the thermodynamic limit, we have always $\tau_1 \rightarrow 0$. In this case, the parameters α and γ coincide with those defined by Grossmann *et al* [7], who have shown how different types of phase transitions can be attributed to certain values of α and γ . They claimed that $\alpha=0$ and $\gamma=0$ correspond to a first-order phase transition, second-order transitions correspond to $0 < \alpha < 1$ with $\gamma=0$ or $\gamma \neq 0$, third-order transitions to $1 \leq \alpha < 2$ with arbitrary values of γ , and that all higher order phase transition correspond to $\alpha > 1$. For macroscopic systems (with $\tau_1 \rightarrow 0$), α cannot be smaller than zero, because this would cause a divergence of the internal energy. However, in small systems with a finite τ_1 this is possible.

In our classification scheme, we therefore define phase transitions in small systems to be of first order for $\alpha \leq 0$, while second- and higher-order transitions are defined in complete analogy to the Grossmann scheme augmented by the third parameter τ_1 . The definition of a critical temperature β_C in small systems is crucial and ambiguous since no thermodynamic properties diverge. Thus, different definitions are possible. We define the critical temperature as $\beta_{\text{cut}} = \beta_1 - \gamma\tau_1$, i.e., the crossing point of the approximated line of zeros with the real temperature axis. An alternative definition is the real part of the first complex zero β_1 . In the thermodynamic limit, both definitions coincide.

Comparing the specific heats calculated for different discrete distributions of zeros shows the advantages of this classification scheme. Figure 2 shows (a) three distributions of zeros lying on straight lines corresponding to a first-order transition ($\alpha=0$ and $\gamma=0$), a second-order transition (α

FIG. 2. Plot of (a) generated zeros lying on straight lines to simulate first- ($\alpha=0$ and $\gamma=0$), second-, ($\alpha=0.5$ and $\gamma=-0.5$), and third- ($\alpha=1.5$ and $\gamma=-1$) order phase transitions and (b) the appropriate specific heats per particle.

$=0.5$ and $\gamma=-0.5$), and a third-order phase transition ($\alpha=1.5$, and $\gamma=-1$) and (b) the pertinent specific heats. In all cases the specific heat exhibits a hump extending over a finite-temperature region and cannot be used to classify the phase transition. In contrast, even for very small systems (large τ_1), the order of the phase transition is extractable from the distribution of zeros.

The zeros of the canonical partition function have a distinct geometrical interpretation, which explains the smoothed curves of the specific heat and other thermodynamic properties in finite systems.

Figure 3 shows (a) the ground-state occupation number $|\eta_0(\mathcal{B})|/N$ in the complex temperature plane and (b) the ground-state occupation number at real temperatures for a finite ideal Bose gas of $N=120$ particles, where $\eta_0(\mathcal{B})$ is given by the derivative of the logarithm of the canonical partition function $Z(\mathcal{B})$ with respect to the ground-state energy ϵ_0 , i.e., $\eta_0(\mathcal{B}) = -(1/\mathcal{B}) \partial_{\epsilon_0} Z(\mathcal{B})/Z(\mathcal{B})$.

Zeros of the partition function are poles of $\eta_0(\mathcal{B})$ and are indicated by dark spots, which influence the value of the ground-state occupation number at real temperatures impres-

FIG. 3. Comparison of (a) $|\eta_0|/N$ with (b) the appropriate value of η_0 at real temperatures for a 120-particle harmonically trapped ideal Bose gas (note that $\hbar=k_B=\omega=1$).

sively. Every pole seems to *radiate* onto the real axis and therefore determines the occupation number at real temperatures. This *radiation* extends over a broad temperature range so that the occupation number for real temperatures does not show a discontinuity but rather a smoothed curve. A closer look at Eq. (2.3) gives the mathematical explanation for this effect. The discrete distribution of zeros, i.e., $\tau_1 > 0$, inhibits the specific heat and all other thermodynamic properties to show a divergency at some critical temperature because the denominators of the arguments of the sum remain finite.

Without going into a detailed analysis, we note that in the thermodynamic limit the parameter α is connected to the critical index for the specific heat by

$$C_V \sim (\beta - \beta_c)^{\alpha-1}. \quad (2.6)$$

However, since critical indices are used to describe the shape of a divergency at the critical point, an extension to small systems seems to be more or less academic.

The introduction of complex temperatures might seem artificial at first sight, but, in fact, the imaginary parts τ_k of the complex zeros $\mathcal{B}_k$ have an obvious quantum-mechanical interpretation. We write the quantum-mechanical partition function as

$$Z(\beta + i\tau/\hbar) = \text{Tr}(\exp(-i\tau\hat{H}/\hbar)\exp(-\beta\hat{H})) \quad (2.7)$$

$$= \langle \Psi_{\text{can}} | \exp(-i\tau\hat{H}/\hbar) | \Psi_{\text{can}} \rangle \quad (2.8)$$

$$= \langle \Psi_{\text{can}}(t=0) | \Psi_{\text{can}}(t=\tau) \rangle, \quad (2.9)$$

introducing a *canonical state* as a sum over Boltzmann-weighted eigenstates $|\Psi_{\text{can}}\rangle = \sum_k \exp(-\beta\epsilon_k/2) |\phi_k\rangle$. We explicitly write the imaginary part as $\tau/\hbar$ since the dimension is $1/[\text{energy}]$ and the imaginary part therefore can be interpreted as time. Then the imaginary parts τ_k of the zeros resemble those times for which the overlap of the initial canonical state with the time-evoluted state vanishes. However, they are not connected to a single system but to a whole ensemble of identical systems in a heat bath with an initial Boltzmann distribution.

III. BEC IN POWER-LAW TRAPS

In this section, we assume a continuous single-particle density of states $\Omega(E) = E^{d-1}$ as an approximation for a d -dimensional harmonic oscillator or a $2d$ -dimensional ideal gas. For example, for the harmonic oscillator this corresponds to the limit of $\hbar\omega \rightarrow 0$ and taking only the leading term of the degeneracy of the single-particle energy levels. The one-particle partition function is given by the Laplace transformation

$$Z_1(\mathcal{B}) = \int dE E^{d-1} \exp(-\mathcal{B}E) = (d-1)! \mathcal{B}^{-d}. \quad (3.1)$$

The canonical partition function for N noninteracting bosons can be calculated by the following recursion [6]:

FIG. 4. Distribution of zeros for Bose-Einstein condensates with continuous one-particle density of states $\Omega(E)=E^{d-1}$ for $d=1-6$.

$$Z_N(\mathcal{B}) = \frac{1}{N} \sum_{k=1}^N Z_1(k\mathcal{B}) Z_{N-k}(\mathcal{B}), \quad (3.2)$$

where $Z_1(k\mathcal{B}) = \sum_i \exp(-k\mathcal{B}\epsilon_i)$ is the one-particle partition function at temperature $k\mathcal{B}$ and $Z_0(\mathcal{B}) = 1$. For small particle numbers, this recursion works fine, even though its numerical effort grows proportional to N^2 .

With Eq. (3.1) as Z_1 , Eq. (3.2) leads to a polynomial of order N in $(1/\mathcal{B})^d$ for Z_N , which can be easily generated using MAPLE or MATHEMATICA. The zeros of this polynomial can be found by standard numerical methods.

Figure 4 displays the zeros of the N -particle partition function for $d=1-6$ in the complex temperature plane for particle numbers $N=25, 50$, and 100 . For $d=2-6$, the zeros approach the positive real axis with increasing particle number and are shifted to higher temperatures, which is already an indicator of phase transitions. For $d=1$, the zeros approach the real axis only at negative temperature. This behavior is consistent with the usual prediction that there is no Bose-Einstein condensation for the one-dimensional harmonic oscillator and the two-dimensional ideal Bose gas [10].

The symmetry of the distributions of zeros is due to the fact that Z_N is a polynomial in $\mathcal{B}^{-d}$. For this reason, it can be inferred that for $d \rightarrow \infty$ the zeros lie on a perfect circle.

FIG. 5. Specific heat scaled by dN of Bose-Einstein condensates with continuous one-particle density of states for $d=1-6$.

Figure 5 shows the corresponding specific heats calculated using Eq. (2.3). As expected, for $d=1$ the specific heat has no hump and approaches with increasing temperature the classical value. We therefore exclude the analysis of $d=1$ from the discussions below. For $d=2-6$, the specific heats show humps or peaks, which get sharper with increasing d and increasing particle number. However, from these smooth curves the orders of the phase transition cannot be deduced.

In Fig. 6, the classification parameters α, γ, τ_1 defined above are plotted for two to six dimensions and particle numbers up to $N=100$. For all values of d , the parameter α is only a slightly varying function of N and approaches very fast an almost constant value. Since α is the primary classification parameter, from Fig. 6(a) we can directly infer that the $d=2$ system exhibits a third-order phase transition ($\alpha > 1$) while the transition for all higher dimensions is of second order ($0 \leq \alpha \leq 1$). For $N=50$, the dependence of α on d is plotted in Fig. 7(a). Since α decreases rather rapidly with increasing d , it can be speculated that systems corresponding to a large d exhibit a phase transition that is almost of first order. As mentioned above, for finite systems even values $\alpha \leq 0$ cannot be excluded for mathematical reasons. We note that two-dimensional Bose gases are an interesting and growing field of research. As is well known, the ideal free Bose gas in two dimensions ($d=2$) does not show a phase

FIG. 6. Classification parameters α , γ , and τ_1 for $d=2-6$ versus particle numbers N .

transition due to thermal fluctuations that destabilize the condensate [14]. Switching on a confining potential greatly influences the properties of the gas, the thermal fluctuations are suppressed, and the gas will show Bose-Einstein condensation. Recent experiments [12] have shown that Bose-Einstein

FIG. 7. Classification parameters for $N=50$ for different densities of states $\Omega(E)=E^{d-1}$ and $d=2-10$.

FIG. 8. Comparison between calculated zeros of the canonical partition function for three-dimensional trap geometries with (a) a continuous single-particle density of states and (b) discrete energy levels for $N=40$.

condensation is possible even though it is called a quasicondensate. In our notion, the quasicondensate is just a third-order phase transition. Thus, our results are in complete agreement with recent experiments and earlier theoretical work. An interesting question in this respect is whether the order of the transition changes for $d=2$ in the limit $N \rightarrow \infty$. Additional calculations for larger N , which are not printed in Fig. 6, indicate that α approaches 1 or might even get smaller. Note that $d=2$ is equivalent to a hypothetical four-dimensional ideal Bose gas or bosons confined in a two-dimensional parabolic trap. Our results indicate that the order of the phase transition depends sensitively on d for values around 2. This might be the reason why phase transitions in three space dimensions are sometimes classified as second- and sometimes as third-order phase transitions.

The parameter τ_1 is a measure of the finite size of the system, i.e., the scaling behavior of τ_1 as a function of N is a measure of how fast a system approaches a true n th-order phase transition in the Ehrenfest sense. The N dependence of τ_1 is displayed in Fig. 6(c). The scaling behavior can be approximated by $\tau_1 \sim N^{-\delta}$ with δ ranging between 1.06 and 1.12 for $d=2-6$.

The d dependence of the classification parameter is visualized in Fig. 7 for 50 particles. For this system size, we found $\alpha \sim d^{-4/3}$ and $\tau_1 \sim d^{-4/3}$.

The results presented above for continuous single-particle densities of states $\Omega(E)=E^{d-1}$ are obtained within semianalytical calculations. In order to compare these results to systems with a discrete level density, we adopt as a reference system the three-dimensional harmonic oscillator with the partition function given by

$$Z(B) = \sum_{n=0}^{\infty} \frac{(n+2)(n+1)}{2} \exp(-B(n+3/2)), \quad (3.3)$$

with $\hbar = \omega = k_B = 1$.

Figure 8(a) displays the zeros of the partition function (3.1) for $d=2$ and $d=3$. Figure 8(b) displays a contour plot of the absolute value of the ground-state occupation number $\eta_0(B) = -(1/B) \partial_{\epsilon_0} Z(B)/Z(B)$ with Z given by Eq. (3.3) calculated using an alternative recursion formula [4]. The zeros of Z are poles of η_0 and are indicated by dark spots in this figure.

Analyzing the distribution of zeros consolidates our speculation that the order of the phase transition depends sensitively on d . The distribution of zeros behaves like the above calculated values for $d=2$ but quantitatively like $d=3$. Since the degeneracy of the three-dimensional harmonically confined ideal Bose gas is a second-order polynomial, the quadratic term is not the only term that must be taken into account. The linear term becomes dominant for lower temperatures, so for very low temperatures the best approximation of a continuous one-particle density of states is $\Omega(E)=E$. The parameter α supports this statement [9], i.e., α resides in a region above 1, whereas the parameter γ behaves like the $d=3$ case. Finally, the parameter τ_1 , which is a measure for the discreteness of the system, shows a $\tau_1 \sim N^{-0.96}$ dependence that is comparable to the one for $d=2$. Thus, for small systems the phase transition is of third order; it can be speculated whether it becomes a second-order transition in the thermodynamic limit.

Our calculations are in very good agreement with recent theoretical works, not only qualitatively but also quantitatively [15,16]. Comparing the *critical* temperature, which we defined in Sec. II, with the usually utilized temperature of the peak of the specific heat $\beta(C_{V,\max})$ or the grand canonically calculated $T_C \sim N^{1/3}$ confirms our approach. In Fig. 9, three possible definitions of the critical temperature are given that all coincide in the thermodynamic limit. All definitions show a $\beta \sim N^{-\rho}$ dependence with ρ ranging between $\frac{2}{5}$ and $\frac{1}{3}$.

IV. CONCLUSION

Starting with the old ideas of Yang and Lee and Grossmann *et al.*, we have developed a classification scheme for phase transitions in finite systems. Based on the analytic continuation of the inverse temperature β into the complex

FIG. 9. Comparison between three different approaches to define a critical temperature for phase transitions in finite systems.

plane, we have shown the advantages of this approach. The distribution of the so-called Fisher zeros β_k draws enlightening pictures even for small systems, whereas the usually referred to thermodynamic properties such as the specific heat fail to classify the phase transitions properly. The classification scheme presented in this paper enables us to name the order of the transition in a nonambiguous way. The complex parts τ_k of the zeros β_k resemble times for which a whole ensemble of identical systems under consideration in a heat bath with an initial Boltzmann distribution loses its memory.

We have applied this to ideal noninteracting Bose gases confined in power-law traps. We have found that the order of the phase transition sensitively depends on the single-particle density of states generated by the confining potential. The distribution of zeros exactly reveals the order of the phase transition in finite systems.

-
- [1] S. Bose, Z. Phys. **26**, 178 (1924); A. Einstein, Sitzungsber. K. Preuss. Akad. Wiss. **1925**, 3 (1925).
 - [2] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Science **269**, 198 (1995); C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, Phys. Rev. Lett. **75**, 1687 (1995); K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, *ibid.* **75**, 3969 (1995).
 - [3] D. Stamper-Kurn, M. R. Andrews, A. P. Chikkatur, S. Inouye, H.-J. Miesner, J. Stenger, and W. Ketterle, Phys. Rev. Lett. **80**, 2027 (1998).
 - [4] P. Borrmann, J. Harting, O. Mülken, and E. Hilf, Phys. Rev. A **60**, 1519 (1999).
 - [5] S. Grossmann and M. Holthaus, Phys. Rev. Lett. **79**, 3557 (1997); S. Grossmann and M. Holthaus, Opt. Express **1**, 262 (1997); P. Navez *et al.*, Phys. Rev. Lett. **79**, 1789 (1997); M. Gajda and K. Razazewski, *ibid.* **78**, 2686 (1997).
 - [6] P. Borrmann and G. Franke, J. Chem. Phys. **98**, 2484 (1993).
 - [7] S. Grossmann and W. Rosenhauer, Z. Phys. **207**, 138 (1967); **218**, 437 (1969); S. Grossmann and W. Rosenhauer, *ibid.* **218**, 449 (1969).
 - [8] M. E. Fisher, in *Lectures in Theoretical Physics*, edited by W. E. Brittin (University of Colorado Press, Boulder, 1965), Vol. 7c.
 - [9] P. Borrmann, O. Mülken, and J. Harting, Phys. Rev. Lett. **84**, 3511 (2000).
 - [10] V. Bagnato and D. Kleppner, Phys. Rev. A **44**, 7439 (1991).
 - [11] H. Perez Rojas, Phys. Lett. A **234**, 13 (1997); S. Pearson, T. Pang, and C. Chen, Phys. Rev. A **58**, 4811 (1998); Z. Yan, *ibid.* **59**, 4657 (1999).
 - [12] A. I. Safonov *et al.*, Phys. Rev. Lett. **81**, 4545 (1998).
 - [13] C. N. Yang and T. Lee, Phys. Rev. **97**, 404 (1952); **87**, 410 (1952).
 - [14] W. J. Mullin, J. Low Temp. Phys. **106**, 1405 (1997); **110**, 167 (1997); F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999).
 - [15] N. L. Balazs and T. Bergeman, Phys. Rev. A **58**, 2359 (1998).
 - [16] S. Grossmann and M. Holthaus, Z. Naturforsch. A. Phys. Sci. **50**, 921 (1995).