

## On the set of divisors of a number

***Citation for published version (APA):***

Bruijn, de, N. G., Ebbenhorst Tengbergen, van, C., & Kruyswijk, D. (1951). On the set of divisors of a number. *Nieuw Archief voor Wiskunde, serie 2, 23*, 191-193.

***Document status and date:***

Published: 01/01/1951

***Document Version:***

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

***Please check the document version of this publication:***

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

***General rights***

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

***Take down policy***

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

# ON THE SET OF DIVISORS OF A NUMBER

BY

N. G. DE BRUIJN,

(Delft)

C<sup>A</sup>. VAN EBBENHORST TENGBERGEN

(Den Haag)

and D. KRUYSWIJK

(Amsterdam)

Theorem 1 below was stated as a conjecture and proposed as a prize problem of the Wiskundig Genootschap for 1949. Correct solutions were given by the second and the third author. The proof presented here depends on theorem 2, which is of some interest in itself. The theorems are purely combinatorial, but they are stated in an arithmetical form for the sake of a simple terminology.

If  $m$  is a natural number, and if  $m = p_1^{\lambda_1} \dots p_v^{\lambda_v}$  is its canonical factorisation, then the sum  $\lambda_1 + \dots + \lambda_v$  will be called the *degree* of  $m$ .

Consider the set  $U$  of all divisors of a fixed number  $m$  whose degree equals  $[\frac{1}{2}n]$ , where  $n$  is the degree of  $m$ . It is clear that no element of this set divides any other element of the set. Theorem 1 expresses that the set just considered is a maximal set with this property.

**Theorem 1.** *Let  $m$  be a natural number, and let  $n$  be its degree. Let  $d_1, \dots, d_l$  be a set of divisors of  $m$  with the property that no  $d_i$  divides any  $d_j$  ( $i \neq j$ ). Then we have  $l \leq L$ , where  $L$  is the number of divisors of  $m$  whose degree equals  $[\frac{1}{2}n]$ .*

The special case of squarefree  $m$ , that is,  $m$  is the product of  $n$  different primes, is a theorem of E. SPERNER<sup>1)</sup>.

<sup>1)</sup> Math. Zeitschrift 27, 544—548 (1928).

A sequence  $d_1, \dots, d_n$  of divisors of  $m$  will be called a *symmetrical chain* whenever it joins the following properties:

- a. The degree of  $d_1$  equals the one of  $m/d_n$ .
- b. For  $1 \leq i < h$  the quotient  $d_{i+1}/d_i$  is a prime <sup>2)</sup>.

**Theorem 2.** *The set of divisors of  $m$  can be completely divided into a number of disjoint symmetrical chains.*

**Proof.** We proceed by induction with respect to the number  $\nu(m)$  of different primes dividing  $m$ . For  $\nu(m) = 0$  the theorem is trivial. Now assume that the theorem has been proved for the number  $m$ ; we proceed to prove it for the number  $m_1 = m\phi^\lambda$  where  $\phi$  does not divide  $m$ .

Let  $d_1, \dots, d_n$  be a symmetrical chain of divisors of  $m$ . Then consider the divisors  $d_i\phi^\alpha$  ( $i = 1, \dots, n; \alpha = 0, \dots, \lambda$ ) of the number  $m_1$ . Our induction will be accomplished by showing that this set can be sub-divided into symmetrical chains of divisors of  $m_1$ . This is exhibited by the following scheme:

$$\begin{array}{c}
 d_1 \\
 d_1\phi \\
 \vdots \\
 \vdots \\
 d_1\phi^{\lambda-1} \\
 d_1\phi^\lambda
 \end{array}
 \left| \begin{array}{c}
 d_2 \\
 d_2\phi \\
 \vdots \\
 \vdots \\
 d_2\phi^{\lambda-1} \\
 d_2\phi^\lambda
 \end{array} \right.
 \begin{array}{c}
 \dots\dots d_n \\
 \dots\dots d_n\phi \\
 \vdots \\
 \vdots \\
 d_n\phi^{\lambda-1} \\
 d_n\phi^\lambda
 \end{array}$$

It is easily seen that the sequences  $\{d_1, d_1\phi, \dots, d_1\phi^\lambda, d_2\phi^\lambda, \dots, d_n\phi^\lambda\}$ ,  $\{d_2, d_2\phi, \dots, d_2\phi^{\lambda-1}, d_3\phi^{\lambda-1}, \dots, d_n\phi^{\lambda-1}\}$ , etc. are symmetrical chains.

**Proof of theorem 1.** Let the set of divisors of  $m$  be divided into symmetrical chains, and let  $U$  be the set of the  $L$  divisors of  $m$  of degree  $[\frac{1}{2}n]$ . Any element of  $U$  is contained in just one chain, and from the definition of the chain it follows that each chain contains just one element of  $U$ . Consequently the number of chains is  $L$ . Finally, if the set  $d_1, \dots, d_n$  has the property mentioned in theorem 1, it is

<sup>2)</sup> If  $h = 1$ , the latter condition vanishes. Hence, if  $n$  is even, any divisor of degree  $\frac{1}{2}n$  is a symmetrical chain of one element.

clear that each chain contains at most one of these  $d$ 's. Hence  $l \leq L$ .

The following theorem is a simple application of theorem 2. A set  $S$  of divisors of  $m$  will be called *convex* whenever

$$d_1 \in S, d_2 \in S, d_1 | d_3 | d_2 \text{ imply } d_3 \in S.$$

**THEOREM 3.** Let  $\nu$  be the number of different primes dividing  $m$ , and let  $S$  be a convex set of divisors of  $m$ . Then we have, if  $\mu$  indicates the Möbius function:

$$\left| \sum_{d \in S} \mu(d) \right| \leq \binom{\nu}{\lfloor \frac{1}{2}\nu \rfloor}.$$

**PROOF.** Since  $\mu(d) = 0$  whenever  $d$  is not squarefree, we may restrict ourselves to the case that  $m$  is squarefree.

Thus  $\nu$  is the degree of  $m$ , and clearly  $L = \binom{\nu}{\lfloor \frac{1}{2}\nu \rfloor}$  is the number of elements in the set  $U$ . Hence it equals the number of chains into which the set of divisors of  $m$  can be divided. Now write  $S = S_1 + \dots + S_L$ , where  $S_i$  is a subset of the  $i$ -th chain.  $S$  being convex, we infer that each  $S_i$  is either empty or it consists of a number of consecutive elements of the chain. If  $d$  runs through the elements of a chain, then  $\mu(d)$  assumes the values  $+1$  and  $-1$  alternately. Hence  $\sum \mu(d) = 0$  or  $1$  or  $-1$  whenever  $d$  runs through any set  $S_i$ . Finally

$$\left| \sum_S \mu(d) \right| \leq \sum_{i=1}^L \left| \sum_{S_i} \mu(d) \right| \leq L.$$

The special case where  $S$  consists of all divisors of  $m$  belonging to some interval  $a \leq d \leq b$ , was established by P. ERDÖS<sup>3)</sup>. His argument can also be used in order to prove the general case. To this end one has to add the following remark: if  $S$  is a convex subset of the set  $U$  of all divisors of  $m$ , then we can find two convex sets  $V_1$  and  $V_2$  such that  $1 \in V_1$ ,  $m \in V_2$ ,  $V_1 \cap V_2 = S$ ,  $V_1 \cup V_2 = U$ .

(Ingekomen 6.10.50).

<sup>3)</sup> Math. Student **17**, 32—33 (1950).