Magnonic black holes

Citation for published version (APA):

Roldán-Molina, A., Nunez, A. S., & Duine, R. A. (2017). Magnonic black holes. *Physical Review Letters*, 118(6), 061301. [061301]. <https://doi.org/10.1103/PhysRevLett.118.061301>

DOI:

[10.1103/PhysRevLett.118.061301](https://doi.org/10.1103/PhysRevLett.118.061301)

Document status and date:

Published: 08/02/2017

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Magnonic Black Holes

A. Roldán-Molina,^{1,2} Alvaro S. Nunez,² and R. A. Duine^{3,4}

¹*Centro para el Desarrollo de la Nanociencia y la Nanotecnología, CEDENNA, Avenida Ecuador 3493, Santiago 9170124, Chile*

²*Departamento de Física, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Casilla 487-3, Santiago, Chile*

³*Institute for Theoretical Physics, Utrecht University, Leuvenlaan 4, 3584 CE Utrecht, Netherlands*

⁴*Department of Applied Physics, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, Netherlands*

(Received 19 October 2016; published 8 February 2017)

We show that the interaction between the spin-polarized current and the magnetization dynamics can be used to implement black-hole and white-hole horizons for magnons—the quanta of oscillations in the magnetization direction in magnets. We consider three different systems: easy-plane ferromagnetic metals, isotropic antiferromagnetic metals, and easy-plane magnetic insulators. Based on available experimental data, we estimate that the Hawking temperature can be as large as 1 K. We comment on the implications of magnonic horizons for spin-wave scattering and transport experiments, and for magnon entanglement.

DOI: 10.1103/PhysRevLett.118.061301

Introduction.—Hawking’s 1974 postulation [1] that black holes evaporate by radiating particles with a thermal spectrum has triggered an enormous amount of scientific research and debate. It showed that black holes have a temperature—now called the Hawking temperature—confirming earlier ideas by Bekenstein on black-hole entropy and black-hole thermodynamics [2]. Computing the black-hole entropy from a microscopic statistical-physics description has been a key test of candidates for quantum-gravity theories ever since [3]. At the foundational level, the scaling of the black-hole entropy with area rather than volume led to the formulation of the so-called holographic principle [4] and to debates concerning the black-hole information paradox [5].

Despite these developments, Hawking radiation from gravitational black holes has not been observed yet. This is in part due to the low Hawking temperatures associated with astronomical black holes. Creating small black holes—which should have a higher Hawking temperature—seems experimentally impossible, and, if one were to succeed, they would evaporate rapidly. To circumvent such problems and to shed light on conceptual issues in the theoretical treatment of Hawking radiation—such as the so-called trans-Planckian problem—Unruh [6] suggested experimentally creating black-hole-horizon analogues. A black-hole horizon for sound waves in a flowing medium is created by a transition from subsonic to supersonic flow, such that waves along the flow and incoming from the subsonic region cannot escape from the supersonic region. A white-hole horizon is then a region where the flow changes from supersonic to subsonic. In this case, a wave traveling against the flow from the subsonic region cannot penetrate the supersonic part. Unruh’s original proposal concerned waves in water and motivated experiments in that direction [7]. This system cannot be driven into the quantum regime where the temperature is much lower than

the Hawking temperature. Nonetheless, by measuring the energy dependence of reflection and transmission amplitudes of waves scattering off the horizon in the classical regime, the Hawking spectrum can be determined up to normalization as the underlying physics is linear. This was experimentally implemented in Ref. [8].

Unruh’s work motivated theoretical proposals for black-hole-horizon analogues based on different systems in different regimes [9,10]. These include theoretical proposals for superfluid helium [11], atomic Bose-Einstein condensates [12], light in dispersive media [13], electromagnetic waveguides [14], ultracold fermions [15], trapped-ion rings [16], exciton-polariton condensates [17], light in nonlinear liquids [18], and, most recently, Weyl semimetals [19]. Experimental observations of various aspects of horizons have been reported for Bose-Einstein condensates [20], optical systems [21], and exciton-polariton condensates [22]. The essential ingredients for analogue horizons are linearly dispersing waves at long wavelengths and a background flow velocity which can exceed the velocity of the waves.

In this Letter, we propose a solid-state realization of a black-hole-horizon analogue. We outline how to use spin transfer torques, i.e., torques arising from the interplay between the spin current and the magnetization dynamics [23], to implement a black hole for magnons—the quanta of spin waves. In short, our proposal is based on the result that a spin-polarized electric current through a magnetic conductor interacts with the magnetization dynamics to give the spin waves a Doppler shift with effective “spin-drift” velocity v_s —as was experimentally detected in Ref. [24]. “Supersonic” and “subsonic” regions are then regions where the velocity v_s is larger and smaller (in absolute value) than the spin-wave velocity c , respectively.

Our proposal is distinct from other implementations in that the background flow for the excitations is not provided

by a moving medium but rather by a separate “fluid”—the spin current—that is controlled electrically and interacts with the magnetization and its excitations. (The interaction between magnons and spin superfluid currents was discussed in the context of magnon condensates in Ref. [25].) In addition, the dissipation and, in particular, the dissipative coupling of the spin current to the magnetization—or, in the language of analogue gravity, the dissipative coupling between excitations and background flow—is well understood, which facilitates understanding its interplay with Hawking radiation. Moreover, the interaction between solitonic excitations—magnetic domain walls—and the spin current allows for control over the position of domain walls as has been demonstrated experimentally with the long-term goal of building the magnetic “racetrack” memory [26]. This allows for a controlled study of the interaction between domain walls and the magnonic horizon.

From a practical point of view, this system is attractive, as it can be embedded in a device, can be electrically contacted, and has properties that are controlled by magnetic fields and electrical currents. While this facilitates experiments, a magnonic black-hole horizon may, in the longer term, also serve as an on-chip resource of entangled magnons for magnon-based quantum computation and information purposes [27]—as pairs of Hawking particles emitted from the horizon are entangled [1,28]. This was experimentally demonstrated very recently in Bose-Einstein condensates [20]. Below, we outline our proposals, provide estimates for their Hawking temperatures, and comment on experimental implications in the classical and quantum regime. We discuss two systems: ferromagnetic metals and antiferromagnetic metals. In the first of these, experiments have advanced furthest, while experiments on antiferromagnetic metals are rapidly catching up. In the Supplemental Material [29], we also discuss a proposal based on magnetic insulators.

Easy-plane ferromagnetic metal.—We consider a ferromagnetic metal far below its Curie temperature, such that the unit vector $\mathbf{n}(\mathbf{x}, t)$ along the direction of the magnetic order parameter is the appropriate degree of freedom at low energies and long wavelengths. Spintronics research over the past decade [23,35–48] has established that, in the presence of a steady-state transport current, it obeys the Landau-Lifshitz-Gilbert equation with spin transfer torques given by

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_s \cdot \nabla\right) \mathbf{n} - \mathbf{n} \times \mathbf{H}_{\text{eff}} = -\alpha \mathbf{n} \times \left(\frac{\partial}{\partial t} + \frac{\beta}{\alpha} \mathbf{v}_s \cdot \nabla\right) \mathbf{n}, \quad (1)$$

provided that intrinsic spin-orbit coupling is not very strong. In this equation, the velocity $\mathbf{v}_s = -gP\mu_B \mathbf{j}/2eM_s$, which is proportional to the electrical transport-current density $\mathbf{j}$ parametrizes the reactive and dissipative spin transfer torques, corresponding to the terms proportional to

$\mathbf{v}_s$ on the left- and right-hand sides of the above equation, respectively. Here, g is the Landé factor, P the spin polarization of the current, μ_B the Bohr magneton, e minus the electric charge, and M_s the saturation magnetization. The Gilbert damping parameter is given by α . Usually, the dissipative coefficients $\beta \sim \alpha$ because of approximate Galilean invariance, and they are of the order 10^{-2} . The above equation accurately describes experiments on current-driven domain wall motion in permalloy and other magnetic materials [49–55], and it also predicts the spin-wave Doppler shift that was measured in Ref. [24].

The effective field $\mathbf{H}_{\text{eff}} = -\delta E/(\hbar \delta \mathbf{n})$ is determined as the functional derivative of the energy $E[\mathbf{n}]$ and acquires contributions from exchange, anisotropies, and external fields. Here, we consider an easy-plane configuration and a field in the z direction such that

$$E[\mathbf{n}] = \int \frac{d\mathbf{x}}{a^3} \left(-\frac{J_s}{2} \mathbf{n} \cdot \nabla^2 \mathbf{n} + \frac{K}{2} n_z^2 + B n_z \right), \quad (2)$$

with a^3 being the volume of a unit cell, J_s the spin stiffness, and B the external field (absorbing all prefactors). Finally, K is the anisotropy constant that enforces the easy-plane anisotropy. Minimization of this energy yields a magnetization direction $\mathbf{n} = -\hat{z}$, with $\hat{z}$ being the unit vector in the z direction, for fields $B > K$. In this regime the magnons disperse quadratically and have a gap $\sim B - K$. For $B < K$ the magnetization direction deviates from the z direction and acquires a component in the $x - y$ plane. In that case we find that $n_z = 1 - B/K$, with the $x - y$ component determined by normalization. This latter so-called polar phase can be interpreted as a Bose-Einstein-condensed phase of magnons [56] and will turn out to have linearly dispersing magnons.

The Landau-Lifshitz-Gilbert equation is rewritten as a dissipative Gross-Pitaevskii equation by introducing the complex field $\psi(\mathbf{x}, t)$ by means of $\mathbf{n} = (\sqrt{\mu_B/2M_s} \text{Re}[\psi], \sqrt{\mu_B/2M_s} \text{Im}[\psi], \mu_B |\psi|^2/M_s)$, which corresponds to a classical linearized version of the usual Holstein-Primakoff transformation. We find that

$$i\hbar \left(\frac{\partial}{\partial t} + \mathbf{v}_s \cdot \nabla \right) \psi = (-J_s \nabla^2 - \mu + g|\psi|^2) \psi + \hbar \alpha \left(\frac{\partial}{\partial t} + \frac{\beta}{\alpha} \mathbf{v}_s \cdot \nabla \right) \psi, \quad (3)$$

with the chemical potential $\mu = K - B$ and the contact interaction $g = K\mu_B/M_s$. In the polar phase when $B < K$, and thus $\mu > 0$, we insert $\psi = \sqrt{n_c} + \delta\psi$, with $n_c = \mu/g$ being the effective condensate density, into the above. Linearizing with respect to $\delta\psi$ and $\delta\psi^*$ leads to two coupled equations for $\delta\psi$ and $\delta\psi^*$. Using the Bogoliubov ansatz, we write $\delta\psi = u(\mathbf{x})e^{-i\omega t} - v^*(\mathbf{x})e^{+i\omega t}$. The resulting equations are, up to the

dissipative corrections proportional to α and β , equivalent to the equation describing Bogoliubov excitations on top of a Bose-Einstein condensate flowing with the velocity $\mathbf{v}_s$ [10,12,29]. Note, however, that here the velocity is given not by a superflowing condensate but by the electrons providing the nonzero charge current and the resulting nonzero spin current. Taking, for the moment, $\mathbf{v}_s$ to be constant, we find in the long-wave limit and to leading order in α and β the magnon dispersion relation $(\omega_{\mathbf{k}} - \mathbf{v}_s \cdot \mathbf{k}) = c\mathbf{k} - i\alpha c\mathbf{k} - i(\alpha - \beta)\mathbf{v}_s \cdot \mathbf{k}$, with the spin-wave velocity $c = \sqrt{2J_s(K - B)}/\hbar$.

Black-hole and white-hole horizons are now implemented by regions where the velocity $\mathbf{v}_s$ changes from $|\mathbf{v}_s| < c$ to $|\mathbf{v}_s| > c$, and vice versa. We specifically consider the setups in Fig. 1 that involve a wire geometry. A current density is flowing from right to left, such that the velocity $\mathbf{v}_s$ is typically pointing from left to right (depending on the sign of the spin polarization P). A narrow region in a wire leads to an increase in the current density, and thus in $|\mathbf{v}_s|$ with respect to a wider region. If the current density in this narrow region is such that in the narrow region $|\mathbf{v}_s| > c$, while in the wider regions of the wire we have $|\mathbf{v}_s| < c$, there will be a black-hole horizon for magnons coming from the wide region on the left (traveling “downstream,” i.e., to the right) in Fig. 1(a). Similarly, there

will be a white-hole horizon for magnons traveling to the left from the right “upstream” region in Fig. 1(b). A dent in the wire creates a pair of horizons, a black and a white one [see Fig. 1(c)].

Since α and β are small, we ignore, in the first instance, dissipation. We focus in the following on black-hole horizons. The Hawking temperature T_H of the black-hole horizon is then given by $k_B T_H = \frac{\hbar}{2\pi} \partial(|\mathbf{v}_s - c|)/\partial r$ [10], where the derivative is taken at the horizon and in the direction perpendicular to it, and where k_B is Boltzmann’s constant. Taking a typical value of $J_s = 10^{-39}$ J m⁻² for the exchange interactions, and B/k_B and K/k_B to be on the order of 1 K [26], we estimate $c \sim 10^3$ m/s, although it can be made arbitrarily small by tuning $B \uparrow K$. The critical current density j_c required for $|\mathbf{v}_s|$ to exceed c is $j_c \sim M_s |e| c / \mu_B \sim 10^{11}$ A/m², where we took $g \sim P \sim 1$, $M_s / \mu_B \sim 1$ nm⁻³, and $c = 1000$ m/s. We note here that such large, or even larger, current densities are quite common in experiments on current-driven domain wall motion [26]. Assuming now that the current density changes over a length scale of a $d = 1$ nm—which can be achieved by nanofabrication techniques—we find that $T_H \sim \hbar c / k_B d \sim 1$ K. At zero temperature, pairs of magnons are created with one magnon being absorbed by the black hole. The black-hole horizon will emit magnons with a thermal spectrum determined by T_H into the subsonic region left of the magnonic black-hole horizon in Fig. 1(a). Of course, the current density leads to an increase in temperature because of Joule heating such that zero or small temperatures are difficult to achieve. By tuning the field B to approach K , one can lower c and the required critical current j_c . The Hawking temperature will go down accordingly, but the Joule heating is quadratic in temperature, whereas the change in Hawking temperature is linear, allowing disentanglement of both effects.

There are also signatures of the physics of Hawking radiation in the classical regime, i.e., at temperatures $T \gg T_H$, as the underlying processes are linear. Following the arguments of Ref. [8], we have it that the ratio of spin-wave transmission (t) and reflection (r) amplitudes off the black-hole horizon is given by

$$\frac{|t(\omega)|^2}{|r(\omega)|^2} = \exp\left(-\frac{\hbar\omega}{k_B T_H}\right). \quad (4)$$

Spin-wave scattering experiments are standard in the field of magnonics [57] and may thus provide a first step towards observing the nontrivial features of magnonic black-hole horizons. The presence of the horizon itself can, of course, also be detected with a spin-wave scattering experiment.

Using the above expression for the transmission and reflection coefficients, along with standard Landauer-Büttiker expressions for magnon transport properties, we find that magnon transport coefficients are proportional to $I_n = \int d\epsilon \epsilon^n |t(\epsilon)|^2 (-\partial n_B / \partial \epsilon)$, with $n_B(\epsilon) = [e^{\epsilon/k_B T} - 1]^{-1}$

FIG. 1. Setup for creating magnonic black-hole and white-hole horizons. A narrow region of a wire that carries a steady-state current leads to an increased current density such that the background velocity $\propto -\mathbf{j}$ exceeds the magnon velocity if the current exceeds j_c . (a) A magnonic black-hole (MBH) horizon for magnons incoming from the left. (b) A magnon white-hole (MWH) horizon for magnons incoming from the right. (c) A pair of MBH and MWH horizons. Incoming and scattered spin waves are illustrated.

being the Bose-Einstein distribution function at the temperature T . Here, I_0 is proportional to the spin conductance and I_1 to the magnon contribution to the heat conductance. Using Eq. (4) and a conservation of the norm, we conclude that, at low temperatures, the transport coefficients behave as if the actual temperature T is replaced by T^* , with $1/T^* = 1/T + 1/T_H$, so that $T^*/T = (T_H/T)/(1 + T_H/T)$. For the purpose of this estimate, we have replaced the Bose-Einstein distribution function by the Boltzmann one. When $T_H \ll T$ the transport coefficients thus behave as if the temperature is equal to T_H , while in the opposite limit $T^* \approx T$. This may provide a transport signature of the Hawking radiation.

When $|\mathbf{v}_s| > c$ the ferromagnetic ground state may become unstable towards the formation of a modulated state [40,41]. This will not affect the physics in the subsonic region (i.e., left of the black-hole horizon in Fig. 1). Taking into account the dissipative terms, we find that the magnons are linearly stable, however, when $|(\alpha - \beta)\mathbf{v}_s| < \alpha c$, which provides a large window for stability since α and β are usually approximately equal.

Isotropic antiferromagnetic metal.—Our next proposal concerns an isotropic antiferromagnetic metal and may also be implemented using a synthetic antiferromagnet—i.e., two ferromagnetic layers separated by a normal metal—provided the interlayer exchange coupling is sufficiently strong. The interaction between spin-polarized current and the magnetization dynamics in antiferromagnets has been studied theoretically over the past decade [58–60]. Recently, electrical switching of an antiferromagnet was reported using strong spin-orbit coupling [61,62]. In the opposite limit of strong exchange interactions between electron spins and magnetization, the equation of motion for the Néel vector $\mathbf{n}$ of the antiferromagnet is given by [58–60]

$$\mathbf{n} \times [\ddot{\mathbf{n}} - c_a^2 \nabla^2 \mathbf{n} + (\mathbf{v} \cdot \nabla)^2 \mathbf{n} + 2(\mathbf{v} \cdot \nabla) \dot{\mathbf{n}}] = \mathbf{n} \times \mathbf{h}_d, \quad (5)$$

where the velocity $\mathbf{v}$ plays the same role for antiferromagnets as the velocity $\mathbf{v}_s$ does for ferromagnets. Furthermore, $\mathbf{h}_d \propto \alpha[\dot{\mathbf{n}} + \beta(\mathbf{v} \cdot \nabla)\mathbf{n}/\alpha]$ describes relaxation and will, as with ferromagnets, be ignored in the first instance since it is usually small. Generally, we have it that $\mathbf{v} \propto \mathbf{j}$, with the prefactor determined by microscopic physics. Estimates [59,63] show that velocities similar to the case of ferromagnets can be obtained; i.e., $|\mathbf{v}|$ can be on the order of 1000–10000 m/s for current densities $j \sim 10^{11-12}$ A/m². Since the antiferromagnetic spin-wave velocity c_a is on the order of $c_a \sim 1000$ m/s, we conclude that black-hole horizons can be created for magnons in antiferromagnets, as $|\mathbf{v}|$ can exceed c_a .

Linearizing Eq. (5) around a collinear state $\mathbf{n}_0$ by means of $\mathbf{n} = \mathbf{n}_0 + \delta\mathbf{n}$, we find that

$$\frac{1}{c_a^2} \left(\frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right)^2 \Phi - \nabla^2 \Phi = 0, \quad (6)$$

where $\Phi = \mathbf{n}_0 \times \delta\mathbf{n}$. This equation shows that antiferromagnetic magnons interacting with a transport current are described analogously to sound waves propagating in a medium with nonzero velocity, albeit antiferromagnetic magnons have two polarizations. Following the arguments of Ref. [6] and considering the setup in Fig. 1, we find the same expression for the Hawking temperature as for the ferromagnetic metal (with c replaced by c_a and $\mathbf{v}_s$ replaced by $\mathbf{v}$). The possibilities for experimental detection in the classical and quantum regimes are also similar.

Discussion, conclusion and outlook.—In conclusion, we have shown that the interaction between spin current and magnetization dynamics can give rise to black-hole and white-hole horizons for magnons. For the metallic easy-plane ferromagnet, we have discussed the effect of relaxation and how it stabilizes the homogeneous magnetic ground state. While an extensive investigation into the effects of dissipation in the quantum regime is beyond the scope of this paper, we expect that it gives rise to a characteristic length scale $\hbar c / \alpha k_B T$ over which the system needs to be quantum coherent to observe spontaneous magnon pair creation. In our discussions, we have neglected the effects of unwanted anisotropies that give the magnons a gap. Such anisotropies can be neglected as long as the gap is smaller than $k_B T_H$. Given the experimental control over anisotropies by doping, sample shape, and material composition, we expect that this will not pose a severe limitation. For example, in a thin wide wire, the easy-plane anisotropy—required for obtaining the linear dispersion at long wavelengths—is much larger than the unwanted shape anisotropy that ultimately pins the magnetization within the easy plane.

One of the most interesting aspects of Hawking radiation is that the emitted particle pairs are entangled. For our case of magnetic systems, the quantity

$$\langle \hat{S}_{-k}^- \hat{S}_k^- \rangle \langle \hat{S}_{-k}^+ \hat{S}_k^+ \rangle - \langle \hat{S}_k^+ \hat{S}_k^- \rangle \langle \hat{S}_{-k}^+ \hat{S}_{-k}^- \rangle \quad (7)$$

exceeds its classical value of zero if the emitted pairs are entangled [28]. Here, $\hat{S}_k^+$ and $\hat{S}_k^-$ are the usual spin raising and lowering operators at magnon momentum $\mathbf{k}$, and the magnons forming the pairs are emitted with momenta $+\mathbf{k}$ and $-\mathbf{k}$. Using the results of Ref. [64], we expect that the above correlation function can, in principle, be measured from a spin-spin correlation function, e.g., by neutron scattering. Another possibility is to detect the magnon entanglement via spin current noise [65]. In future work we will investigate possible quantum-information devices exploiting the entanglement between the magnonic Hawking partners. Other interesting directions for future research include the interaction of ferromagnetic solitons, i.e., domain walls, with the horizons, the inclusion of strong

spin-orbit coupling, and the development of a transport theory that treats the magnon transport beyond the estimates made here.

R. D. would like to acknowledge a discussion with W. G. Unruh during his Kramers professorship at the Institute for Theoretical Physics in Utrecht that contributed to the ideas presented here. We also acknowledge helpful interactions with Erik van der Wurff, Henk Stoof, and Stefan Vandoren. R. D. is a member of the Delta Institute for Theoretical Physics (D-ITP) consortium, a program of the Netherlands Organisation for Scientific Research (NWO) that is funded by the Dutch Ministry of Education, Culture and Science (OCW). This work is in part funded by the Stichting voor Fundamenteel Onderzoek der Materie (FOM). A. S. N. would like to acknowledge funding from Fondecyt Grant No. 1150072 and support from Financiamiento Basal para Centros Científicos y Tecnológicos de Excelencia (Chile), under Project No. FB 0807.

-
- [1] S. W. Hawking, *Nature (London)* **248**, 30 (1974); *Commun. Math. Phys.* **43**, 199 (1975).
- [2] J. D. Bekenstein, *Phys. Rev. D* **7**, 2333 (1973).
- [3] See, for example, A. Strominger and C. Vafa, *Phys. Lett. B* **379**, 99 (1996).
- [4] L. Susskind, *J. Math. Phys. (N.Y.)* **36**, 6377 (1995); G. 't Hooft, in *Proceedings of the Conference on Highlights of Particle and Condensed Matter Physics (SALAMFEST), Trieste, Italy, 1993*, edited by A. Ali, J. Ellis, and S. Randjbar-Daemi (World Scientific, River Edge, NJ, 1993).
- [5] L. Susskind, *Nat. Phys.* **2**, 665 (2006).
- [6] W. G. Unruh, *Phys. Rev. Lett.* **46**, 1351 (1981).
- [7] G. Rousseaux, C. Mathis, P. Maïssa, T. Philbin, and U. Leonhardt, *New J. Phys.* **10**, 053015 (2008).
- [8] S. Weinfurter, E. W. Tedford, M. C. J. Penrice, W. G. Unruh, and G. A. Lawrence, *Phys. Rev. Lett.* **106**, 021302 (2011).
- [9] *Artificial Black Holes*, edited by M. Novello, M. Visser, and G. E. Volovik (World Scientific, Singapore, 2002).
- [10] R. Balbinot, in *Analogue Gravity Phenomenology*, edited by D. Faccio *et al.* (Springer, Heidelberg, 2013), Chap. 9.
- [11] T. A. Jacobson and G. E. Volovik, *Phys. Rev. D* **58**, 064021 (1998).
- [12] L. J. Garay, J. R. Anglin, J. I. Cirac, and P. Zoller, *Phys. Rev. Lett.* **85**, 4643 (2000).
- [13] U. Leonhardt and P. Piwnicki, *Phys. Rev. Lett.* **84**, 822 (2000).
- [14] R. Schützhold and W. G. Unruh, *Phys. Rev. Lett.* **95**, 031301 (2005).
- [15] S. Giovanazzi, *Phys. Rev. Lett.* **94**, 061302 (2005).
- [16] B. Horstmann, B. Reznik, S. Fagnocchi, and J. I. Cirac, *Phys. Rev. Lett.* **104**, 250403 (2010).
- [17] D. D. Solnyshkov, H. Flayac, and G. Malpuech, *Phys. Rev. B* **84**, 233405 (2011).
- [18] M. Elazar, V. Fleurov, and S. Bar-Ad, *Phys. Rev. A* **86**, 063821 (2012).
- [19] G. E. Volovik, *Pis'ma Zh. Eksp. Teor. Fiz.* **104**, 660 (2016).
- [20] J. Steinhauer, *Nat. Phys.* **10**, 864 (2014); *Nat. Phys.* (to be published).
- [21] T. G. Philbin, C. Kuklewicz, S. Robertson, S. Hill, F. König, and U. Leonhardt, *Science* **319**, 1367 (2008); F. Belgiorno, S. L. Cacciatori, M. Clerici, V. Gorini, G. Ortenzi, L. Rizzi, E. Rubino, V. G. Sala, and D. Faccio, *Phys. Rev. Lett.* **105**, 203901 (2010); W. G. Unruh and R. Schützhold, *Phys. Rev. D* **86**, 064006 (2012); S. Liberati, A. Prain, and M. Visser, *Phys. Rev. D* **85**, 084014 (2012).
- [22] H. S. Nguyen, D. Gerace, I. Carusotto, D. Sanvitto, E. Galopin, A. Lemaître, I. Sagnes, J. Bloch, and A. Amo, *Phys. Rev. Lett.* **114**, 036402 (2015).
- [23] For a review, see D. C. Ralph and M. D. Stiles, *J. Magn. Magn. Mater.* **320**, 1190 (2008).
- [24] V. Vlaminc and M. Bailleul, *Science* **322**, 410 (2008).
- [25] M. Clovecko, E. Gazo, M. Kupka, and P. Skyba, *J. Phys. Conf. Ser.* **150**, 032016 (2009).
- [26] S. Parkin and S.-H. Yang, *Nat. Nanotechnol.* **10**, 195 (2015).
- [27] S. N. Andrianov and S. A. Moiseev, *Phys. Rev. A* **90**, 042303 (2014).
- [28] X. Busch and R. Parentani, *Phys. Rev. D* **89**, 105024 (2014).
- [29] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.118.061301>, which includes Refs. [10,30–34], for more details on the derivation of the Bogoliubov equations and a discussion of the implementation of horizons in a magnetic insulator.
- [30] B. I. Halperin and P. C. Hohenberg, *Phys. Rev.* **188**, 898 (1969).
- [31] E. B. Sonin, *Adv. Phys.* **59**, 181 (2010).
- [32] Y. M. Bunkov and G. E. Volovik, *Int. ser. monogr. phys.* **156**, 253 (2013).
- [33] S. Takei and Y. Tserkovnyak, *Phys. Rev. Lett.* **112**, 227201 (2014).
- [34] S. Takei, B. I. Halperin, A. Yacoby, and Y. Tserkovnyak, *Phys. Rev. B* **90**, 094408 (2014).
- [35] J. C. Slonczewski, *J. Magn. Magn. Mater.* **159**, L1 (1996).
- [36] L. Berger, *Phys. Rev. B* **54**, 9353 (1996).
- [37] M. Tsoi, A. G. M. Jansen, J. Bass, W.-C. Chiang, M. Seck, V. Tsoi, and P. Wyder, *Phys. Rev. Lett.* **80**, 4281 (1998).
- [38] E. B. Myers, D. C. Ralph, J. A. Katine, R. N. Louie, and R. A. Buhrman, *Science* **285**, 867 (1999).
- [39] Y. B. Bazaliy, B. A. Jones, and S.-C. Zhang, *Phys. Rev. B* **57**, R3213 (1998).
- [40] J. Fernández-Rossier, M. Braun, A. S. Núñez, and A. H. MacDonald, *Phys. Rev. B* **69**, 174412 (2004).
- [41] G. Tatara and H. Kohno, *Phys. Rev. Lett.* **92**, 086601 (2004); **96**, 189702 (2006).
- [42] S. Zhang and Z. Li, *Phys. Rev. Lett.* **93**, 127204 (2004).
- [43] S. E. Barnes and S. Maekawa, *Phys. Rev. Lett.* **95**, 107204 (2005); S. E. Barnes, *Phys. Rev. Lett.* **96**, 189701 (2006).
- [44] Y. Tserkovnyak, H. J. Skadsem, A. Brataas, and G. E. W. Bauer, *Phys. Rev. B* **74**, 144405 (2006).
- [45] H. Kohno, G. Tatara, and J. Shibata, *J. Phys. Soc. Jpn.* **75**, 113706 (2006).
- [46] F. Piéchon and A. Thiaville, *Phys. Rev. B* **75**, 174414 (2007).
- [47] R. A. Duine, A. S. Núñez, J. Sinova, and A. H. MacDonald, *Phys. Rev. B* **75**, 214420 (2007).
- [48] G. Tatara, H. Kohno, and J. Shibata, *Phys. Rep.* **468**, 213 (2008).

- [49] J. Grollier, P. Boulenc, V. Cros, A. Hamzić, A. Vaurès, A. Fert, and G. Faini, *Appl. Phys. Lett.* **83**, 509 (2003).
- [50] M. Tsoi, R. E. Fontana, and S. S. P. Parkin, *Appl. Phys. Lett.* **83**, 2617 (2003).
- [51] A. Yamaguchi, T. Ono, S. Nasu, K. Miyake, K. Mibu, and T. Shinjo, *Phys. Rev. Lett.* **92**, 077205 (2004).
- [52] M. Kläui, C. A. F. Vaz, J. A. C. Bland, W. Wernsdorfer, G. Faini, E. Cambril, L. J. Heyderman, F. Nolting, and U. Rüdiger, *Phys. Rev. Lett.* **94**, 106601 (2005).
- [53] G. S. D. Beach, C. Knutson, C. Nistor, M. Tsoi, and J. L. Erskine, *Phys. Rev. Lett.* **97**, 057203 (2006).
- [54] M. Hayashi, L. Thomas, C. Rettner, R. Moriya, and S. S. P. Parkin, *Nat. Phys.* **3**, 21 (2007).
- [55] M. Yamanouchi, D. Chiba, F. Matsukura, and H. Ohno, *Nature (London)* **428**, 539 (2004).
- [56] B. Flebus, S. A. Bender, Y. Tserkovnyak, and R. A. Duine, *Phys. Rev. Lett.* **116**, 117201 (2016).
- [57] V. V. Kruglyak, S. O. Demokritov, and D. Grundler, *J. Phys. D* **43**, 264001 (2010).
- [58] A. S. Núñez, R. A. Duine, P. Haney, and A. H. MacDonald, *Phys. Rev. B* **73**, 214426 (2006).
- [59] A. C. Swaving and R. A. Duine, *Phys. Rev. B* **83**, 054428 (2011).
- [60] K. M. D. Hals, Y. Tserkovnyak, and A. Brataas, *Phys. Rev. Lett.* **106**, 107206 (2011).
- [61] J. Zelezny, H. Gao, K. Vyborny, J. Zemen, J. Masek, A. Manchon, J. Wunderlich, J. Sinova, and T. Jungwirth, *Phys. Rev. Lett.* **113**, 157201 (2014).
- [62] P. Wadley *et al.*, *Science* **351**, 587 (2016).
- [63] Y. Yamane, J. Ieda, and J. Sinova, *Phys. Rev. B* **94**, 054409 (2016).
- [64] J. Steinhauer, *Phys. Rev. D* **92**, 024043 (2015).
- [65] A. Kamra and W. Belzig, *Phys. Rev. B* **94**, 014419 (2016).