

On fiber orientation distribution peak selection for diffusion MRI fiber tractography

Citation for published version (APA):

Zhylka, A., Leemans, A., Pluim, J., & De Luca, A. (2019). *On fiber orientation distribution peak selection for diffusion MRI fiber tractography*. 14. Abstract from 36th Annual Scientific Meeting of the European Society for Magnetic Resonance in Medicine and Biology, (ESMRMB2019), Rotterdam, Netherlands.
<https://doi.org/10.1007/s10334-019-00753-3>

DOI:

[10.1007/s10334-019-00753-3](https://doi.org/10.1007/s10334-019-00753-3)

Document status and date:

Published: 01/09/2019

Document Version:

Accepted manuscript including changes made at the peer-review stage

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

On fiber orientation distribution peak selection for diffusion MRI fiber tractography

A. Zhylyka, A. Leemans, J. Plum, A. De Luca

Introduction

With diffusion MRI (dMRI) based fiber tractography (FT), brain white matter fiber pathways can be reconstructed in vivo and noninvasively². Recent work has shown that reliable and consistent computation of these fiber trajectories is challenging as there are various algorithms with different user-defined settings⁴. In particular, it is well known that probabilistic FT approaches can produce more false positive reconstructions than the deterministic ones. On the other hand, while deterministic FT approaches can sample dominant peak orientations from the fiber orientation distribution (FOD) in a robust way, they typically produce more false negative pathways (Fig. 1).

We developed a novel way to reconstruct pathways by considering a peak selection strategy that also integrates the orientations with a higher angular deviation. By incorporating prior knowledge of the location of seed and target regions of interest (ROI) while maintaining a deterministic view of FT, we can provide a more balanced tradeoff between true and false positive reconstructions.




Fig. 1: Current FT methods only use a part of the FOD peak information, reconstructing a subset of the pathways (blue). We aim to use the neglected information, to reconstruct the missing branches (red). Background image is from [5] (with permission).

Methods

The peak selection strategy introduced in this work is implemented for the constrained spherical deconvolution (CSD) based FT method¹ and consists of the following steps. Firstly, conventional CSD FT is performed (with FOD threshold of 0.1 and angular deviation threshold of 45°). Secondly, every point of each pathway that has more than one FOD peak is considered as a new seed, but now with the other peak orientations as the initial directions for reconstructing new pathways. This procedure can be performed recursively with predefined convergence criteria. Finally, the resulting pathways are selected based on prior knowledge of the anatomical location of the ROI in combination with the topology of the fiber tracts.

The dMRI data used in this work were acquired with a 3T MRI system: 60 volumes at $b=3000\text{s/mm}^2$, 6 volumes at $b=0\text{s/mm}^2$; 2.5mm isotropic voxel size. Visualizations are performed with *ExploreDTI*³.

Results

In Fig. 2a, a conventional pathway is reconstructed with the seed point defined in the spine area (only one seed point is taken). Fig. 2b shows the “unconstrained” result with the additional seed definition according to proposed strategy. Fig. 2c shows the final result with target ROI selected. Fig. 3 shows the results in a more realistic scenario with more seeds.




Fig. 2: (a) Example pathway with color-encoding according to number of fiber peaks along its trajectory. (b) pathways crossing the original pathway in (a). (c) Final result showing the additional branches (see white target ROI).




Fig. 3: (a) Conventional FT result (similar as Fig. 2(a), but more seed points). (b) The final result after including our proposed peak selection strategy. Notice the increase of true positives (lateral tract branching as indicated by the white arrow).

Conclusion

We developed a novel way to perform FT by considering a peak selection strategy that also integrates the orientations with a higher angular deviation and that incorporates prior knowledge of fiber tract anatomy. In doing so, we can obtain a more complete representation of brain pathways.

Acknowledgements

Andrey Zhylyka is supported by the funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 765148.

References:

1. Jeurissen B., et al. *Hum Brain Mapp*2011;32(3):461-79.
2. Jeurissen B., et al. *NMR Biomed*2019;32(4):e3785.
3. Leemans A., et al. *ISMRM* 2009;3537.
4. Maier-Hein K., et al. *Nat Comm*2017;8(1):1349.
5. Tournier J.-D., et al. *MRM*2011;65(6):1532-56.