

Direct numerical simulation of laminarization in the atmospheric boundary layer

Citation for published version (APA):

Donda, J. M. M., van de Wiel, B. J. H., van Heijst, G. J. F., & Clercx, H. J. J. (2020). *Direct numerical simulation of laminarization in the atmospheric boundary layer*. Paper presented at 14th European Turbulence Conference, ETC 2013, Lyon, France.

Document status and date:

Published: 01/01/2020

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

**DIRECT NUMERICAL SIMULATION
OF LAMINARIZATION
IN THE ATMOSPHERIC BOUNDARY LAYER**

Judith M. M. Donda¹, B.J.H. van de Wiel¹, G.J.F. Van Heijst¹, H.J.J. Clercx¹
¹Fluid Dynamics Lab., Eindhoven Technical University, Netherlands

Abstract A well-known phenomenon in the atmospheric boundary layer is the facts that winds may become very weak in the evening after a clear sunny day. In these quiet conditions usually hardly any turbulence is present. Consequently, this type of boundary layer is referred to as the quasi-laminar boundary layer. In spite of its omnipresence, the appearance of the laminar boundary layer is poorly understood and forms a long-standing problem in meteorological research. In the present study we investigate an analogue problem in the form of a stably stratified channel flow. The flow is studied by using direct numerical simulations (DNS). Simulations reveal that flow laminarization occurs when the normalized surface heat extraction h/L is larger than 1.23. In a companioning study this laminarisation is explained by the maximum sustainable heat flux theory (MSHF), which will be validated in the present research.

MODEL SET UP

In the present work we follow the setup of Nieuwstadt (2005) [1] and Flores and Riley (2011) [2]. A pressure driven non-stratified flow is suddenly cooled from below. The default Reynolds number is: $Re_* = u_* h / \nu = 360$, with $u_* \equiv \sqrt{-(1/\rho)(\partial P/\partial x)h}$. The surface heat flux H_0 is prescribed by imposing the external stability parameter h/L , with $h/L = (\kappa g h H_0) / (T_{ref} \rho c_p u_*^3)$. A schematic picture of the configuration is given in Figure 1 below.




Figure 1. schematic picture of the channel flow configuration with the pressure gradient force and prescribed surface heat extraction as external parameters. Decreasing temperature is indicated by increasing grey-scale.

In the DNS the Navier-Stokes equations in the Boussinesq approximation are solved on a rectangular grid. Numerical characteristics: the domain size is $5h$ in the horizontal directions and $1h$ in the wall-normal direction; Both Reynolds number ($360, 720, \dots$) and resolution ($100^3, 200^3, 400^3, \dots$) are varied. Periodic conditions in the horizontal directions are applied and the initial condition is a neutral, steady flow.

RESULTS

For a case with moderate surface cooling ($h/L=1$) the evolution of the mean velocity and temperature profiles is given in Figure 2.




Figure 2. velocity (right) and temperature (left) profiles at $t=0, 10, 20$ and 25 , for a dimensionless surface heat extraction $h/L=1$.

Because turbulent exchange of momentum is less efficient in stratified than in neutral conditions, large wind shears are needed to oppose the horizontal pressure gradient. The profile attains a log-linear shape resembling typical atmospheric flows [3,4]. The temperature profiles approach a quasi-steady equilibrium after $t \sim 20$.

The simulations were repeated for various values of the surface heat extraction. When the value of the surface heat extraction exceeded a critical value of $h/L=1.24$ turbulence in the flow cannot survive and the flow laminarizes. This is illustrated by Figures 3a,b and 4, below. In a companioning study this laminarisation is explained by the maximum sustainable heat flux theory (MSHF [3,4]), which will be validated in the present research.




Figure 3 Snapshot of temperature distribution a) in typical turbulent case (with surface heat extraction $h/L=1$) and a typical laminar case ($h/L=1.24$).




Figure 4. turbulent kinetic energy as a function of time. BLUE : surface heat extraction : $h/L=1.0$; RED : $h/L=1.24$.

References

- [1] F.T.M. Nieuwstadt. Direct numerical simulation of stable channel flow at large stability. *Boundary Layer Meteorology* **116**: 277–299, 2005.
- [2] O. Flores, J.J. Riley. Analysis of turbulence collapse in the stably stratified surface layer using direct numerical simulation. *Boundary Layer Meteorology* **139**: 241–259, 2011.
- [3] B. J. H. Van de Wiel, A.F. Moene, H.J.J. Jonker. The cessation of continuous turbulence as precursor of the very stable nocturnal boundary layer. *Journal of the Atmospheric Sciences* **69**: 3097–3115, 2012.
- [4] B. J. H. Van de Wiel, A.F. Moene, H.J.J. Jonker et al. The minimum wind speed for sustainable turbulence in the nocturnal boundary layer. *Journal of the Atmospheric Sciences* **69**: 3116–3127, 2012.