

# Hilbert spaces of harmonic functions in which differentiation operators are continuous or compact

**Citation for published version (APA):**

Liu, G. Z. (1989). *Hilbert spaces of harmonic functions in which differentiation operators are continuous or compact*. (RANA : reports on applied and numerical analysis; Vol. 8909). Technische Universiteit Eindhoven.

**Document status and date:**

Published: 01/01/1989

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

**Eindhoven University of Technology**  
**Department of Mathematics and Computing Science**


RANA 89-09

June 1989

**HILBERT SPACES OF HARMONIC  
FUNCTIONS IN WHICH DIFFERENTIATION  
OPERATORS ARE CONTINUOUS OR COMPACT**

by

Liu Gui-Zhong



Reports on Applied and Numerical Analysis  
Department of Mathematics and Computing Science  
Eindhoven University of Technology  
P.O. Box 513  
5600 MB Eindhoven  
The Netherlands

# HILBERT SPACES OF HARMONIC FUNCTIONS IN WHICH DIFFERENTIATION OPERATORS ARE CONTINUOUS OR COMPACT

Liu Gui-Zhong\*

Department of Mathematics and Computing Science

Eindhoven University of Technology

5600 MB Eindhoven

The Netherlands

## Abstract

Hilbert spaces of harmonic functions are presented wherein differentiation operators are continuous or even compact

## 1. Introduction

We all say that differentiation operators are unbounded or equivalently not continuous in a Hilbert or Banach space of functions. In fact, differentiation operators are a standard illustrative example in the undergraduate text books on functional analysis, which is used to show that there are unbounded operators in Banach spaces. In the present note, however, we are going to give some naturally arising Hilbert spaces of harmonic functions wherein the differentiation operators are continuous or even compact.

Let us fix a few notations.  $\mathbb{R}^q$  is the Euclidean space of dimension  $q \geq 2$  with inner product  $x \cdot y = \sum_{i=1}^q x_i y_i$  and corresponding norm  $|x| = (x \cdot x)^{1/2}$  for  $x, y \in \mathbb{R}^q$ .  $B^q(R) = \{x \mid x \in \mathbb{R}^q, |x| < R\}$ ;  $B^q \equiv B^q(1)$ .  $S^{q-1} = \{x \in \mathbb{R}^q \mid |x| = 1\}$  is the unit sphere in  $\mathbb{R}^q$ . The points on  $S^{q-1}$  are denoted  $\zeta, \eta$  et al. As usual  $L^2(S^{q-1})$  stands for the  $L^2$  space of functions defined on  $S^{q-1}$  with inner product

$$(u, v) = \int_{S^{q-1}} u(\zeta) \overline{v(\zeta)} d\sigma_{q-1}(\zeta)$$

and corresponding norm  $\|u\| = [(u, u)]^{1/2}$ . Here  $d\sigma_{q-1}$  represents the usual Lebesgue-measure element and  $\sigma_{q-1} = \int_{S^{q-1}} 1 \cdot d\sigma_{q-1}(\zeta)$  is the total measure of  $S^{q-1}$ .

\* Permanent address: Department of Mathematics, Xi'an Jiaotong University, Xi'an, China.

## 2. Identifications of Weighted Hilbert Spaces of Harmonic Functions on $\mathbb{R}^q$ with Domains of Positive Self-adjoint Operators in $L^2(S^{q-1})$

Let  $\mu : (0, \infty) \rightarrow (0, \infty)$  be a Lebesgue measurable function which is nonnegative almost everywhere. It is easy to see that

$$HA^q(\mu) = \{u \mid u(x) \text{ harmonic on } \mathbb{R}^q \text{ and } \|u\|_\mu = \left[ \int_{\mathbb{R}^q} |u(x)|^2 \mu(|x|) dx \right]^{1/2} < \infty\} \quad (1)$$

is a pre-Hilbert space with corresponding inner product

$$(u, v)_\mu = \int_{\mathbb{R}^q} u(x) \overline{v(x)} \mu(|x|) dx. \quad (2)$$

On the other hand let  $\{\lambda_m \mid m \in \mathbb{N}_0\}$  be a sequence of nonnegative numbers. Recall that we have the standard identity decomposition  $L^2(S^{q-1}) = \bigoplus_{m=0}^{\infty} \mathbf{H}_m^q$  (cf. e.g., [7]). So, associated with the given sequence  $\{\lambda_m \mid m \in \mathbb{N}_0\}$  a positive self-adjoint operator  $\Lambda$  in  $L^2(S^{q-1})$  is well defined as follows:

$$\begin{aligned} u &= \sum_{m=0}^{\infty} P_m u \\ \Lambda u &= \sum_{m=0}^{\infty} \lambda_m P_m u \end{aligned} \quad (4)$$

and

$$D(\Lambda) = \{u \in L^2(S^{q-1}) \mid \sum_{m=0}^{\infty} \lambda_m^2 \|P_m u\|^2 < \infty\} \quad (5)$$

where  $P_m$  is the projection of  $L^2(S^{q-1})$  onto  $\mathbf{H}_m^q$ . The domain  $D(\Lambda)$ , equipped with the graph inner product

$$(u, v)_\Lambda = \sum_{m=0}^{\infty} \lambda_m^2 (P_m u, P_m v) \quad (6)$$

and corresponding norm

$$\|u\|_\Lambda = \left( \sum_{m=0}^{\infty} \lambda_m^2 \|P_m u\|^2 \right)^{1/2} \quad (7)$$

is a Hilbert space.

Our main aim in this section is to identify a space  $HA^q(\mu)$  with a space  $D(\Lambda)$  for an admissible

pair  $(\mu, \{\lambda_m\})$  of weight function and nonnegative sequence. In order to formulate and prove the result we need an auxiliary lemma.

**Lemma 1.** (cf. [2] and [3] Theorem IV.1.10)

(i)  $u(\zeta) \in L^2(S^{q-1})$  can be extended to a harmonic function  $u(x)$  on  $B^q(R)$  ( $R > 1$ ) iff

$$\sum_{m=0}^{\infty} r^{2m} \|P_m u\|^2 < \infty \text{ for any } r \in (0, R). \quad (8)$$

(ii) If  $u(x)$  is a harmonic function on  $B^q(R)$  ( $R > 1$ ) then

$$\sum_{m=0}^{\infty} r^m (P_m u)(\zeta) = \sum_{m=0}^{\infty} (P_m u)(x) = u(x) \quad (9)$$

where the series converges uniformly on each ball  $B^q(r)$  with  $r \in (0, R)$ . □

Now we can give

**Theorem 2.** Assume that a weight function  $\mu : (0, \infty) \rightarrow (0, \infty)$  and a sequence  $\{\lambda_m \mid m \in \mathbb{N}_0\}$  satisfy the conditions below

- i)  $\mu_m \equiv \left( \int_0^{\infty} r^{2m+q-1} \mu(r) dr \right)^{1/2} < \infty, \forall m \in \mathbb{N}_0$
- ii)  $R^m / \lambda_m = O(1), \forall R > 0$
- iii)  $\{\lambda_m\} \sim \{\mu_m\}$ , i.e.  $0 < \liminf \frac{\lambda_m}{\mu_m} \leq \limsup \frac{\lambda_m}{\mu_m} < \infty$ .

Then the space  $HA^q(\mu)$  is isomorphic to the space  $D(\Lambda)$  as normed spaces; the isomorphism is exactly the restriction-extension correspondence as in Lemma 1 above. Moreover, if, instead of iii) above,  $\lambda_m = \mu_m$  for all  $m \in \mathbb{N}_0$ , then the isomorphism is actually an isometry.

*Proof.* Let  $u(\zeta) \in D(\Lambda)$ . Because of condition ii) Lemma 1 ensures that  $u(\zeta)$  extends uniformly to a harmonic function  $u(x)$  on  $\mathbb{R}^q$ , namely

$$u(x) = u(r \zeta) = \sum_{m=0}^{\infty} r^m (P_m u)(\zeta). \quad (10)$$

It is then clear that

$$\int_{S^{q-1}} |u(r \zeta)|^2 d\sigma_{q-1}(\zeta) = \sum_{m=0}^{\infty} r^{2m} \|P_m u\|^2. \quad (11)$$

Therefore

$$\begin{aligned} \|u\|_{\mu}^2 &= \int_{\mathbb{R}^q} \mu(|x|) |u(x)|^2 dx \\ &= \int_0^{\infty} r^{q-1} \mu(r) dr \int_{S^{q-1}} |u(r \zeta)|^2 d\sigma_{q-1}(\zeta) \\ &= \int_0^{\infty} r^{q-1} \mu(r) dr \sum_{m=0}^{\infty} r^{2m} \|P_m u\|^2 \\ &= \sum_{m=0}^{\infty} \mu_m^2 \|P_m u\|^2 \end{aligned} \quad (12)$$

which implies that  $u(x) \in HA^q(\mu)$ .

Conversely, if  $u(x) \in HA^q(\mu)$ , then Lemma 1 (ii) implies that equality (10) is valid, so are (11) and (12).

Thus, by condition iii) and equality (12) we conclude that the spaces  $H^q(\mu)$  and  $D(\Lambda)$  are isomorphic as normed spaces via the natural restriction-extension correspondence. In particular, if  $\lambda_m = \mu_m$  for all  $m \in \mathbb{N}_0$ , then they are isometric.  $\square$

**Corollary 3.** Under the conditions in Theorem 2 the space  $HA^q(\mu)$  is complete. So it is a Hilbert space.  $\square$

**Corollary 4.** Under the conditions in Theorem 2 we have  $HA^q(\mu) = \bigoplus_{m=0}^{\infty} H_m^q$ . Moreover, if  $\{e_{m,j}^q(\zeta) \mid 1 \leq j \leq N(q,m)\}$  is an orthonormal basis in  $H_m^q$ , then  $\{\frac{1}{\mu_m} e_{m,j}^q(x) \mid 1 \leq j \leq N(q,m)\}$  is an orthonormal basis in  $H_m^q$  considered a subspace of  $HA^q(\mu)$ .

*Proof.* For  $p_m \in H_m^q$  and  $q_n \in H_n^q$  we have

$$(p_m, q_n)_{\mu} = \int_{\mathbb{R}^q} \mu(|x|) p_m(x) \overline{q_n(x)} dx = \left( \int_0^{\infty} r^{m+n+q-1} \mu(r) dr \right) (p_m, q_n). \quad (13)$$

Hence  $H_m^q \perp H_n^q$  in  $HA^q(\mu)$  if  $m \neq n$ , for  $H_m^q \perp H_n^q$ . Equality (13) implies in particular that

$$\|p_m\|_{\mu}^2 = \mu_m^2 \|p_m\|^2. \quad (14)$$

So equality (12) becomes

$$\|u\|_{\mu}^2 = \sum_{m=0}^{\infty} \|P_m u\|_{\mu}^2. \quad (15)$$

Thus  $HA^q(\mu) = \bigoplus_{m=0}^{\infty} HA_m^q$ . The last assertion in the present corollary follows from (13) and (14) directly.  $\square$

**Corollary 5.** Under the conditions in Theorem 2 the Hilbert space  $HA^q(\mu)$  has a reproducing kernel  $K_{\mu}^q(x, y)$ . Explicitly  $K_{\mu}^q(x, y) = \sum_{m=0}^{\infty} K_{\mu, m}^q(x, y)$ , where, for each  $m$ ,  $K_{\mu, m}^q(x, y)$  is the reproducing kernel of the subspace  $H_m^q$ , namely

$$K_{m, k}^q(x, y) = \frac{N(q, m)}{\sigma_{q-1} \mu_m^2} |x|^m |y|^m P_m^q \left[ \frac{x}{|x|} \cdot \frac{y}{|y|} \right]. \quad (16)$$

Here  $P_m^q$  are the Legendre polynomials (cf. [6] and [7]).

*Proof.* Of the basic theory of reproducing kernel we refer to [1], [8] and [5].

Let  $u \in HA^q(\mu)$ . Then  $u(x) = \sum_{m=0}^{\infty} r^m (P_m u)(\zeta)$ . In view of the estimate  $N(q, m) \leq K_q m^{q-2}$  (cf. [Mü]) and condition ii) in Theorem 2 we have

$$\begin{aligned} |u(x)| &\leq \sum_{m=0}^{\infty} r^m |(P_m u)(\zeta)| \\ &\leq \sum_{m=0}^{\infty} r^m \left[ \frac{N(q, m)}{\sigma_{q-1}} \right]^{\frac{1}{2}} \|P_m u\| \\ &\leq \left[ \sum_{m=0}^{\infty} \frac{N(q, m) r^{2m}}{\sigma_{q-1} \lambda_m^2} \right]^{\frac{1}{2}} \left( \sum_{m=0}^{\infty} \lambda_m^2 \|P_m u\|^2 \right)^{\frac{1}{2}} \\ &\leq C(r) \|u\|_{\mu} \end{aligned} \quad (17)$$

where  $C(r)$  is a constant depending only on  $q$ ,  $\mu$  and  $\{\lambda_m\}$ . The above inequality shows that for each fixed  $x \in \mathbb{R}^q$ , the functional  $u \mapsto u(x)$  is continuous on  $HA^q(\mu)$ . Therefore the Hilbert space  $HA^q(\mu)$  possesses a reproducing kernel  $K_{\mu}^q$ .

Let  $\{e_{m,j}^q \mid 1 \leq j \leq N(q,m)\}$  be an orthonormal basis in  $H_m^q$ . Corollary 4 above says that  $\{\frac{1}{\mu_m} e_{m,j}^q(x) \mid 1 \leq j \leq N(q,m)\}$  is an orthonormal basis in  $H_m^q$ . So its reproducing kernel  $K_{\mu,m}^q$  is given by

$$\begin{aligned} K_{\mu,m}^q(x,y) &= \sum_{j=1}^{N(q,m)} \frac{1}{\mu_m} e_{m,j}^q(x) \cdot \overline{\frac{1}{\mu_m} e_{m,j}^q(y)} \\ &= \frac{|x|^m |y|^m}{\mu_m^2} \sum_{j=1}^{N(q,m)} e_{m,j}^q \left[ \frac{x}{|x|} \right] \overline{e_{m,j}^q \left[ \frac{y}{|y|} \right]} \\ &= \frac{N(q,m)}{\sigma_{q-1} \mu_m^2} |x|^m |y|^2 P_m^q \left[ \frac{x}{|x|} \cdot \frac{y}{|y|} \right]. \end{aligned} \quad (18)$$

Since  $HA^q(\mu) = \bigoplus_{m=0}^{\infty} H_m^q$ , the reproducing kernel  $K_{\mu}^q(x,y)$  for  $HA^q(\mu)$  equals  $\sum_{m=0}^{\infty} K_{\mu,m}^q(x,y)$ .  $\square$

**Corollary 6.**

(i) For  $u \in H_m^q$  holds the estimate

$$|u(x)| \leq \left[ \frac{N(q,m)}{\sigma_{q-1}} \right]^{\frac{1}{2}} \frac{|x|^m}{\mu_m} \|u\|_{\mu}. \quad (19)$$

(ii) For  $u \in HA^q(\mu)$  holds the estimate

$$|u(x)| \leq \left[ \sum_{m=0}^{\infty} \frac{N(q,m)}{\sigma_{q-1}} \frac{|x|^{2m}}{\mu_m^2} \right]^{\frac{1}{2}} \|u\|_{\mu}. \quad (20)$$

(iii) For  $u \in HA^q(\mu)$  and  $m \in \mathbb{N}_0$

$$(P_m u)(x) = \frac{N(q,m) |x|^m}{\sigma_{q-1} \mu_m^2} \int_{\mathbb{R}^q} |y|^m P_m^q \left[ \frac{x}{|x|} \cdot \frac{y}{|y|} \right] u(y) dy. \quad (21)$$

*Proof.* Assertion (i) and (ii) follow directly from Corollary 5. On multiplying both sides of (18) by  $u(y)$  and integrating over  $\mathbb{R}^q$  we obtain equality (21) immediately.  $\square$



### 3. Conditions for Differential Operators to be Continuous or Compact in $HA^q(\mu)$

In this section Theorem 2 in the previous section is applied to give conditions such that differentiation operators are continuous or compact in the Hilbert space  $HA^q(\mu)$ .

**Theorem 7.** Suppose the conditions in Theorem 2 are satisfied.

- (i) If  $\lambda_m^2 / \lambda_{m+1}^2 = O(m^{-q})$ , then the differentiation operator  $\partial_k (1 \leq k \leq q)$  is continuous on  $HA^q(\mu)$ .
- (ii) If  $\lambda_m^2 / \lambda_{m+1}^2 = o(m^{-q})$ , then the differentiation operator  $\partial_k (1 \leq k \leq q)$  is compact on  $HA^q(\mu)$ .  $\square$

In the proof of the theorem we need an estimate of spherical harmonics given in [4]:

**Lemma 8.** For  $h \in H_m^q$  holds the estimate

$$|(\partial_k h)(\zeta)| \leq m \left[ \frac{N(q, m)}{\sigma_{q-1}} \right]^{1/2} \|h\|.$$

$\square$

*Proof of Theorem 7.* Let  $u(x)$  be a harmonic function on  $\mathbb{R}^q$ . The  $u(x) = \sum_{m=0}^{\infty} (P_m u)(x)$  where the convergence is uniform on compacta of  $\mathbb{R}^q$  and  $\sum_{m=0}^{\infty} R^{2m} \|P_m u\|^2 < \infty$  for any  $R > 0$  (Lemma 1). By Lemma 8 we have

$$\begin{aligned} |(\partial_k [(P_m u)(x)])| &\leq r^{m-1} |(\partial_k (P_m u))(\zeta)| \\ &\leq r^{m-1} m \left[ \frac{N(q, m)}{\sigma_{q-1}} \right]^{1/2} \|P_m u\| \\ &\leq \left[ \frac{K_q}{\sigma_{q-1}} \right]^{1/2} R^{-1} m^{q/2} \left[ \frac{r}{R} \right]^{m-1} \cdot R^m \|P_m u\| \quad (0 < r < R). \end{aligned} \quad (22)$$

Therefore the series  $\sum_{m=0}^{\infty} \partial_k [(P_m u)(x)]$  converges uniformly on compacta of  $\mathbb{R}^q$ . Hence

$$(\partial_k u)(x) = \sum_{m=0}^{\infty} \partial_k [(P_m u)(x)]. \quad (23)$$

In particular, for  $u \in HA^q(\mu)$  we have

$$\begin{aligned}
 \|\partial_k u\|_\Lambda^2 &= \sum_{m=0}^{\infty} \lambda_m^2 \|P_m(\partial_k u)\|^2 \\
 &= \sum_{m=0}^{\infty} \lambda_m^2 \|\partial_k(P_{m+1} u)\|^2 \\
 &\leq \sum_{m=0}^{\infty} \lambda_m^2 \int_{S^{q-1}} |[\partial_k(P_{m+1} u)](\zeta)|^2 d\sigma_{q-1}(\zeta) \\
 &\leq \sum_{m=0}^{\infty} \lambda_m^2 (m+1)^2 \frac{N(q,m)}{\sigma_{q-1}} \|P_{m+1} u\|^2 \sigma_{q-1} \quad (\text{Lemma 8}) \\
 &\leq \sum_{m=0}^{\infty} K_q (m+1)^q \frac{\lambda_m^2}{\lambda_{m+1}^2} \cdot \lambda_{m+1}^2 \|P_{m+1} u\|^2. \tag{24}
 \end{aligned}$$

(i) If  $\frac{\lambda_m^2}{\lambda_{m+1}^2} = O(m^{-q})$ , then the above inequality (24) shows that  $(\partial_k u)(\zeta) \in D(\Lambda)$ , and  $\|\partial_k u\|_\Lambda \leq c\|u\|_\Lambda$  for some constant  $c$  independent on  $u$ . By Theorem 2 we then conclude that  $(\partial_k u)(x) \in HA^q(\mu)$  and  $\|\partial_k u\|_\mu \leq c'\|u\|_\mu$  with  $c'$  another constant. Thus the differentiation operator  $\partial_k$  is continuous on  $HA^q(\mu)$ .

(ii) If  $\frac{\lambda_m^2}{\lambda_{m+1}^2} = o(m^{-q})$ , then the inequality (24) implies that for a given bounded set  $\mathbf{B}$  in  $D(\Lambda)$  the set  $\{\partial_k u \mid u \in \mathbf{B}\}$  is bounded in  $D(\Lambda)$  and for any  $\varepsilon > 0$  there exists  $M$  such that  $\sum_{m \geq M} \lambda_m^2 \|P_m \partial_k u\|^2 < \varepsilon$  for all  $u \in \mathbf{B}$ . Thus  $\{\partial_k u \mid u \in \mathbf{B}\}$  is a relatively compact set in  $D(\Lambda)$ .

Therefore  $\partial_k$  is a compact operator on  $D(\Lambda)$ , so is it on  $HA^q(\mu)$  by Theorem 2.  $\square$

#### 4. Examples

Let  $\mu(r) = e^{-ar^b}$  ( $a, b > 0$ ). Then the differentiation operators  $\partial_k$  ( $1 \leq k \leq q$ ) are compact on  $HA^q(\mu)$  for  $b < \frac{2}{q}$  and continuous on  $HA^q(\mu)$  for  $b = \frac{2}{q}$ . This assertion follows from Theorem 2 and Theorem 7 above and the following two lemmas.

**Lemma 9.**

$$u_m^2 = \int_0^\infty e^{-ar^b} r^{2m+q-1} dr = (2\pi)^{1/2} b^{-1/2} (eab)^{-\frac{2m+q}{b}} (2m+q)^{\frac{2m+q}{b}-\frac{1}{2}}. \quad (25)$$

*Proof.* Substituting  $s$  for  $ar^b$  we have

$$\begin{aligned} u_m^2 &= \int_0^\infty e^{-ar^b} r^{2m+q-1} dr \\ &= \int_0^\infty e^{-s} a^{-\frac{2m+q}{b}} b^{-1} s^{\frac{2m+q}{b}-1} ds \\ &= a^{-\frac{2m+q}{b}} b^{-1} \Gamma\left[\frac{2m+q}{b}\right]. \end{aligned}$$

Then an invocation to Stirling's formula

$$\Gamma(\theta) = (2\pi)^{1/2} \theta^{\theta-1/2} e^{-\theta} \quad (\theta \rightarrow \infty)$$

yields relation (25). □

**Lemma 10.** Set  $\lambda_m^2 = \left[ \frac{2m}{eab} \right] \frac{2}{b^m} m^{\frac{q}{b}-\frac{1}{2}}$  ( $m \in \mathbb{N}_0$ ). Then  $\frac{R^{2m}}{\lambda_m^2} = O(1)$  for all  $R > 0$  and

$$\{\lambda_m\} \sim \{\mu_m\}.$$

Moreover

$$\frac{\lambda_m^2}{\lambda_{m+1}^2} \sim m^{-\frac{2}{b}}.$$

□

Further examples of Hilbert spaces of type  $HA^q(\mu)$  wherein differentiation operators are continuous or compact are given in [3].

## References

- [1] Aronszajn, N., Theory of reproducing kernels, Trans. AMS, **68**, 337-404 (1950).
- [2] Graaf, J. de, Two spaces of generalized functions based on harmonic polynomials, in C. Brezinski, et al (ed.), Polynômes orthogonaux et Applications, Proc. Bar-le-Duc 1984, Lecture Notes in Mathematics 1171, Springer, Berlin et al., 1984.
- [3] Liu. G.-Z., Scales of Banach Spaces and Evolution Equations, Ph. D Thesis, Eindhoven University of Technology, Eindhoven, Netherlands, June 1989.
- [4] Liu, G.-Z., Martens F.J.L., Improvement on an Estimate of Spherical Harmonics, preprint, Eindhoven, Netherlands, 1989.
- [5] Martens, F.J.L., Spaces of Analytic Functions on Inductive/Projective Limits of Hilbert Spaces, Ph. D Thesis, Eindhoven University of Technology, Eindhoven, November 1988.
- [6] Müller, C., Spherical Harmonics, Lecture Notes in Mathematics 17, Springer, Berlin et al., 1966.
- [7] Stein E.M., Weiss G., Introduction to Fourier Analysis on Euclidean Spaces, Princeton University Press, Princeton, New Jersey, 1971.
- [8] Yosida, K., Functional Analysis, Grundlehren der Mathematischen Wissenschaften 123, Sixth Edition, Springer, 1980.

## Acknowledgement:

The author would like to express his gratitude to Prof. De Graaf for his inspiring advices.

Key Words: Differential operators; spherical harmonics; Hilbert spaces of harmonic functions; reproducing kernels; Stirling's formula.

1980 AMS Classification: 46E20, 47F05, 47B38, 31B05, 33A45, 33A15