

Hitting times for multiplicative growth-collapse processes

Citation for published version (APA):

Löpker, A. H., & Leeuwaarden, van, J. S. H. (2007). *Hitting times for multiplicative growth-collapse processes*. (Report Eurandom; Vol. 2007049). Eurandom.

Document status and date:

Published: 01/01/2007

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

HITTING TIMES FOR MULTIPLICATIVE GROWTH-COLLAPSE PROCESSES

Andreas H. Löpker[†]Johan S. H. van Leeuwen[‡]

Abstract: We consider a stochastic process $(X_t)_{t \geq 0}$ that grows linearly in time and experiences collapses at times governed by a Poisson process with rate λ . The collapses are modeled by multiplying the process level by a random variable supported on $[0, 1)$. For the hitting time defined as $\tau_y = \inf\{t > 0 | X_t = y\}$ we derive power series for the Laplace transform and all moments. We further discuss the asymptotic behavior of the mean of τ_y as y tends to infinity.

1. INTRODUCTION

We investigate a growth-collapse process $(X_t)_{t \geq 0}$ with deterministic growth and a multiplicative collapse structure. The process increases linearly with rate one and at the time T_i of the i th collapse, the process jumps down to $Q_i \cdot X_{T_i}$, where Q, Q_1, Q_2, \dots are i.i.d. random variables with distribution function F_Q supported on $[0, 1)$.

The collapse times $(T_i)_{i \in \mathbb{N}}$ are governed by a Poisson process with rate λ . We define the hitting time of level y by

$$\tau_y = \inf\{t > 0 | X_t = y\}. \quad (1)$$

We shall establish a formula for the Laplace transform of τ_y , as well as for its moments, and the asymptotic behavior of its mean.

Growth-collapse processes are real-valued processes that grow between random collapse times, at which they jump down according to some distribution depending on their current level. This evolutionary pattern is encountered in a large variety of physical phenomena, see [16], like build-up of friction, earthquakes, avalanches, neuron firing, and shot noise, as well as in other fields like insurance mathematics [29], queueing theory [7] and mathematical finance [9]. Some properties of Markovian growth collapse models have been studied in [12]. The case where Q is a constant (usually $Q = 1/2$) is used as a model for the Transmission Control Protocol (TCP), the dominant protocol for data transfer over the internet, cf. Section 7 and [4, 5, 8, 15, 19, 27]. Picking items on a circle [23, 24] and DNA replication [22] are further applications of this case. Moreover, as indicated in [26] (see also [19]), the process $(X_t)_{t \geq 0}$ is equivalent to an exponential functional ([9, 13]) of a Lévy process.

In the literature, emphasis lies on analyzing the stationary behavior of the growth-collapse processes. The results we present for the hitting times are some of the few results obtained on the transient behavior of growth-collapse processes.

[†]EURANDOM, P.O. Box 513 - 5600 MB Eindhoven, The Netherlands.
Email address: lopker@eurandom.tue.nl

[‡]Eindhoven University of Technology and EURANDOM, P.O. Box 513 - 5600 MB Eindhoven, The Netherlands. Email address: j.s.h.v.leeuwen@tue.nl

Another noteworthy contribution on transient behavior is [15] in which results are obtained for the mean hitting time of a growth-collapse process with Q a constant and the Poisson process having rate $\lambda \cdot X_t$ (level-dependent).

To obtain our results, we invoke techniques from the field of piecewise deterministic Markov processes (see [14]). We consider the infinitesimal generator of the strong Markov process $(X_t)_{t \geq 0}$ and apply the well known Dynkin formula. In Theorem 2 we find for τ_y , starting from level $x < y$, that its mean can be expressed as

$$E_x \tau_y = \sum_{n=1}^{\infty} \frac{y^n - x^n}{n!} \prod_{i=1}^{n-1} \theta_i$$

and its Laplace transform as

$$E_x e^{-s\tau_y} = \frac{1 + s \sum_{n=1}^{\infty} \frac{x^n}{n!} \prod_{i=1}^{n-1} (s + \theta_i)}{1 + s \sum_{n=1}^{\infty} \frac{y^n}{n!} \prod_{i=1}^{n-1} (s + \theta_i)},$$

where $\theta_i = \lambda(1 - E(Q^i))$.

We also investigate the asymptotic behavior of $E_x \tau_y$ as $y \rightarrow \infty$. Clearly the asymptotic behavior will strongly depend on the behavior of $F_Q(x)$ as $x \rightarrow 1$. It turns out that the asymptotic relations for $E_x \tau_y$ all have as their leading behavior $e^{\lambda y}$ as $y \rightarrow \infty$, and so one might argue that the mean hitting times are almost exponential. In the context of Markov chains on finite state spaces there is a vast literature on almost exponential hitting times (see [1] for an overview). As advocated by Aldous [3], hitting times of rarely-visited sets are approximately exponentially distributed if stationarity is reached rapidly. In [26] the authors obtained expressions for the transient moments of $(X_t)_{t \geq 0}$. These moments were shown to converge to their stationary counterparts exponentially fast.

A hitting time is a valuable characteristic, as it indicates the ability of the process to recover after a collapse. We therefore introduce the closely related notion of *recovery time* σ_x , defined as the time to reach level x conditioned on the fact that a collapse took place when the process was exactly in x . As for the hitting time, we obtain for the recovery time explicit expressions for the Laplace transform and its mean. Moreover, we show that the recovery time and hitting time have the same asymptotic behavior.

We have structured the paper as follows. A formal introduction to the model is presented in Section 2, including the infinitesimal generator, Dynkin's formula and a specific class of test functions. In Section 3 we derive explicit expressions for the means and Laplace transforms of the hitting- and recovery times. Section 4 is concerned with the asymptotic behavior of the mean hitting time. The main theorem, covering various modes of asymptotics, is proved in Section 5. We conclude with two special choices of Q in Sections 6 and 7. In Section 6 we treat the case where Q follows a Beta distribution. Several characteristics are reformulated in terms of hypergeometric functions and exponential integrals. In Section 7 we consider the case where Q is some constant in $[0, 1)$, which gives rise to q -calculus.

In what follows let P_x denote as usual the conditional probability given that the process starts at $X_0 = x$. Moreover, let $P_{(x)}$ be the probability given that the process starts with a jump from x and let E_x and $E_{(x)}$ be the respective expectations. Note that $E_{(x)}$ is obtained from E_x by conditioning on the first jump,

$$E_{(x)}(\cdot) = \int_0^1 E_{xy}(\cdot) dF_Q(y). \quad (2)$$

For a probability distribution function F , let $\bar{F} = 1 - F$. For the asymptotic behavior of functions we sometimes write $f(x) \asymp g(x)$ if there is a constant $C \in (0, \infty)$ such that $\lim f(x)/g(x) = C$. In particular, if $C = 1$ we use the usual notation $f(x) \sim g(x)$ for asymptotic equivalence. The infimum over the empty set is defined to be infinite, empty sums are zero, empty products are one.

We already introduced the coefficients $\theta_a = \lambda(1 - E(Q^a))$. We further introduce $\pi_0(s) = 1$ and the product

$$\pi_k(s) = \prod_{i=1}^k (\theta_i + s).$$

2. THE MODEL

We thus consider the process $(X_t)_{t \geq 0}$, a Markov process with linear deterministic increase and multiplicative jumps. The process jumps down at times $(T_i)_{i \in \mathbb{N}}$ that are governed by a Poisson process $(N_t)_{t \geq 0}$ with intensity λ . At T_i the process jumps from X_{T_i-} to $X_{T_i} = Q_i \cdot X_{T_i-}$ where Q, Q_1, Q_2, \dots are i.i.d. random variables with distribution function F_Q supported on $[0, 1)$. We exclude the case where $P(Q = 1) > 0$, although it could be included by changing the jump intensity from λ to $P(Q < 1) \cdot \lambda$.

The process is a special case of a piecewise deterministic Markov process introduced by Davis [14]. The state space \mathcal{S} consists of all non-negative real numbers. The extended generator of the strong Markov process $(X_t)_{t \geq 0}$ is given by

$$\mathcal{A}f(x) = f'(x) - \lambda f(x) + \lambda \int_0^1 f(xy) dF_Q(y), \quad x \in \mathcal{S} = [0, \infty).$$

The domain of \mathcal{A} contains absolutely continuous functions $f : [0, \infty) \rightarrow \mathbb{R}$ that are either locally bounded or for which $f(xy) \leq f(x)f(y)$ holds (see [26]). The defining property of the extended generator is that for all functions f in the domain of \mathcal{A} the stochastic process $f(X_t) - \int_0^t \mathcal{A}f(X_s) ds$ is a martingale.

Our central observation is that for all a for which $E(Q^a) < \infty$ the application of the generator to power functions leads to a sum of two powers. That is,

$$\mathcal{A}x^a = ax^{a-1} - x^a \theta_a. \quad (3)$$

It has been shown in [26] that (3) can be used to obtain a formula for the transient moments of the process. In this paper we will demonstrate how (3) can be utilized to find expressions for the mean and the Laplace transform of the hitting- and recovery times. The preservation of powers property in (3) is crucial in proving Theorem 1, which gives a solution to the generator equation $\mathcal{A}f(x) = sf(x) + z(x)$, where z is some arbitrary function.

3. HITTING- AND RECOVERY TIMES

We have defined the *hitting time* of the level $y \in \mathcal{S}$ as $\tau_y = \inf\{t > 0 | X_t = y\}$. We have furthermore defined the *recovery time* σ_x as the time the process needs to reach level x , conditionally on the fact that a jump takes place when the process is in level x . It has been shown in [12] and [26] that $E_x \tau_y < \infty$ for all $x, y \in \mathcal{S}$. Obviously, $E_{(x)} \sigma_x \leq E_0 \tau_x < \infty$ for all $x \in \mathcal{S}$.

To prepare the ground for our main result we provide a solution to the generator equation $\mathcal{A}f(x) = sf(x) + z(x)$. The result is more general than needed, but perhaps of some interest for future study of this model.

Theorem 1. Let $z : \mathbb{R} \rightarrow \mathbb{R}$ be a function analytic in a neighborhood zero. A solution of $\mathcal{A}f(x) = sf(x) + z(x)$ with $f(0) = a_0 \in \mathbb{R}$ is given by

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(sa_0 + \sum_{k=0}^{n-1} \frac{z^{(k)}(0)}{\pi_k(s)} \right) \pi_{n-1}(s) \frac{x^n}{n!}.$$

Proof. It is straightforward to show that for a function $h(x) = \sum_{n=0}^{\infty} a_n \pi_{n-1}(s) \frac{x^n}{n!}$ we get

$$\mathcal{A}h(x) = (1-s)h(0) + sh(x) + \sum_{n=1}^{\infty} \Delta a_n \pi_{n-1}(s) \frac{x^{n-1}}{(n-1)!},$$

where $\Delta a_n = a_n - a_{n-1}$. Thus the equation $\mathcal{A}f(x) = sf(x) + z(x)$ can be written as

$$(1-s)a_0 + \sum_{n=1}^{\infty} \Delta a_n \pi_{n-1}(s) \frac{x^{n-1}}{(n-1)!} = \sum_{n=1}^{\infty} z^{(n-1)}(0) \frac{x^{n-1}}{(n-1)!}.$$

It follows from comparison of the coefficients that $z(0) = (1-s)a_0 + (a_1 - a_0) = a_1 - sa_0$ and $\Delta a_n = z^{(n-1)}(0)$, and finally that $a_n = sa_0 + \sum_{k=1}^n \frac{z^{(k-1)}(0)}{\pi_{k-1}(s)}$ for $n \geq 1$. \square

We now present our main result. Let the function $R_s(x)$ be given by

$$R_s(x) = \sum_{n=1}^{\infty} \pi_{n-1}(s) \frac{x^n}{n!}. \quad (4)$$

Theorem 2. For $x \leq y$ the means and Laplace transforms of τ_y and σ_x are given by

$$E_x \tau_y = R_0(y) - R_0(x), \quad (5)$$

$$E_x e^{-s\tau_y} = \frac{1 + sR_s(x)}{1 + sR_s(y)}, \quad (6)$$

$$E_{(x)} \sigma_x = \frac{1}{\lambda} (R'_0(x) - 1), \quad (7)$$

$$E_{(x)} e^{-s\sigma_x} = 1 + \frac{1}{\lambda} \left(s - \frac{R'_s(x)}{1 + sR_s(x)} \right). \quad (8)$$

Proof. We utilize the fact that, if f is in the domain of \mathcal{A} ,

$$f(X_t) - \int_0^t \mathcal{A}f(X_s) ds \quad (9)$$

and

$$f(X_t) \exp \left(- \int_0^t \mathcal{A}f(X_s)/f(X_s) ds \right) \quad (10)$$

are martingales (see e.g. [15, 17]). It follows from Theorem 1 that $R_s(x)$ is the solution of the equation $\mathcal{A}f(x) = sf(x) + 1$ with $f(0) = 0$. In particular, $\mathcal{A}R_0(x) = 1$, so that $R_0(X_t) - t$ is a martingale and optional stopping, if allowed, yields $E_x \tau_y = R_0(y) - R_0(x)$. On $\{t < \tau_y\}$ we have that $|R_0(X_t) - t| \leq R_0(y) + t = O(t)$ as $t \rightarrow \infty$. Since $E_x \tau_y < \infty$ it follows that

$$E_x(R_0(X_t) - t; t < \tau_y) = E_x(R_0(X_t) - t | t < \tau_y) o(1/t) \rightarrow 0$$

as $t \rightarrow \infty$ and thus the optional stopping theorem can be applied (see [17] for this criterion) and (5) follows.

Since $\mathcal{A}(1 + sR_s(x)) = s(1 + sR_s(x))$ it follows that $e^{-st}(1 + sR_s(X_t))$ is a martingale and optional stopping yields (6). Because $R_s(X_t)$ is bounded for $t < \tau_y$, optional stopping is allowed here (see also [21]).

We can find an expression for the mean of σ_x in terms of the function R_0 by applying the conditioning formula (2), i.e.,

$$E_{(x)}\sigma_x = \int_0^1 E_{xy}\tau_x dF_Q(y) = \int_0^1 (R_0(x) - R_0(xy)) dF_Q(y).$$

Since $\mathcal{A}R_0(x) = 1 = R'_0(x) - \lambda \int_0^1 (R_0(x) - R_0(xy)) dF_Q(y)$, it follows that

$$E_{(x)}\sigma_x = \frac{R'_0(x) - \mathcal{A}R_0(x)}{\lambda} = \frac{R'_0(x) - 1}{\lambda}.$$

A similar method works for the Laplace transform. Using (2) and (6) yields

$$E_{(x)}e^{-s\sigma_x} = E_{(x)}e^{-s\tau_x} = \int_0^1 \frac{1 + sR_s(xy)}{1 + sR_s(x)} dF_Q(y).$$

Since $\mathcal{A}(1 + sR_s(x)) = s(1 + sR_s(x))$ it follows that

$$s = \frac{\mathcal{A}(1 + sR_s(x))}{1 + sR_s(x)} = \frac{R'_s(x)}{1 + sR_s(x)} - \lambda + \lambda \int_0^1 \frac{1 + sR_s(xy)}{1 + sR_s(x)} dF_Q(y).$$

Consequently,

$$E_{(x)}e^{-s\sigma_x} = 1 + \frac{\mathcal{A}(1 + sR_s(x)) - R'_s(x)}{\lambda(1 + sR_s(x))} = 1 + \frac{s}{\lambda} - \frac{R'_s(x)}{\lambda(1 + sR_s(x))}.$$

This completes the proof. \square

To derive formulas for the higher moments we define

$$\pi_k^{(m)}(s) = \sum_{\substack{K \subseteq \{1, \dots, k\} \\ |K|=k-m}} \prod_{i \in K} (\theta_i + s),$$

and $\kappa_n(x) = \sum_{k=n}^{\infty} \frac{x^k}{k!} \pi_{k-1}^{(n-1)}(0)$. Note that $\pi_k(s) = \pi_k^{(0)}(s)$ and thus $\kappa_1(x) = R_0(x)$. The next lemma shows that these functions are solutions of the equation $\mathcal{A}^k \kappa_k(x) = 1$, where \mathcal{A}^k denotes the repeated application of the operator \mathcal{A} . Moreover, although we will not use this fact in the sequel, it is shown that the κ_k are the coefficients of the power series expansion of R_s in s .

Lemma 3. $\mathcal{A} \kappa_n(x) = \kappa_{n-1}(x)$ and $R_s(x) = \sum_{n=1}^{\infty} \kappa_n(x) s^{n-1}$.

Proof. We have

$$\begin{aligned} R_s(x) &= \sum_{n=1}^{\infty} \prod_{k=1}^{n-1} (\theta_k + s) \frac{x^n}{n!} = \sum_{n=1}^{\infty} \left(\sum_{k=0}^{n-1} \pi_{n-1}^{(k)}(0) s^k \right) \frac{x^n}{n!} \\ &= \sum_{n=1}^{\infty} s^{n-1} \sum_{k=n}^{\infty} \pi_{k-1}^{(n-1)}(0) \frac{x^k}{k!} = \sum_{n=1}^{\infty} s^{n-1} \kappa_n(x). \end{aligned}$$

That $\mathcal{A} \kappa_n(x) = \kappa_{n-1}(x)$ follows from $\mathcal{A} R_s(x) = 1 + sR_s(x)$. \square

Theorem 4. For $x \leq y$ the higher moments of τ_y and σ_x are given via the recursions

$$E_x \tau_y^n = n! \left(\kappa_n(x) - \kappa_n(y) - \sum_{k=1}^{n-1} (-1)^k \frac{\kappa_k(y)}{(n-k)!} E_x \tau_y^{n-k} \right), \quad (11)$$

$$E_{(x)} \sigma_x^n = (-1)^n n! \left(\frac{\kappa_{n-1}(x) - \kappa'_n(x)}{\lambda} - \sum_{k=1}^{n-1} (-1)^k \frac{\kappa_k(x)}{(n-k)!} E_{(x)} \sigma_x^{n-k} \right). \quad (12)$$

Proof. It is shown in [25] that (11) holds if $\mathcal{A}\kappa_n(x) = \kappa_{n-1}(x)$. Equation (12) is immediate if we use (2). We have

$$\begin{aligned} E_{(x)}\sigma_x^n &= (-1)^n n! \int_0^1 \left(\kappa_n(xu) - \kappa_n(x) - \sum_{k=1}^{n-1} (-1)^k \frac{\kappa_k(x)}{(n-k)!} E_{xu}\tau_x^{n-k} \right) dF_Q(u) \\ &= (-1)^n n! \left(\frac{\kappa_{n-1}(x) - \kappa'_n(x)}{\lambda} - \sum_{k=1}^{n-1} (-1)^k \frac{\kappa_k(x)}{(n-k)!} E_{(x)}\sigma_x^{n-k} \right). \quad \square \end{aligned}$$

4. ASYMPTOTIC BEHAVIOR OF THE HITTING TIME AND RECOVERY TIME

Our first asymptotic result gives an upper bound for the mean of τ_y and σ_y and states that as $y \rightarrow \infty$ both values grow equally fast in the sense of asymptotic equivalence (denoted by \sim). Both statements are not difficult to prove and we will see later that the actual growth may differ considerably from these bounds.

Proposition 5. *We have*

$$\frac{e^{\theta_1 y} - 1}{\theta_1} \leq E_0\tau_y \leq \frac{e^{\lambda y} - 1}{\lambda} \quad (13)$$

and $\frac{1}{\lambda} (e^{\theta_1 y} - 1) \leq E_{(y)}\sigma_y \leq \frac{1}{\lambda} (e^{\lambda y} - 1)$. Moreover, as $y \rightarrow \infty$,

$$E_{(y)}\sigma_y \sim E_x\tau_y. \quad (14)$$

Proof. It is immediate from the definition of R_s and $\theta_1 \leq \theta_n \leq \lambda$ that

$$\frac{e^{\theta_1 y} - 1}{\theta_1} = \sum_{n=1}^{\infty} \left(\prod_{k=1}^{n-1} \lambda(1 - EQ) \right) \frac{y^n}{n!} \leq R_s(y) \leq \sum_{n=1}^{\infty} \lambda^{n-1} \frac{y^n}{n!} = \frac{e^{\lambda y} - 1}{\lambda}.$$

Next, we have $E_{(y)}\sigma_y = \frac{1}{\lambda} (R'_0(y) - 1)$ and

$$e^{\theta_1 y} = \sum_{n=0}^{\infty} \left(\prod_{k=1}^n \theta_1 \right) \frac{y^n}{n!} \leq R'_0(y) \leq \sum_{n=0}^{\infty} \lambda^n \frac{y^n}{n!} = e^{\lambda y}.$$

Finally, with $\pi_n = \pi_n(0)$,

$$R'_0(y) = \sum_{n=0}^{\infty} \pi_n \frac{y^n}{n!} = \theta_1 + \sum_{n=1}^{\infty} \frac{\pi_n}{\pi_{n+1}} \pi_{n+1} \frac{y^n}{n!},$$

so that relation (14) follows from Lemma 8 and the fact that $\pi_{n+1}/\pi_n = \theta_{n+1}$ converges to λ as $n \rightarrow \infty$. \square

The next result shows that there is no smaller coefficient than λ in the linear term in the exponent.

Proposition 6. *For every $\varepsilon > 0$,*

$$e^{(1-\varepsilon)\lambda y} \leq E_0\tau_y \leq \frac{e^{\lambda y} - 1}{\lambda}, \quad (15)$$

for large enough values of y .

Proof. The upper bound in (15) is copied from (13). For an entire function $f(x) = \sum_{n=0}^{\infty} a_n x^n$ for which $e^{c(1-\varepsilon)x^p} \leq f(x) \leq e^{c(1+\varepsilon)x^p}$ holds ultimately for every $\varepsilon > 0$ the coefficients ρ and c are called *order* and *type* of f . One can calculate the order and

type by using the formulas $\rho = -\lim_{n \rightarrow \infty} n \log n / \log |a_n|$ and $c = \lim_{n \rightarrow \infty} \frac{n}{e\rho} |a_n|^{\rho/n}$, respectively (see e.g. [11]). Stirling's formula and $\theta_n^{n-1} \leq \pi_n \leq \theta_1^{n-1}$ yield

$$\begin{aligned} \rho^{-1} &= \lim_{n \rightarrow \infty} \frac{\log n! - \log \pi_{n-1}}{n \log n} = \lim_{n \rightarrow \infty} \left(1 - \frac{\log \pi_{n-1}}{n \log n} \right) = 1. \\ c &= \lim_{n \rightarrow \infty} \frac{n}{e} (\pi_{n-1}/n!)^{1/n} = \lim_{n \rightarrow \infty} |\pi_{n-1}|^{1/n} = \lim_{n \rightarrow \infty} \frac{\sum_{i=1}^{n-1} \log \theta_i}{n} = \lambda. \end{aligned}$$

Thus (15) follows. \square

For $q \in [0, 1]$, let $(q)_\infty = \prod_{k=1}^{\infty} (1 - q^k)$ denote the infinite q -series (see Askey et al. [2]) and define $x_Q = \sup\{x \in [0, 1] \mid F_Q(x) < 1\}$. A measurable function $L : [0, \infty) \rightarrow [0, \infty)$ is said to be *slowly varying* at 0 if $L(cx)/L(x) \rightarrow 1$ as $x \rightarrow 0$ for all $c > 0$ (cf. [10]). The central result of this section is the following theorem, which gives the asymptotic behavior of $E_x \tau_y$ as y tends to infinity in some important cases.

Theorem 7. *Let F be a probability distribution function on $[0, 1]$ with*

$$\bar{F}(x) = (1 - x)^\beta L(1 - x)$$

and $L(x)$ slowly varying as $x \rightarrow 0$. The following asymptotic relations hold as $y \rightarrow \infty$.

1. *If either $x_Q < 1$ or $F_Q(x) \geq F(x)$, $\beta = 1$ and $\int_1^\infty L(1/u)/u \, du < \infty$ then*

$$E_x \tau_y \sim \frac{D}{\lambda} \cdot e^{\lambda y} \quad (16)$$

with $D = \prod_{k=1}^{\infty} (1 - E(Q^k))$. Moreover, $(x_Q)_\infty \leq D \leq (EQ)_\infty$.

2. *If $F_Q(x) \sim F(x)$ as $x \rightarrow 1$ and $\beta \in (\frac{1}{2}, 1]$ then $E_x \tau_y \sim C \cdot \exp(\int_0^y \theta_{\lambda t} \, dt)$ with some $C > 0$. In particular,*
 - (a) *if $\beta = 1$ then*

$$E_x \tau_y \sim \exp \left(\lambda y - \int_1^{\lambda y} \frac{L(1/t)}{t} \, dt \cdot (1 + o(1)) \right). \quad (17)$$

- (b) *if $\beta \in (\frac{1}{2}, 1)$ then*

$$E_x \tau_y \sim \exp \left(\lambda y - (\lambda y)^{1-\beta} L(1/(\lambda y)) \cdot \left(\frac{\Gamma(1+\beta)}{1-\beta} + o(1) \right) \right). \quad (18)$$

Some remarks are in order.

Firstly, for the third assertion in Theorem 7, we rely heavily on a depoissonization lemma of Foss & Korshunov [18] (see Lemma 11). Although this result may be used also for our $\beta \in (0, \frac{1}{2})$ case, the results in [18] are not easy to state in this case. We refer to the original paper for the interested reader. Further depoissonization methods can be found in [20, 28].

Secondly, note the appearance of the $o(1)$ term in (17) and (18). As $y \rightarrow \infty$ the quotients

$$E_x \tau_y / \exp \left(\lambda y - \int_1^{\lambda y} \frac{L(1/t)}{t} \, dt \right)$$

and

$$E_x \tau_y / \exp \left(\lambda y - (\lambda y)^{1-\beta} L(1/(\lambda y)) \cdot \left(\frac{\Gamma(1+\beta)}{1-\beta} \right) \right)$$

may considerably deviate from constants. It seems complicated though to derive more accurate results.

5. PROOF OF THEOREM 7

We need several lemmas. The first lemma is well known from calculus and shows that functions expanded in power series with asymptotically equivalent coefficients are asymptotically equivalent.

Lemma 8. *Let (a_n) and (b_n) be two non-negative real sequences and let $A(x) = \sum_{k=1}^{\infty} a_k x^k$ and $B(x) = \sum_{k=1}^{\infty} b_k x^k$ be convergent for all x . If $a_k \sim c \cdot b_k$ as $k \rightarrow \infty$ then $A(x) \sim c \cdot B(x)$ as $x \rightarrow \infty$. In particular, if $a_k \asymp b_k$ then $A(x) \asymp B(x)$.*

Proof. First note that $A(x)$ and $B(x)$ tend to ∞ as $x \rightarrow \infty$. For all $\varepsilon > 0$ there is an $m \in \mathbb{N}$ such that $b_k(1 - \varepsilon/2) \leq a_k \leq b_k(1 + \varepsilon/2)$. Thus

$$A(x) \leq \sum_{k=1}^{m-1} x^k a_k + \sum_{k=m}^{\infty} x^k b_k (1 + \varepsilon/2)$$

Let $a'_m = \sum_{k=1}^{m-1} x^k a_k$ and $b'_m = \sum_{k=1}^{m-1} x^k b_k$. Then

$$\frac{A(x)}{B(x)} \leq \frac{a'_m}{B(x)} + (1 + \varepsilon/2) \left(1 - \frac{b'_m}{B(x)}\right).$$

Thus for every $\varepsilon > 0$, $A(x)/B(x) \leq (1 + \varepsilon)$ ultimately as $x \rightarrow \infty$. It can be shown similarly that $A(x)/B(x) \geq (1 - \varepsilon)$. \square

The next result gives the asymptotics of the coefficients $\pi_k(s) = \prod_{i=1}^k (\theta_i + s)$ in terms of the moments $E(Q^k)$.

Lemma 9. *Let M be the smallest integer such that $\sum_{j=1}^{\infty} E(Q^j)^M < \infty$. Then*

$$\pi_k(s) \asymp (\lambda + s)^k \exp \left(- \sum_{i=1}^{M-1} \frac{1}{i} \left(\frac{\lambda}{\lambda + s} \right)^i \sum_{j=1}^k E(Q^j)^i \right). \quad (19)$$

Proof. For brevity we write $\sigma_k(s) = (\lambda + s)^{-k} \pi_k(s)$. Note that $\sigma_k(s) = \prod_{j=1}^k (1 - \mu_j)$, with $\mu_t = \frac{\lambda}{\lambda + s} E(Q^t)$. By taking logarithms we obtain $\log \sigma_k(s) = \sum_{j=1}^k \log(1 - \mu_j)$. We have $\mu_j \leq c = \frac{\lambda}{\lambda + s} EQ < 1$ and from Taylor's formula $\log(1 - x) = - \sum_{i=1}^n \frac{x^i}{i} - \frac{(x/z)^{n+1}}{n+1}$ for all $x \in [0, c)$ and some $z = z(x) \in [1 - c, 1)$. Hence

$$\log \sigma_k(s) = - \sum_{i=1}^{M-1} \frac{1}{i} \sum_{j=1}^k \mu_j^i - d^M C$$

with $d \in [1, \frac{1}{1-c}]$ and $C = \frac{1}{M} \sum_{j=1}^k \mu_j^M < \infty$, and so (19) follows. \square

The next result relates $F_Q(x)$ at $x = 1$ to the Mellin transform of Q at infinity.

Lemma 10. *Let $\beta > 0$. Then the following relations are equivalent:*

$$\begin{aligned} \overline{F}_Q(x) &\sim (1 - x)^\beta L(1 - x) && (x \rightarrow 1) \\ E(Q^t) &\sim \Gamma(\beta + 1) t^{-\beta} L(1/t) && (t \rightarrow \infty) \end{aligned}$$

Proof. Let $W(x) = \overline{F}_Q(e^{-x})$ be the probability distribution function of $W = -\log Q$. Since $(1 - e^{-x})^\beta \sim x^\beta$ and

$$\frac{L(1 - e^{-1/x})}{L(1/x)} = \frac{L(1/x + O(1/x^2))}{L(1/x)} \rightarrow 1, \quad (20)$$

we know that $W(x) \sim x^\beta L(1/x)$ as $x \rightarrow 0$. Therefore, by Karamata's Tauberian theorem (cf. [10])

$$E(Q^t) = \int_0^\infty w^t dF_Q(w) = \int_0^\infty e^{-tw} dW(w) \sim \Gamma(\beta + 1)t^{-\beta}L(1/t),$$

which concludes the proof. \square

Lemma 11 (Foss & Korshunov [18]). *Let $(\xi_n)_{n \in \mathbb{N}}$ be a sequence of i.i.d. non-negative random variables with $E\xi_1^{1/\gamma} < \infty$ for some $\gamma \in (\frac{1}{2}, 1]$ and let $\lambda = E(\xi_1)$. Let N_t be the associated renewal counting process and let X be a random time with tail function $\bar{G}(t) = P(X > t)$. If $\bar{G}(n + h(n)) \sim \bar{G}(n)$ as $n \rightarrow \infty$ for all functions $h(n) = o(n^\gamma)$, then $P(X > \sum_{i=1}^n \xi_i) \sim \bar{G}(\lambda n)$.*

Lemma 12. *Let g be a non-increasing function such that $g(x) \rightarrow 0$ and $g(x+h(x)) \sim g(x)$ as $x \rightarrow \infty$ for all functions $h(n) = o(n^\gamma)$ for some $\gamma \in (\frac{1}{2}, 1]$. Then, as $x \rightarrow \infty$,*

$$\sum_{k=1}^{\infty} g(k) \frac{x^k}{k!} \sim e^x g(x).$$

Proof. We may assume that $g(0) = 1$ and set $\bar{G}(x) = P(X > x) = g(x)$ for some random variable X . Let $(N_t)_{t \geq 0}$ be a Poisson process with intensity one and let $\xi_i = N_i - N_{i-1}$. Then clearly $E\xi_i^a < \infty$ for all $a > 0$, and according to Lemma 11,

$$g(n) \sim P(X > \sum_{i=1}^n \xi_i) = P(X > N_n) = \sum_{k=1}^{\infty} e^{-n} g(k) \frac{n^k}{k!}.$$

Finally note that $P(N_{n+1} < X) \leq P(N_x < X) \leq P(N_n < X)$ for $x \in [n, n+1]$. It follows that $P(N_{n+1} < X)/P(N_n < X) \rightarrow 1$ and $g(n)/g(x) \rightarrow 1$ as $n, x \rightarrow \infty$. \square

Proof of Theorem 7. Clearly if $F_Q(x) \geq F(x)$ then $E_x \tau_y \geq E_x \tau'_y$, where τ'_y is the modified hitting time of a process where F_Q has been replaced by F . Since $\int_1^\infty L(u)/u du < \infty$ it follows from Lemma 10 that $\sum_{k=1}^{\infty} EQ^k < \infty$. In particular the variable M in Lemma 9 is equal to one and thus $\pi_k \asymp \lambda^k$. Applying Lemma 8 yields $E_x \tau_y \asymp e^{\lambda y}$. The inequalities for D are immediate from $x_Q^k \leq E(Q^k) \leq (EQ)^k$. This proves assertion 1.

To prove assertion 2 note that it follows from Lemma 10 that $E(Q^t) \sim \Gamma(\beta + 1)t^{-\beta}L(1/t)$, so that $\int_1^\infty E(Q^t) dt$ is finite and thus $\sum_{k=1}^{\infty} (EQ^k)^2$ is finite. Hence, application of Lemma 9 with $M = 2$ gives

$$\pi_k(s) \asymp \lambda^k \exp \left(- \sum_{j=1}^k E(Q^j) \right) \quad (21)$$

as $k \rightarrow \infty$. Since the function $t \mapsto EQ^t$ is decreasing it follows that

$$\int_1^k E(Q^t) dt \leq \sum_{j=1}^k E(Q^j) \leq \int_1^k E(Q^t) dt + EQ,$$

so that the sum in (21) can be approximated by an integral without changing the asymptotic behavior (w.r.t. \asymp). Let $h(x) = o(x^{\beta-\delta})$ with $\delta \in (0, \beta - \frac{1}{2})$. It follows from Lemma 10 and the mean value theorem that for x large enough there are

constants $c, c' > 0$, such that

$$\begin{aligned} \int_x^{x+h(x)} E(Q^t) dt &\leq c \cdot \int_x^{x+h(x)} \frac{L(1/t)}{t^\beta} dt \leq c \cdot \int_x^{x+h(x)} \frac{1}{t^{\beta-\delta/2}} dt \\ &\leq c' \cdot h(x) \cdot x^{\delta/2-\beta} = o(x^{-\delta/2}). \end{aligned}$$

For $g(x) = \exp(-\int_1^x E(Q^t) dt)$ and $\gamma = \beta - \delta \in (\frac{1}{2}, 1]$, the conditions of Lemma 12 are fulfilled. The first part of assertion 2 follows from

$$\sum_{k=1}^{\infty} \pi_{k-1}(s) \frac{x^k}{k!} \asymp \sum_{k=1}^{\infty} \exp\left(-\int_1^k E(Q^t) dt\right) \frac{(\lambda x)^k}{k!} \sim \exp\left(\lambda x - \int_1^{\lambda x} E(Q^t) dt\right)$$

and

$$\lambda x - \int_1^{\lambda x} E(Q^t) dt = 1 + \frac{1}{\lambda} \int_1^{\lambda x} \lambda(1 - E(Q^t)) dt = 1 + \int_1^x \lambda(1 - E(Q^{\lambda t})) dt.$$

Moreover, Lemma 10 yields

$$\int_1^{\lambda x} E(Q^t) dt \sim \Gamma(1 + \beta) \int_1^x \frac{L(1/t)}{t^\beta} dt,$$

which proves assertion 2a in the $\beta = 1$ case. To see that 2b is true for $\beta < 1$, observe that

$$\Gamma(1 + \beta) \int_1^x \frac{L(1/t)}{t^\beta} dt \sim \frac{\Gamma(1 + \beta)}{1 - \beta} \lambda^{1-\beta} x^{1-\beta} L(1/(\lambda x))$$

from Karamata's integration theorem (see [10]). \square

6. SPECIAL CASE: BETA DISTRIBUTED Q

We assume that Q has a beta distribution with density

$$f_Q(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)},$$

for $x \in [0, 1]$, where $B(\alpha, \beta) = \int_0^1 x^{\alpha-1}(1-x)^{\beta-1} dx = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ and $\alpha > 0, \beta > 0$.

In this case we have

$$\theta_t = \lambda \left(1 - \frac{B(\alpha + t, \beta)}{B(\alpha, \beta)}\right) = \lambda \left(1 - \frac{\Gamma(\alpha + t)\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\alpha + \beta + t)}\right).$$

Lemma 10 yields $E(Q^t) \sim \Gamma(\alpha + \beta)/\Gamma(\alpha) \cdot t^{-\beta}$ as $t \rightarrow \infty$. According to Theorem 7 we then have, for $\beta \in (\frac{1}{2}, 1)$,

$$E_x \tau_y = C \cdot \exp\left(\lambda y - (\lambda y)^{1-\beta} \frac{\Gamma(\alpha + \beta)}{(1 - \beta)\Gamma(\alpha)} + o(y^{1-\beta})\right).$$

In case $\beta = 1$, when $Q \stackrel{d}{=} U^{1/\alpha}$, we can obtain more explicit results and sharper asymptotics.

Corollary 13. *For $x \leq y$ and the mean hitting- and recovery times are given by*

$$E_x \tau_y = \int_x^y \alpha \lambda^{-\alpha} u^{-\alpha} e^{\lambda u} \gamma(\alpha, \lambda u) du, \quad (22)$$

$$E_{(x)} \sigma_x = \alpha \lambda^{-\alpha-1} x^{-\alpha} e^{\lambda x} \gamma(\alpha, \lambda x) - \frac{1}{\lambda}, \quad (23)$$

where $\gamma(\alpha, \lambda u) = \int_0^{\lambda u} t^{\alpha-1} e^{-t} dt$.

Proof. Since $\theta_t = \frac{\lambda t}{\alpha + t}$ it follows that

$$\pi_k(s) = (\lambda + s)^k \frac{(\frac{\alpha s}{\lambda + s} + 1)_k}{(\alpha + 1)_k}, \quad (24)$$

where $(x)_k = x \cdot (x + 1) \cdots (x + k - 1)$ is the Pochhammer symbol. We thus have

$$R'_0(x) = \sum_{k=1}^{\infty} \frac{(\lambda x)^{k-1}}{(k-1)!} \frac{(k-1)!}{(\alpha + 1)_{k-1}} = \alpha \lambda^{-\alpha} x^{-\alpha} e^{\lambda x} \gamma(\alpha, \lambda x).$$

With $E_x \tau_y = R_0(y) - R_0(x)$ and $E_{(x)} \sigma_x = \frac{1}{\lambda} (R'_0(x) - 1)$ this gives the result. \square

Expression (23) leads, using $\lim_{x \rightarrow \infty} \gamma(\alpha, \lambda x) = \Gamma(\alpha)$, to a sharp result for the asymptotics.

Corollary 14. *For $x \leq y$ and $Q \stackrel{d}{=} U^{1/\alpha}$ we have the asymptotics*

$$E_{(y)} \sigma_y \sim E_x \tau_y \sim \frac{\Gamma(\alpha + 1)}{\lambda^{\alpha+1} y^\alpha} e^{\lambda y}. \quad (25)$$

In the special case $Q \stackrel{d}{=} U$, i.e. if $\alpha = 1$, we obtain $R_0(x) = C + \frac{1}{\lambda} (\text{Ei}(\lambda x) - \log x)$ with $\text{Ei}(x) = -\int_{-x}^{\infty} \frac{1}{u} e^{-u} du$ the exponential integral. Consequently,

$$E_x \tau_y = \frac{\text{Ei}(\lambda y) - \text{Ei}(\lambda x) - \log(y/x)}{\lambda} \quad (26)$$

and

$$E_{(x)} \sigma_x = \frac{e^{\lambda x} - 1 - \lambda x}{\lambda^2 x}. \quad (27)$$

The asymptotic result $E_{(y)} \sigma_y \sim E_x \tau_y \sim \frac{1}{\lambda^2 y} e^{\lambda y}$ follows immediately from (25) with $\alpha = 1$. Alternatively, it follows from (26) and the fact that $\text{Ei}(x) \sim \frac{1}{x} e^x$ (see [2], p. 231).

Let ${}_1F_1(a; b; x) = \sum_{k=0}^{\infty} \frac{(a)_k x^k}{(b)_k k!}$ denote Kummer's hypergeometric function (see e.g. [6]).

Corollary 15. *For $x \leq y$ and $Q \stackrel{d}{=} U^{1/\alpha}$ the Laplace transforms of the hitting- and recovery times are given by*

$$\begin{aligned} E_x e^{-s \tau_y} &= \frac{{}_1F_1(\frac{\alpha s}{\lambda + s}; \alpha; (\lambda + s)y)}{{}_1F_1(\frac{\alpha s}{\lambda + s}; \alpha; (\lambda + s)x)}, \\ E_x e^{-s \sigma_x} &= 1 + \frac{s}{\lambda} \left(1 - \frac{{}_1F_1(\frac{\alpha s}{\lambda + s} + 1; \alpha + 1; (\lambda + s)x)}{{}_1F_1(\frac{\alpha s}{\lambda + s}; \alpha; (\lambda + s)x)} \right). \end{aligned}$$

Proof. We get from (4) and (24) that

$$\frac{1}{s} R'_s(x) = {}_1F_1(\frac{\alpha s}{\lambda + s} + 1; \alpha + 1; (\lambda + s)x)$$

and since ${}_1F_1(a; b; x) = \frac{b-1}{a-1} \cdot \frac{d}{dx} {}_1F_1(a-1; b-1; x)$ and $R_s(0) = 1$,

$$1 + s R_s(x) = {}_1F_1(\frac{\alpha s}{\lambda + s}; \alpha; (\lambda + s)x).$$

Application of Theorem 2 then completes the proof. \square

7. SPECIAL CASE: DETERMINISTIC Q

In this section we discuss the case where Q is a constant, say $Q = q \in [0, 1)$. The case $q = 1/2$ corresponds to the standard model for TCP, which is the dominant protocol for data transmission over the internet. The case $q = 0$ describes the age process of renewal theory with exponential renewal epochs.

In general, we have $\theta_a = \lambda(1 - q^a)$ and

$$\prod_{j=1}^k (\theta_{a+j} + u) = (\lambda + u)^k \prod_{j=0}^{k-1} \left(1 - \frac{\lambda q^{a+1}}{\lambda + u} q^j\right) = (\lambda + u)^k \left(\frac{\lambda}{\lambda + u} q^{a+1}; q\right)_k,$$

where $(a; q)_k = \prod_{i=0}^{k-1} (1 - aq^i)$ is the so called q -series (see Askey et al. [2]). Recall that $(q)_\infty = \lim_{k \rightarrow \infty} (q; q)_k$. Specializing Theorem 2 to the deterministic case immediately yields the following result.

Corollary 16. *For $x \leq y$ and $Q = q \in [0, 1)$ the means and Laplace transforms of τ_y and σ_x are given by*

$$E_x \tau_y = \frac{1}{\lambda} \sum_{k=1}^{\infty} \frac{(\lambda y)^k - (\lambda x)^k}{k!} (q; q)_{k-1}, \quad (28)$$

$$E_x e^{-s\tau_y} = \frac{1 + \sum_{k=1}^{\infty} (x(\lambda + s))^k / k! \cdot \left(\frac{\lambda}{\lambda + s}; q\right)_k}{1 + \sum_{k=1}^{\infty} (y(\lambda + s))^k / k! \cdot \left(\frac{\lambda}{\lambda + s}; q\right)_k}, \quad (29)$$

$$E_{(x)} \sigma_x = \frac{1}{\lambda} \sum_{k=1}^{\infty} \frac{(\lambda x)^k}{k!} (q; q)_k, \quad (30)$$

$$E_{(x)} e^{-s\sigma_x} = 1 + \frac{s}{\lambda} \left(1 - \frac{1 + \sum_{k=1}^{\infty} ((\lambda + s)x)^k / k! \cdot \left(\frac{\lambda}{\lambda + s}; q\right)_k}{1 + \sum_{k=1}^{\infty} ((\lambda + s)x)^k / k! \cdot \left(\frac{\lambda}{\lambda + s}; q\right)_k}\right). \quad (31)$$

The asymptotic result below is an immediate consequence of Theorem 7, part 1.

Corollary 17. *For $x \leq y$ and $Q = q \in [0, 1)$ we have the asymptotics*

$$E_{(y)} \sigma_y \sim E_x \tau_y \sim \frac{(q)_\infty}{\lambda} e^{\lambda y}. \quad (32)$$

8. ACKNOWLEDGMENT

We would like to thank Svevolod Shneer for stimulating discussions and for pointing out the connection to the depoissonization Lemma 11.

REFERENCES

- [1] M. Abadi and A. Galves. Inequalities for the occurrence times of rare events in mixing processes. The state of the art. *Markov Process. Relat. Fields*, 7(1):97–112, 2001.
- [2] M. Abramowitz and I.A. Stegun. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Washington: John Wiley & Sons, 1984.
- [3] D.J. Aldous. Markov chains with almost exponential hitting times. *Stochastic Process. Appl.*, 13(3):305–310, 1982.
- [4] E. Altman, K. Avrachenkov, Barakat Ch., and R. Nunez Queija. State-dependent M/G/1 Type Queueing Analysis for Congestion Control in Data Networks. In *INFOCOM*, pages 1350–1359, 2001.
- [5] E. Altman, K. Avrachenkov, A. Kherani, and B. Prabhu. Performance Analysis and Stochastic Stability of Congestion Control Protocols. Technical Report RR-5262, INRIA, Sophia-Antipolis, France, July 2004.

- [6] G.E. Andrews, R. Askey, and R. Roy. *Special Functions*, volume 71 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, 2000.
- [7] S. Asmussen. *Applied Probability and Queues*. John Wiley & Sons, 1987.
- [8] F. Baccelli, D.R. McDonald, and J. Reynier. A mean-field model for multiple TCP connections through a buffer implementing RED. *Perform. Eval.*, 49(1-4):77–97, 2002.
- [9] J. Bertoin and M. Yor. Exponential functionals of Lévy processes. *Probability Surveys*, 2:191–212, 2005.
- [10] N.H. Bingham, C.M. Goldie, and J.L. Teugels. *Regular Variation*, volume 27 of *Encyclopedia of Mathematics and Applications*. Cambridge University Press, 1987.
- [11] R.P. Boas. *Entire Functions*. Pure and applied mathematics, A series of monographs and textbooks. New York: Academic Press Inc., 1954.
- [12] O. Boxma, D. Perry, W. Stadje, and S. Zacks. A Markovian growth-collapse model. *Adv. Appl. Probab.*, 38(1):221–243, 2006.
- [13] P. Carmona, F. Petit, and M. Yor. Exponential functionals of Lévy processes. Barndorff-Nielsen, Ole E. et al. (ed.), *Lévy processes. Theory and applications*. Boston: Birkhäuser. 39-55, 2001.
- [14] M.H.A. Davis. *Markov Models and Optimization.*, volume 49 of *Monographs on Statistics and Applied Probability*. London: Chapman & Hall, 1993.
- [15] V. Dumas, F. Guillemin, and Ph. Robert. A Markovian analysis of additive-increase, multiplicative-decrease (AIMD) algorithms. *Adv. Appl. Probab.*, 34(1):85–111, 2002.
- [16] I. Eliazar and K. Klafter. A growth-collapse model: Lévy inflow, geometric crashes, and generalized Ornstein-Uhlenbeck dynamics. *Physica A*, 334:1–21, 2004.
- [17] S.N. Ethier and T.G. Kurtz. *Markov Processes. Characterization and Convergence*. John Wiley & Sons, 1986.
- [18] S. Foss and D. Korshunov. Sampling at a random time with a heavy-tailed distribution. *Markov Process. Relat. Fields*, 6(4):543–568, 2000.
- [19] F. Guillemin, Ph. Robert, and B. Zwart. AIMD algorithms and exponential functionals. *Ann. Appl. Probab.*, 14(1):90–117, 2004.
- [20] Ph. Jacquet and W. Szpankowski. Analytical depoissonization and its applications. *Theor. Comput. Sci.*, 201(1-2):1–62, 1998.
- [21] O. Kella and W. Stadje. On hitting times for compound Poisson dams with exponential jumps and linear release rate. *J. Appl. Probab.*, 38(3):781–786, 2001.
- [22] A. Lachal. First exit time from a bounded interval for a certain class of additive functionals of Brownian motion. *J. Theor. Probab.*, 13(3):733–775, 2000.
- [23] N. Litvak and I.J.B.F. Adan. The travel time in carousel systems under the nearest item heuristic. *J. Appl. Probab.*, 38:45–54, 2001.
- [24] N. Litvak and W.R. van Zwet. On the minimal travel times needed to collect n items on a circle. *Ann. Appl. Probab.*, 14(2):881–902, 2004.
- [25] A.H. Löpker. Some martingales for Markov processes. Technical Report 2006-36, ISSN 1389-2355, Eurandom, Eindhoven, 2006.
- [26] A.H. Löpker and J.S.H. van Leeuwen. Transient moments of the window size in TCP. Technical Report 2007-19, ISSN 1389-2355, Eurandom, Eindhoven, 2007.
- [27] T. Ott, J. Kemperman, and M. Mathis. The stationary behavior of ideal TCP congestion avoidance. <ftp://ftp.bellcore.com/pub/tjo/TCPwindow.ps>.
- [28] Ph. Robert. On the asymptotic behavior of some algorithms. *Random Struc. Alg.*, 27:235–250, 2005.
- [29] T. Rolski, H. Schmidli, V. Schmidt, and J. Teugels. *Stochastic Processes for Insurance and Finance*. New York: John Wiley & Sons, 1999.