

On the number of positive integers $\leq x$ and free of prime factors $> y$

Citation for published version (APA):

Erdős, P., & van Lint, J. H. (1966). On the number of positive integers $\leq x$ and free of prime factors $> y$. *Simon Stevin : Wis- en Natuurkundig Tijdschrift*, 40(11), 73-76.

Document status and date:

Published: 01/01/1966

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

ON THE NUMBER OF POSITIVE INTEGERS
 $\leq x$ AND FREE OF PRIME FACTORS $> y$

P. ERDÖS

University of Illinois
Urbana, Illinois

and

J. H. VAN LINT

Bell Telephone Laboratories, Incorporated
Murray Hill, New Jersey

1. Introduction

Let $\Psi(x, y)$ denote the number of integers specified in the title. A number of estimates and asymptotic formulae for this function have been given (cf. [1] and the literature mentioned there). Recently DE BRUIJN ([2]) proved an asymptotic formula for $\log \Psi(x, y)$ which holds uniformly for $2 < y \leq x$. Part of the proof consisted of showing that

$$\Psi(x, y) \geq \binom{\pi(y) + u}{u} \text{ where } u = [(\log x)/(\log y)].$$

It is the purpose of this note to extend this inequality to an asymptotic formula (which is weaker than DE BRUIJN's result). In fact we shall prove:

Theorem 1: For $2 < y \leq x$ we have for $x \rightarrow \infty$, uniformly in y ,

$$\log \Psi(x, y) \sim \log \binom{\pi(y) + u}{u} \quad (1)$$

where

$$u = [(\log x)/(\log y)].$$

We remark that this of course follows from DE BRUIJN's theorem. Our interest lies mainly in giving a short fairly straightforward proof. For some ranges of values of y (1) is nearly trivial and the most interesting part of the proof concerns the range $(\log x)^{\varepsilon} < y < (\log x)^{1+\varepsilon}$.

2. Proof of Theorem 1

We shall prove (1) by showing that for every $\varepsilon > 0$ we have

$$\binom{\pi(y)+u}{u} < \Psi(x,y) < \binom{\pi(y)+u}{u}^{1+\varepsilon} \quad \text{for } x > x_0(\varepsilon). \quad (2)$$

- a. The first inequality immediately follows from the fact that $\binom{\pi(y)+u}{u}$ represents the number of solutions of

$$\sum_{p \leq y} \alpha_p \leq u = [(\log x)/(\log y)]$$

in nonnegative integers α_p and this number is less than the number of solutions of

$$\sum_{p \leq y} \alpha_p \log p \leq \log x$$

which is $\Psi(x,y)$ by definition.

In sections **b**, **c** and **d** we prove the second inequality of (2).

- b. We now consider $y < (\log x)^{1+\varepsilon}$. We first remark that (2) is trivial for very small values of y , for example $y < (\log x)^{\varepsilon/2}$, because

$$\Psi(x,y) \leq \left(\frac{\log x}{\log 2} + 1\right)^{\pi(y)} < \binom{\pi(y)+u}{u}^{1+\varepsilon} \quad \text{if } y < (\log x)^{\varepsilon/2}.$$

Hence we can assume that $y > (\log x)^{\varepsilon/2}$.

Let N_1 denote the number of integers $\leq x$, free of prime factors $> y^{1-\varepsilon}$ and let N_2 denote the number of integers $\leq x$ all of whose prime factors satisfy $y^{1-\varepsilon} < p \leq y$. Then $\Psi(x,y) \leq N_1 N_2$.

Trivially $N_1 < \left(\frac{\log x}{\log 2} + 1\right)^{\pi(y^{1-\varepsilon})}$. Furthermore

$\binom{\pi(y)+u}{u} > \left(\frac{\pi(y)+u}{\pi(y)}\right)^{\pi(y)}$. From this it follows that

$$\frac{\log N_1}{\log \binom{\pi(y)+u}{u}} = O\left(\frac{\log \log x}{(\log x)^{\frac{1}{2}\varepsilon^2}}\right)$$

i.e. $N_1 < \binom{\pi(y)+u}{u}^\varepsilon$ for $x > x_1(\varepsilon)$.

Now N_2 is less than the number of solutions of

$$\sum_{y^{1-\varepsilon} < p \leq y} \alpha_p \leq \frac{\log x}{(1-\varepsilon)\log y}$$

in nonnegative integers α_p and this number does not exceed

$$\binom{\pi(y)+u'}{u'} \text{ where } u' = \left\lfloor \frac{\log x}{(1-\varepsilon)\log y} \right\rfloor.$$

We now use the fact that if a, b and $b(1+\varepsilon)$ are positive integers then

$$\binom{a+b}{a}^{1+\varepsilon} = \prod_{i=0}^{a-1} \left(1 + \frac{b}{a-i}\right)^{1+\varepsilon} > \prod_{i=0}^{a-1} \left(1 + \frac{b(1+\varepsilon)}{a-i}\right) = \binom{a+b(1+\varepsilon)}{a}.$$

It follows that $N_2 < \binom{\pi(y)+u}{u}^{1+o(\varepsilon)}$.

Combining the estimates for N_1 and N_2 we find

$$\Psi(x, y) < \binom{\pi(y)+u}{u}^{1+o(\varepsilon)}$$

proving (2) for $y < (\log x)^{1+\varepsilon}$.

c. For $y > (\log x)^{n(\varepsilon)}$ where for instance $n(\varepsilon) = 2/\varepsilon$ the right-hand side of (2) is trivial because

$$\binom{\pi(y)+u}{u} > \left(\frac{\pi(y)}{u}\right)^u \text{ which implies } \binom{\pi(y)+u}{u}^{1+\varepsilon} > x.$$

d. The case $(\log x)^{1+\varepsilon} < y < (\log x)^{2/\varepsilon}$ will be treated by writing $y = (\log x)^\alpha$ and proving

$$\Psi(x, y) = x^{1-1/\alpha+o(1)} \tag{3}$$

which implies (2).

We first remark that $\binom{\pi(y)+u}{u} = x^{1-1/\alpha+o(1)}$ and the same holds for $\binom{\pi(y)}{u}$. So it remains to show that

$$\Psi(x, y) < x^{1-1/\alpha+o(1)}.$$

To do this we split the integers counted by $\Psi(x, y)$ into two classes. First those with at least u distinct prime factors. Their number is less than

$$\frac{x}{u!} \left(\sum_{p < x} \frac{1}{p} \right)^u < x \left(\frac{2e \log \log x}{u} \right)^u = x^{1-1/u+o(1)}.$$

The number of integers $\leq x$ with less than u distinct prime factors $\leq y$ is less than

$$\binom{\pi(y)}{u} \binom{\left(\frac{\log x}{\log 2} + u\right)}{u}$$

because there are $\binom{\pi(y)}{u}$ different u -tuples of primes $\leq y$ and for each of these the sum of the exponents is less than $\frac{\log x}{\log 2}$. We have

$$\binom{\left(\frac{\log x}{\log 2} + u\right)}{u} = x^{o(1)} \left(\text{this follows from } \binom{n}{k} < \frac{n^k}{k!} < \left(\frac{ne}{k}\right)^k \right).$$

So the number of integers in the second class is also $x^{1-1/u+o(1)}$.

This completes the proof of Theorem 1.

REFERENCES

- [1] N.G. DE BRUIJN, On the number of positive integers $\leq x$ and free of prime factors $> y$, *Proc. Kon. Ned. Akad. v. Wetensch.*, 54, 50-59 (1951).
- [2] N.G. DE BRUIJN, On the number of positive integers $\leq x$ and free of prime factors $> y$, II, *Proc. Kon. Ned. Akad. v. Wetensch.*, 69, 335-348 (1966).