

Novel ultrasound-contrast-agent dilution method for assessment of left ventricular ejection fraction

Citation for published version (APA):

Jansen, A. H. M., Misch, M., Dantzig, van, J. M., Peels, C. H., Bracke, F. A. L. E., Hemel, van, N. M., & Korsten, H. H. M. (2006). Novel ultrasound-contrast-agent dilution method for assessment of left ventricular ejection fraction. *Circulation*, 114(18 SII), 479-479.

Document status and date:

Published: 01/01/2006

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

(*Circulation*. 2006;114:II_479.)

© 2006 American Heart Association, Inc.

Echocardiography: Stress, Contrast and Cardiac Function I

Abstract 2340: Novel Ultrasound-Contrast-Agent Dilution Method for Assessment of Left Ventricular Ejection Fraction

Annemieke H Jansen¹; Massimo Mischi²; Jan melle van Dantzig³;
Frank Bracke³; Kathinka Peels³; Norbert M van Hemel⁴;
Hendrikus H Korsten⁵

¹ Catharina Hosp, Eindhoven, The Netherlands

² Technical Univ Eindhoven, Eindhoven, The Netherlands

³ Catharina Hosp, Eindhoven, The Netherlands

⁴ Univ of Utrecht, Utrecht, The Netherlands

⁵ Catharina Hosp, Eindhoven, The Netherlands

Background: Left ventricular ejection fraction (LVEF) is an important determinant of prognosis. We evaluated the accuracy of a novel fast method for LVEF quantification based on indicator dilution curve (IDC) principles (figure 1) and compared the results with contrast-enhanced biplane LVEF assessment.

Method: a 10 ml diluted (1:100) ultrasound-contrast bolus (SonoVue ®) was injected intravenously in thirty patients (pts) (20 male, age 65 ± 10) with known or suspected heart disease. In 22 pts multiple recordings were made and in 12 pts injections were repeated after implantation of a biventricular pacemaker, leading to 68 measurements. The developed algorithm used the left atrium and LV IDC for the LVEF measurement. For the biplane enhanced LVEF measurements a 0.5 ml pure ultrasound-contrast bolus (SonoVue ®) was administered to obtain multiple four- and two-chamber recordings.

Results: according to contrast enhanced biplane assessments, the LVEF ranged from 10 to 75 % while the LV end-diastolic volume ranged from 80 to 521 ml. The mean LVEF measured by the biplane and the IDC method was 33 ± 17 % and 35 ± 18 %, respectively. A correlation coefficient r=0.91 was observed between the two methods (figure 2).

This Article

- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

Services

- ▶ [Email this article to a friend](#)
- ▶ [Similar articles in this journal](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)
- ▶ [Request Permissions](#)

Google Scholar

- ▶ [Articles by Jansen, A. H](#)
- ▶ [Articles by Korsten, H. H](#)

PubMed

- ▶ [Articles by Jansen, A. H](#)
- ▶ [Articles by Korsten, H. H](#)

Conclusion: A new fast method for LVEF assessment based on IDC principles is described and comparison with contrast enhanced biplane LVEF quantification shows accurate results. The proposed method also allows simultaneous quantification of right ventricular EF based on the same IDC principles.

