

Variational characterization of the critical curve for pinning of random polymers

Citation for published version (APA):

Cheliotis, D., & Hollander, den, W. T. F. (2010). *Variational characterization of the critical curve for pinning of random polymers*. (Report Eurandom; Vol. 2010024). Eurandom.

Document status and date:

Published: 01/01/2010

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EURANDOM PREPRINT SERIES
2010-024

**Variational characterization of the critical curve
for pinning of random polymers**

D. Cheliotis, F. den Hollander
ISSN 1389-2355

Variational characterization of the critical curve for pinning of random polymers

D. Cheliotis ¹
F. den Hollander ^{2 3}

May 18, 2010

Abstract

In this paper we look at the pinning of a directed polymer by a one-dimensional linear interface carrying random charges. There are two phases, localized and delocalized, depending on the inverse temperature and on the disorder bias. Using quenched and annealed large deviation principles for the empirical process of words drawn from a random letter sequence according to a random renewal process (Birkner, Greven and den Hollander [6]), we derive variational formulas for the quenched, respectively, annealed critical curve separating the two phases. These variational formulas are used to obtain a necessary and sufficient criterion, stated in terms of relative entropies, for the two critical curves to be different at a given inverse temperature, a property referred to as relevance of the disorder. This criterion in turn is used to show that the regimes of relevant and irrelevant disorder are separated by a unique inverse critical temperature. Subsequently, upper and lower bounds are derived for the inverse critical temperature, from which sufficient conditions under which it is strictly positive, respectively, finite are obtained. The former condition is believed to be necessary as well, a problem that we will address in a forthcoming paper.

Random pinning has been studied extensively in the literature. The present paper opens up a window with a variational view. Our variational formulas for the quenched and the annealed critical curve are new and provide valuable insight into the nature of the phase transition. Our results on the inverse critical temperature drawn from these variational formulas are not new, but they offer an alternative approach that is flexible enough to be extended to other models of random polymers with disorder.

MSC 2000. Primary 60F10, 60K37; Secondary 82B27, 82B44.

Key words and phrases. Random polymer, random charges, localization vs. delocalization, quenched vs. annealed large deviation principle, quenched vs. annealed critical curve, relevant vs. irrelevant disorder, critical temperature.

Acknowledgment. The research in this paper was carried out while DC was a postdoc at EURANDOM. It was partially supported by the DFG-NWO Bilateral Research Group “Mathematical Models from Physics and Biology”.

¹Department of Mathematics, University of Athens, Panepistimiopolis, 15784 Athens, Greece,
dcheliotis@math.uoa.gr

²Mathematical Institute, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands,
denholla@math.leidenuniv.nl

³EURANDOM, P.O. Box 513, 5600 MB Eindhoven, The Netherlands,

1 Introduction and main results

1.1 Introduction

I. Model. Let $S = (S_n)_{n \in \mathbb{N}_0}$ be a Markov chain on a countable state space \mathcal{S} in which a given point is marked 0 ($\mathbb{N}_0 = \mathbb{N} \cup \{0\}$). Write \mathbf{P} to denote the law of S given $S_0 = 0$ and \mathbf{E} the corresponding expectation. Let K denote the distribution of the first return time of S to 0, i.e.,

$$K(n) := \mathbf{P}(S_n = 0, S_m \neq 0 \forall 0 < m < n), \quad n \in \mathbb{N}. \quad (1.1)$$

We will assume that $\sum_{n \in \mathbb{N}} K(n) = 1$ (i.e., 0 is a recurrent state) and

$$\lim_{n \rightarrow \infty} \frac{\log K(n)}{\log n} = -(1 + \alpha) \quad \text{for some } \alpha \in [0, \infty). \quad (1.2)$$

Let $\omega = (\omega_k)_{k \in \mathbb{N}_0}$ be i.i.d. \mathbb{R} -valued random variables with marginal distribution μ_0 . Write $\mathbb{P} = \mu_0^{\otimes \mathbb{N}_0}$ to denote the law of ω , and \mathbb{E} to denote the corresponding expectation. We will assume that

$$M(\lambda) := \mathbb{E}(e^{\lambda \omega_0}) < \infty \quad \forall \lambda \in \mathbb{R}, \quad (1.3)$$

and that μ_0 has mean 0 and variance 1.

Let $\beta \in [0, \infty)$ and $h \in \mathbb{R}$, and for fixed ω define the law $\mathbf{P}_n^{\beta, h, \omega}$ on $\{0\} \times \mathcal{S}^n$, the set of n -steps paths in \mathcal{S} starting from 0, by putting

$$\frac{d\mathbf{P}_n^{\beta, h, \omega}}{d\mathbf{P}_n}((S_k)_{k=0}^n) := \frac{1}{Z_n^{\beta, h, \omega}} \exp \left[\sum_{k=0}^{n-1} (\beta \omega_k - h) 1_{\{S_k=0\}} \right] 1_{\{S_n=0\}}, \quad (1.4)$$

where \mathbf{P}_n is the projection of \mathbf{P} onto $\{0\} \times \mathcal{S}^n$. Here, β plays the role of the inverse temperature, h the role of the disorder bias, while $Z_n^{\beta, h, \omega}$ is the normalizing partition sum. Note that $k = 0$ contributes to the sum while $k = n$ does not, and that the path is tied to 0 at both ends. This is done for later convenience.




Figure 1: A directed polymer sampling random charges at an interface.

Remark 1.1. Note that (1.2) implies $p := \gcd[\text{supp}(K)] = 1$. If $p \geq 2$, then the model can be trivially restricted to $p\mathbb{N}$, so there is no loss of generality. Moreover, if $\sum_{n \in \mathbb{N}} K(n) < 1$, then the model can be reduced to the recurrent case by a shift of h . Similarly, the restriction to μ_0 with mean 0 and variance 1 can be removed by a scaling of β and a shift of h .

Remark 1.2. The key example of the above setting is a simple random walk on \mathbb{Z} , for which $p = 2$ and $\alpha = \frac{1}{2}$ (Spitzer [20], Section 1). In that case the process $(n, S_n)_{n \in \mathbb{N}_0}$ can be thought of as describing a directed polymer in $\mathbb{N}_0 \times \mathbb{Z}$ that is pinned to the interface $\mathbb{N}_0 \times \{0\}$ by random charges ω (see Fig. 1). When the polymer hits the interface at time k , it picks up a reward $\exp[\beta \omega_k - h]$, which can be either > 1 or < 1 depending on the value of ω_k . For $h \leq 0$ the

polymer tends to intersect the interface with a positive frequency (“localization”), whereas for $h > 0$ large enough it tends to wander away from the interface (“delocalization”). Simple random walk on \mathbb{Z}^2 corresponds to $p = 2$ and $\alpha = 0$, while simple random walk on \mathbb{Z}^d , $d \geq 3$, conditioned on returning to 0 corresponds to $p = 2$ and $\alpha = \frac{d}{2} - 1$ (Spitzer [20], Section 1).

II. Free energy and phase transition. The *quenched free energy* is defined as

$$f^{\text{que}}(\beta, h) := \lim_{n \rightarrow \infty} \frac{1}{n} \log Z_n^{\beta, h, \omega}. \quad (1.5)$$

Standard subadditivity arguments show that the limit exists ω -a.s. and in \mathbb{P} -mean, and is non-random (see e.g. Giacomin [11], Chapter 5, and den Hollander [18], Chapter 11). Moreover, $f^{\text{que}}(\beta, h) \geq 0$ because $Z_n^{\beta, h, \omega} \geq e^{\beta\omega_0 - h} K(n)$, $n \in \mathbb{N}$, and $\lim_{n \rightarrow \infty} \frac{1}{n} \log K(n) = 0$ by (1.2). The lower bound $f^{\text{que}}(\beta, h) = 0$ is attained when S visits the state 0 only rarely. This motivates the definition of two quenched phases:

$$\begin{aligned} \mathcal{L} &:= \{(\beta, h) : f^{\text{que}}(\beta, h) > 0\}, \\ \mathcal{D} &:= \{(\beta, h) : f^{\text{que}}(\beta, h) = 0\}, \end{aligned} \quad (1.6)$$

referred to as the *localized* phase, respectively, the *delocalized* phase.

Since $h \mapsto f^{\text{que}}(\beta, h)$ is non-increasing for every $\beta \in [0, \infty)$, the two phases are separated by a *quenched critical curve*

$$h_c^{\text{que}}(\beta) := \inf \{h : f^{\text{que}}(\beta, h) = 0\}, \quad \beta \in [0, \infty). \quad (1.7)$$

with \mathcal{L} the region below the curve and \mathcal{D} the region on and above. Since $(\beta, h) \mapsto f^{\text{que}}(\beta, h)$ is convex and $\mathcal{D} = \{(\beta, h) : f^{\text{que}}(\beta, h) \leq 0\}$ is a level set of f^{que} , it follows that \mathcal{D} is a convex set and h_c^{que} is a convex function. Since $\beta = 0$ corresponds to a homopolymer, we have $h_c^{\text{que}}(0) = 0$ (see Appendix A). It was shown in Alexander and Sidoravicius [2] that $h_c^{\text{que}}(\beta) > 0$ for $\beta \in (0, \infty)$. Therefore we have the qualitative picture drawn in Fig. 2. We further remark that $\lim_{\beta \rightarrow \infty} h_c^{\text{que}}(\beta)/\beta$ is finite if and only if $\text{supp}(\mu_0)$ is bounded from above.




Figure 2: Qualitative plot of $\beta \mapsto h_c^{\text{que}}(\beta)$. The fine details of this curve are not known.

The mean value of the disorder is $\mathbb{E}(\beta\omega_0 - h) = -h$. Thus, we see from Fig. 2 that for the random pinning model localization may even occur for *moderately negative* mean values of the disorder, contrary to what happens for the homogeneous pinning model, where localization occurs only for strictly positive parameter (see Appendix A). In other words, even a globally repulsive random interface can pin the polymer: all that the polymer needs to do is to hit the positive values of the disorder and avoid the negative values as much as possible.

The *annealed free energy* is defined by

$$f^{\text{ann}}(\beta, h) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{E}(Z_n^{\beta, h, \omega}). \quad (1.8)$$

Since

$$\mathbb{E}(Z_n^{\beta, h, \omega}) = \mathbf{E} \left(\exp \left[\sum_{k=0}^{n-1} [\log M(\beta) - h] 1_{\{S_k=0\}} \right] 1_{\{S_n=0\}} \right), \quad (1.9)$$

we have that $f^{\text{ann}}(\beta, h)$ is the free energy of the homopolymer with parameter $\log M(\beta) - h$. The associated *annealed critical curve*

$$h_c^{\text{ann}}(\beta) := \inf\{h: f^{\text{ann}}(\beta, h) = 0\}, \quad \beta \in [0, \infty), \quad (1.10)$$

therefore equals

$$h_c^{\text{ann}}(\beta) = \log M(\beta). \quad (1.11)$$

Since $f^{\text{que}} \leq f^{\text{ann}}$, we have $h_c^{\text{que}} \leq h_c^{\text{ann}}$. The disorder is said to be *relevant* for a given choice of K , μ_0 and β when $h_c^{\text{que}}(\beta) < h_c^{\text{ann}}(\beta)$, otherwise it is said to be *irrelevant*. Our main focus in the present paper will be on deriving *variational formulas* for h_c^{que} and h_c^{ann} and investigating under what conditions on K , μ_0 and β the disorder is relevant, respectively, irrelevant.

1.2 Main results

This section contains three theorems and four corollaries, all valid subject to (1.2–1.3). To state these we need some further notation.

I. Notation. Abbreviate

$$E := \text{supp}[\mu_0] \subset \mathbb{R}. \quad (1.12)$$

Let $\tilde{E} := \cup_{k \in \mathbb{N}} E^k$ be the set of *finite words* consisting of *letters* drawn from E . Let $\mathcal{P}(\tilde{E}^{\mathbb{N}})$ denote the set of probability measures on *infinite sentences*, equipped with the topology of weak convergence. Write $\tilde{\theta}$ for the left-shift acting on $\tilde{E}^{\mathbb{N}}$, and $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ for the set of probability measures that are invariant under $\tilde{\theta}$.

For $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, let $\pi_{1,1}Q \in \mathcal{P}(E)$ denote the projection of Q onto the first letter of the first word. Define the set

$$\mathcal{C} := \left\{ Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : \int_E |x| d(\pi_{1,1}Q)(x) < \infty \right\}, \quad (1.13)$$

and on this set the function

$$\Phi(Q) := \int_E x d(\pi_{1,1}Q)(x), \quad Q \in \mathcal{C}. \quad (1.14)$$

We also need two rate functions on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, denoted by I^{ann} and I^{que} , which will be defined in Section 2. These are the rate functions of the annealed and the quenched large deviation principle that play a central role in the present paper, and they satisfy $I^{\text{que}} \geq I^{\text{ann}}$.

II. Theorems. With the above ingredients, we obtain the following *characterization of the critical curves*.

Theorem 1.3. Fix μ_0 and K . For all $\beta \in [0, \infty)$,

$$h_c^{\text{que}}(\beta) = \sup_{Q \in \mathcal{C}} [\beta \Phi(Q) - I^{\text{que}}(Q)], \quad (1.15)$$

$$h_c^{\text{ann}}(\beta) = \sup_{Q \in \mathcal{C}} [\beta \Phi(Q) - I^{\text{ann}}(Q)]. \quad (1.16)$$

We know that $h_c^{\text{ann}}(\beta) = \log M(\beta)$. However, the variational formula for $h_c^{\text{ann}}(\beta)$ will be important for the comparison with $h_c^{\text{que}}(\beta)$.

Next, let

$$d\mu_\beta(x) := \frac{1}{M(\beta)} e^{\beta x} d\mu_0(x), \quad x \in E, \quad (1.17)$$

and

$$q_\beta(x_1, x_2, \dots, x_n) := K(n) \mu_\beta(x_1) \mu_0(x_2) \times \dots \times \mu_0(x_n), \quad n \in \mathbb{N}, x_1, x_2, \dots, x_n \in E. \quad (1.18)$$

Further, let $Q_\beta := q_\beta^{\otimes \mathbb{N}} \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$. Then Q_0 is the probability measure under which the words are i.i.d., with length drawn from K and i.i.d. letters drawn from μ_0 , while Q_β differs from Q_0 in that the first letter of each word is drawn from the tilted probability distribution μ_β . We will see that Q_β is the unique maximizer of the supremum in (1.16) (note that $Q_\beta \in \mathcal{C}$ because of (1.3)). This leads to the following *necessary and sufficient criterion for disorder relevance*.

Theorem 1.4. Fix μ_0 and K . For all $\beta \in [0, \infty)$,

$$h_c^{\text{que}}(\beta) < h_c^{\text{ann}}(\beta) \iff I^{\text{que}}(Q_\beta) > I^{\text{ann}}(Q_\beta). \quad (1.19)$$

We will see that also the supremum in (1.15) is attained, which is to be interpreted as saying that there is a localization strategy *at* the quenched critical line.

Disorder relevance is *monotone* in β (see Fig. 3).

Theorem 1.5. For all μ_0 and K there exists a $\beta_c = \beta_c(\mu_0, K) \in [0, \infty]$ such that

$$h_c^{\text{que}}(\beta) \begin{cases} = h_c^{\text{ann}}(\beta) & \text{if } \beta \in [0, \beta_c], \\ < h_c^{\text{ann}}(\beta) & \text{if } \beta \in (\beta_c, \infty). \end{cases} \quad (1.20)$$




Figure 3: Uniqueness of the critical inverse temperature β_c .

III. Corollaries. From Theorems 1.3–1.5 we draw four corollaries. Abbreviate

$$\chi := \sum_{n \in \mathbb{N}} [\mathbf{P}(S_n = 0)]^2, \quad w := \sup[\text{supp}(\mu_0)]. \quad (1.21)$$

Corollary 1.6. *If $\alpha = 0$, then $\beta_c = \infty$ for all μ_0 .*

Corollary 1.7. *If $\alpha \in (0, \infty)$, then the following bounds hold:*

(i) $\beta_c \geq \beta_c^*$ with $\beta_c^* = \beta_c^*(\mu_0, K) \in (0, \infty]$ given by

$$\beta_c^* := \sup \{ \beta: M(2\beta)/M(\beta)^2 < 1 + \chi^{-1} \}. \quad (1.22)$$

(ii) $\beta_c \leq \beta_c^{**}$ with $\beta_c^{**} = \beta_c^{**}(\mu_0, K) \in (0, \infty]$ given by

$$\beta_c^{**} := \inf \{ \beta: h(\mu_\beta | \mu_0) > h(K) \}, \quad (1.23)$$

where $h(\mu_\beta | \mu_0) = \int_E \log(d\mu_\beta/d\mu_0) d\mu_\beta$ is the relative entropy of μ_β w.r.t. μ_0 , and $h(K) := -\sum_{n \in \mathbb{N}} K(n) \log K(n)$ is the entropy of K .

Corollary 1.8. *If $\alpha \in (0, \infty)$ and $\chi < \infty$, then $\beta_c > 0$ for all μ_0 .*

Corollary 1.9. *If $\alpha \in (0, \infty)$, then $\beta_c < \infty$ for all μ_0 with $\mu_0(\{w\}) = 0$ (which includes $w = \infty$).*

We close with a conjecture stating that the condition $\chi < \infty$ in Corollary 1.8 is not only sufficient for $\beta_c > 0$ but also necessary. This conjecture will be addressed in a forthcoming paper.

Conjecture 1.10. *If $\alpha \in (0, \infty)$ and $\chi = \infty$, then $\beta_c = 0$.*

1.3 Discussion

I. What is known from the literature? Before discussing the results in Section 1.2, we give a summary of what is known about the issue of relevant vs. irrelevant disorder from the literature. This summary is drawn from the papers by Alexander [1], Toninelli [21], [22], Giacomin and Toninelli [14], Derrida, Giacomin, Lacoïn and Toninelli [8], Alexander and Zygouras [3], [4], Giacomin, Lacoïn and Toninelli [12], [13], and Lacoïn [19].

Theorem 1.11. *Suppose that condition (1.2) is strengthened to*

$$K(n) = n^{-(1+\alpha)} L(n) \text{ with } \alpha \in [0, \infty) \text{ and } L \text{ strictly positive and slowly varying at infinity.} \quad (1.24)$$

Then

- (1) $\beta_c = 0$ when $\alpha \in (\frac{1}{2}, \infty)$.
- (2) $\beta_c = 0$ when $\alpha = \frac{1}{2}$ and $L(\infty) \in [0, \infty)$ or $\lim_{N \rightarrow \infty} [L(N)]^{-1} \sum_{n=1}^N n^{-1} [L(n)]^{-2} = \infty$.
- (3) $\beta_c > 0$ when $\alpha = \frac{1}{2}$ and $\sum_{n \in \mathbb{N}} n^{-1} [L(n)]^{-2} < \infty$.
- (4) $\beta_c > 0$ when $\alpha \in (0, \frac{1}{2})$.
- (5) $\beta_c = \infty$ when $\alpha = 0$.

The results in Theorem 1.11 hold irrespective of the choice of μ_0 . Toninelli [22] proves that if $\log M(\lambda) \sim C\lambda^\gamma$ as $\lambda \rightarrow \infty$ for some $C \in (0, \infty)$ and $\gamma \in (1, \infty)$, then $\beta_c < \infty$ irrespective of $\alpha \in (0, \infty)$ and L . Note that there is a small gap between cases (2) and (3) at the critical threshold $\alpha = \frac{1}{2}$.

For the cases of relevant disorder, bounds on the gap between $h_c^{\text{ann}}(\beta)$ and $h_c^{\text{que}}(\beta)$ have been derived in the above cited papers subject to (1.24). As $\beta \downarrow 0$, this gap decays like

$$h_c^{\text{ann}}(\beta) - h_c^{\text{que}}(\beta) \asymp \begin{cases} \beta^2, & \text{if } \alpha \in (1, \infty), \\ \beta^2 \psi(1/\beta), & \text{if } \alpha = 1, \\ \beta^{2\alpha/(2\alpha-1)}, & \text{if } \alpha \in (\frac{1}{2}, 1), \end{cases} \quad (1.25)$$

for all choices of L , with ψ slowly varying and vanishing at infinity when $L(\infty) \in (0, \infty)$.

Partial results are known for $\alpha = \frac{1}{2}$. For instance, when $\sum_{n \in \mathbb{N}} n^{-1} [L(n)]^{-2} = \infty$ the gap decays faster than any polynomial, which implies that the disorder can at most be *marginally relevant*, a situation where standard perturbative arguments do not work. When $L(\infty) \in (0, \infty)$, the gap lies between $\exp[-\beta^{-4}]$ and $\exp[-\beta^{-2}]$ for $\beta \in (0, \infty)$ small enough, modulo constants in the exponent. When $L(n) = O([\log n]^{-\frac{1}{2}-\theta})$, $n \rightarrow \infty$, $\theta \in (0, \infty)$, the gap lies above $\exp[-\beta^{-\theta}]$ for all $\theta' \in (0, \theta)$ and $\beta \in (0, \infty)$ small enough. Both cases correspond to marginal relevance.

Remark 1.12. Most of the above mentioned results are proved for Gaussian disorder only, and in the cited papers it is stated somewhat loosely that proofs carry over to arbitrary disorder subject to (1.3).

Remark 1.13. The fact that $\alpha = \frac{1}{2}$ is critical for relevant vs. irrelevant disorder is in accordance with the so-called *Harris criterion* for disordered systems (see Harris [17]): “Arbitrary weak disorder modifies the nature of a phase transition when the order of the phase transition in the non-disordered system is < 2 ”. The order of the phase transition for the homopolymer, which is briefly described in Appendix A, is < 2 precisely when $\alpha \in (\frac{1}{2}, \infty)$ (see Giacomin [11], Chapter 2). This link is emphasized in Toninelli [21].

II. What is new in the present paper? The main importance of our results in Section 1.2 is that they *open up a new window* on the random pinning problem. Whereas the results cited in Theorem 1.11 are derived with the help of a variety of *estimation techniques*, like fractional moment estimates and trial choices of localization strategies, Theorem 1.3 gives a *variational characterization* of the critical curves that is new (it is very rare indeed that critical curves for disordered systems allow for a direct variational representation). Theorem 1.4 gives a necessary and sufficient criterion for disorder relevance that is explicit and tractable, although not easy to handle. Theorem 1.5 shows that uniqueness of the inverse critical temperature is a direct consequence of this criterion, while Corollaries 1.6–1.9 show that the criterion can be used to obtain important information on the inverse critical temperature.

Remark 1.14. Theorem 1.5 was proved in Giacomin, Lacoïn and Toninelli [13] with the help of the FKG-inequality.

Remark 1.15. Corollary 1.6 is the main result in Alexander and Zygouras [4].

Remark 1.16. Since (see Section 8)

$$\lim_{\beta \rightarrow \infty} M(2\beta)/M(\beta)^2 = 1/\mu_0(\{w\}), \quad \lim_{\beta \rightarrow \infty} h(\mu_\beta | \mu_0) = \log [1/\mu_0(\{w\})], \quad (1.26)$$

with the understanding that the second limit is ∞ when $\mu_0(\{w\}) = 0$, Corollary 1.7 implies Corollaries 1.8–1.9.

Remark 1.17. Note that $\chi = \mathbf{E}(|I_1 \cap I_2|)$ with I_1, I_2 two independent copies of the set of return times of S (recall (1.1)). Thus, according to Corollary 1.8 and Conjecture 1.10, $\beta_c > 0$ is expected to be equivalent to the renewal process of *joint return times* to be recurrent. Note that $1/\mathbf{P}(I_1 \cap I_2 \neq \emptyset) = 1 + \chi^{-1}$, the quantity appearing in Corollary 1.7(i).

Remark 1.18. For Gaussian disorder (with $\mu_0(\{w\}) = 0$) we have $\beta_c^* \in (0, \infty)$ if and only if $\chi \in (0, \infty)$, while for Bernoulli disorder (with $\mu_0(\{w\}) = \frac{1}{2}$) we have $\beta_c^* \in (0, \infty)$ if and only if $\chi \in (1, \infty)$. This shows that the condition $\mu_0(\{w\}) = 0$ is not (!) necessary for $\beta_c < \infty$, i.e., Corollary 1.9 is not optimal.

Remark 1.19. As shown in Doney [9], subject to the condition of regular variation in (1.24),

$$\mathbf{P}(S_n = 0) \sim \frac{C_\alpha}{n^{1-\alpha}L(n)} \quad \text{as } n \rightarrow \infty \text{ with } C_\alpha = (\alpha/\pi) \sin(\alpha\pi) \text{ when } \alpha \in (0, 1). \quad (1.27)$$

Hence the condition $\chi < \infty$ in Corollary 1.8 is satisfied exactly for $\alpha \in (0, \frac{1}{2})$ and L arbitrary, and for $\alpha = \frac{1}{2}$ and $\sum_{n \in \mathbb{N}} n^{-1}[L(n)]^{-2} < \infty$. This fits precisely with cases (3) and (4) in Theorem 1.11.

Remark 1.20. Corollary 1.7(ii) is essentially Corollary 3.2 in Toninelli [22], where the condition for relevance, $h(\mu_\beta | \mu_0) > h(K)$, is given in an equivalent form (see Equation (3.6) in [22]). Note that, by (1.2), $h(K) < \infty$ when $\alpha \in (0, \infty)$.

1.4 Outline

In Section 2 we formulate the annealed and the quenched large deviation principle (LDP) that are in Birkner, Greven and den Hollander [6], which are the key tools in the present paper. In Section 3 we use these LDP's to prove Theorem 1.3. In Section 4 we compare the variational formulas for the two critical curves and prove the criterion for disorder relevance stated in Theorem 1.4. In Section 5 we reformulate this criterion to put it into a form that is more convenient for computations. In Section 6 we use the latter to prove Theorem 1.5. In Sections 7–8 we prove Corollaries 1.6–1.9. Appendix A collects a few facts about the homopolymer.

2 Annealed and quenched LDP

In this section we recall the main results from Birkner, Greven and den Hollander [6] that are needed in the present paper. Section 2.1 introduces the relevant notation, while Sections 2.2 and 2.3 state the relevant annealed and quenched LDP's.

2.1 Notation

Let E be a Polish space, playing the role of an alphabet, i.e., a set of *letters*. Let $\tilde{E} := \cup_{k \in \mathbb{N}} E^k$ be the set of *finite words* drawn from E , which can be metrized to become a Polish space.




Figure 4: Cutting words out from a sequence of letters according to renewal times.

Fix $\mu_0 \in \mathcal{P}(E)$, and $K \in \mathcal{P}(\mathbb{N})$ satisfying (1.2). Let $X = (X_k)_{k \in \mathbb{N}_0}$ be i.i.d. E -valued random variables with marginal law μ_0 , and $\tau = (\tau_i)_{i \in \mathbb{N}}$ i.i.d. \mathbb{N} -valued random variables with marginal law K . Assume that X and τ are independent, and write P^* to denote their joint law. Cut words out of the letter sequence X according to τ (see Fig. 4), i.e., put

$$T_0 := 0 \quad \text{and} \quad T_i := T_{i-1} + \tau_i, \quad i \in \mathbb{N}, \quad (2.1)$$

and let

$$Y^{(i)} := (X_{T_{i-1}}, X_{T_{i-1}+1}, \dots, X_{T_i-1}), \quad i \in \mathbb{N}. \quad (2.2)$$

Under the law P^* , $Y = (Y^{(i)})_{i \in \mathbb{N}}$ is an i.i.d. sequence of words with marginal distribution q_0 on \tilde{E} given by

$$\begin{aligned} q_0((x_1, \dots, x_n)) &:= P^*(Y^{(1)} = (x_1, \dots, x_n)) \\ &= K(n) \mu_0(x_1) \times \dots \times \mu_0(x_n), \quad n \in \mathbb{N}, x_1, \dots, x_n \in E. \end{aligned} \quad (2.3)$$

The reverse operation of *cutting* words out of a sequence of letters is *glueing* words together into a sequence of letters. Formally, this is done by defining a *concatenation* map κ from $\tilde{E}^{\mathbb{N}}$ to $E^{\mathbb{N}_0}$. This map induces in a natural way a map from $\mathcal{P}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}(E^{\mathbb{N}_0})$, the sets of probability measures on $\tilde{E}^{\mathbb{N}}$ and $E^{\mathbb{N}_0}$ (endowed with the topology of weak convergence). The concatenation $q_0^{\otimes \mathbb{N}} \circ \kappa^{-1}$ of $q_0^{\otimes \mathbb{N}}$ equals $\mu_0^{\mathbb{N}_0}$, as is evident from (2.3)

2.2 Annealed LDP

Let $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ be the set of probability measures on $\tilde{E}^{\mathbb{N}}$ that are invariant under the left-shift $\tilde{\theta}$ acting on $\tilde{E}^{\mathbb{N}}$. For $N \in \mathbb{N}$, let $(Y^{(1)}, \dots, Y^{(N)})^{\text{per}}$ be the periodic extension of the N -tuple $(Y^{(1)}, \dots, Y^{(N)}) \in \tilde{E}^{\mathbb{N}}$ to an element of $\tilde{E}^{\mathbb{N}}$, and define

$$R_N := \frac{1}{N} \sum_{i=0}^{N-1} \delta_{\tilde{\theta}^i(Y^{(1)}, \dots, Y^{(N)})^{\text{per}}} \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}). \quad (2.4)$$

This is the *empirical process of N -tuples of words*. The following *annealed LDP* is standard (see e.g. Dembo and Zeitouni [7], Section 6.5). For $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, let $H(Q | q_0^{\otimes \mathbb{N}})$ be the *specific relative entropy of Q w.r.t. $q_0^{\otimes \mathbb{N}}$* defined by

$$H(Q | q_0^{\otimes \mathbb{N}}) := \lim_{N \rightarrow \infty} \frac{1}{N} h(\pi_N Q | \pi_N q_0^{\otimes \mathbb{N}}), \quad (2.5)$$

where $\pi_N Q \in \mathcal{P}(\tilde{E}^N)$ denotes the projection of Q onto the first N words, $h(\cdot | \cdot)$ denotes relative entropy, and the limit is non-decreasing.

Theorem 2.1. *The family $P^*(R_N \in \cdot)$, $N \in \mathbb{N}$, satisfies the LDP on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ with rate N and with rate function I^{ann} given by*

$$I^{\text{ann}}(Q) := H(Q | q_0^{\otimes \mathbb{N}}), \quad Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}). \quad (2.6)$$

This rate function is lower semi-continuous, has compact level sets, has a unique zero at $q_0^{\otimes \mathbb{N}}$, and is affine.

2.3 Quenched LDP

To formulate the quenched analogue of Theorem 2.1, we need some more notation. Let $\mathcal{P}^{\text{inv}}(E^{\mathbb{N}_0})$ be the set of probability measures on $E^{\mathbb{N}_0}$ that are invariant under the left-shift θ acting on $E^{\mathbb{N}_0}$. For $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ such that $m_Q := E_Q(\tau_1) < \infty$ (where E_Q denotes expectation under the law Q and τ_1 is the length of the first word), define

$$\Psi_Q := \frac{1}{m_Q} E_Q \left(\sum_{k=0}^{\tau_1-1} \delta_{\theta^k \kappa(Y)} \right) \in \mathcal{P}^{\text{inv}}(E^{\mathbb{N}_0}). \quad (2.7)$$

Think of Ψ_Q as the shift-invariant version of $Q \circ \kappa^{-1}$ obtained after *randomizing* the location of the origin. This randomization is necessary because a shift-invariant Q in general does not give rise to a shift-invariant $Q \circ \kappa^{-1}$.

For $\text{tr} \in \mathbb{N}$, let $[\cdot]_{\text{tr}}: \tilde{E} \rightarrow [\tilde{E}]_{\text{tr}} = \cup_{n=1}^{\text{tr}} E^n$ denote the *truncation map* on words defined by

$$y = (x_1, \dots, x_n) \mapsto [y]_{\text{tr}} := (x_1, \dots, x_{n \wedge \text{tr}}), \quad n \in \mathbb{N}, x_1, \dots, x_n \in E, \quad (2.8)$$

i.e., $[y]_{\text{tr}}$ is the word of length $\leq \text{tr}$ obtained from the word y by dropping all the letters with label $> \text{tr}$. This map induces in a natural way a map from $\tilde{E}^{\mathbb{N}}$ to $[\tilde{E}]_{\text{tr}}^{\mathbb{N}}$, and from $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv}}([\tilde{E}]_{\text{tr}}^{\mathbb{N}})$. Note that if $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, then $[Q]_{\text{tr}}$ is an element of the set

$$\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}}) = \{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : m_Q < \infty\}. \quad (2.9)$$

Theorem 2.2. (Birkner, Greven and den Hollander [6]) *Assume (1.2). Then, for $\mu_0^{\otimes \mathbb{N}_0}$ -a.s. all X , the family of (regular) conditional probability distributions $P^*(R_N \in \cdot | X)$, $N \in \mathbb{N}$, satisfies the LDP on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ with rate N and with deterministic rate function I^{que} given by*

$$I^{\text{que}}(Q) := \begin{cases} I^{\text{fin}}(Q), & \text{if } Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}}), \\ \lim_{\text{tr} \rightarrow \infty} I^{\text{fin}}([Q]_{\text{tr}}), & \text{otherwise,} \end{cases} \quad (2.10)$$

where

$$I^{\text{fin}}(Q) := H(Q | q_0^{\otimes \mathbb{N}}) + \alpha m_Q H(\Psi_Q | \mu_0^{\otimes \mathbb{N}_0}). \quad (2.11)$$

This rate function is lower semi-continuous, has compact level sets, has a unique zero at $q_0^{\otimes \mathbb{N}}$, and is affine.

There is no closed form expression for $I^{\text{que}}(Q)$ when $m_Q = \infty$. For later reference we remark that, for all $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$,

$$\begin{aligned} I^{\text{ann}}(Q) &= \lim_{\text{tr} \rightarrow \infty} I^{\text{ann}}([Q]_{\text{tr}}) = \sup_{\text{tr} \in \mathbb{N}} I^{\text{ann}}([Q]_{\text{tr}}), \\ I^{\text{que}}(Q) &= \lim_{\text{tr} \rightarrow \infty} I^{\text{que}}([Q]_{\text{tr}}) = \sup_{\text{tr} \in \mathbb{N}} I^{\text{que}}([Q]_{\text{tr}}), \end{aligned} \quad (2.12)$$

as shown in [6], Lemma A.1. A remarkable aspect of (2.11) in relation to (2.6) is that it *quantifies* the difference between I^{que} and I^{ann} . Note the explicit appearance of the tail exponent α . Also note that $I^{\text{que}} = I^{\text{ann}}$ when $\alpha = 0$.

3 Variational formulas: Proof of Theorem 1.3

In Section 3.1 we prove (1.16), the variational formula for the annealed critical curve. The proof of (1.15) in Sections 3.2–3.4, the variational formula for the quenched critical curve, is longer. In Section 3.2 we first give the proof for μ_0 with finite support. In Section 3.3 we extend the proof to μ_0 satisfying (1.3). In Section 3.4 we prove three technical lemmas that are needed in Section 3.3.

3.1 Proof of (1.16)

Proof. Recall from (1.17–1.18) that $Q_\beta = q_\beta^{\otimes \mathbb{N}}$, and from (1.11) that $h_c^{\text{ann}}(\beta) = \log M(\beta)$. Below we show that for every $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$,

$$\beta \Phi(Q) - I^{\text{ann}}(Q) = \log M(\beta) - H(Q | Q_\beta). \quad (3.1)$$

Taking the supremum over Q , we arrive at (1.16). Note that the unique probability measure that achieves the supremum in (3.1) is Q_β , which is an element of the set \mathcal{C} defined in (1.13) because of (1.3).

To get (3.1), note that $H(Q|Q_\beta)$ is the limit as $N \rightarrow \infty$ of (recall (1.17–1.18))

$$\begin{aligned} & \frac{1}{N} \int_{\tilde{E}^N} \log \left[\frac{d\pi_N Q}{d\pi_N Q_\beta} (y_1, \dots, y_N) \right] d\pi_N Q(y_1, \dots, y_N) \\ &= \frac{1}{N} \int_{\tilde{E}^N} \log \left[\frac{d\pi_N Q}{d\pi_N Q_0} (y_1, \dots, y_N) \frac{M(\beta)^N}{e^{\beta[c(y_1) + \dots + c(y_N)]}} \right] d\pi_N Q(y_1, \dots, y_N) \\ &= \log M(\beta) + \frac{1}{N} h(\pi_N Q | \pi_N Q_0) - \beta \frac{1}{N} \int_{\tilde{E}^N} [c(y_1) + \dots + c(y_N)] d\pi_N Q(y_1, \dots, y_N), \end{aligned} \quad (3.2)$$

where, $c(y)$ denotes the first letter of the word y . In the last line of (3.2), the limit as $N \rightarrow \infty$ of the second quantity is $H(Q|Q_0) = I^{\text{ann}}(Q)$, while the integral equals $N\Phi(Q)$ by shift-invariance of Q . Thus, (3.1) follows. \blacksquare

3.2 Proof of (1.15) for μ_0 with finite support

Proof. The proof comes in three steps.

Step 1: An alternative way to compute the quenched free energy $f^{\text{que}}(\beta, h)$ from (1.5) is through the radius of convergence $z^{\text{que}}(\beta, h)$ of the power series

$$\sum_{n \in \mathbb{N}} z^n Z_n^{\beta, h, \omega}, \quad (3.3)$$

because

$$z^{\text{que}}(\beta, h) = e^{-f^{\text{que}}(\beta, h)}. \quad (3.4)$$

Write

$$Z_n^{\beta, h, \omega} = \sum_{N \in \mathbb{N}} \sum_{0=k_0 < k_1 < \dots < k_N = n} \prod_{i=1}^N K(k_i - k_{i-1}) e^{\beta \omega_{k_{i-1}} - h}, \quad (3.5)$$

so that, for $z \in (0, 1]$,

$$\sum_{n \in \mathbb{N}} z^n Z_n^{\beta, h, \omega} = \sum_{N \in \mathbb{N}} F_N^{\beta, h, \omega}(z), \quad (3.6)$$

where we abbreviate

$$F_N^{\beta, h, \omega}(z) := \sum_{0=k_0 < \dots < k_N < \infty} \prod_{i=1}^N z^{k_i - k_{i-1}} K(k_i - k_{i-1}) e^{\beta \omega_{k_{i-1}} - h}. \quad (3.7)$$

Step 2: Return to the setting of Section 2. The letter space is E , the word space is $\tilde{E} = \cup_{k \in \mathbb{N}} E^k$, the sequence of letters is $\omega = (\omega_k)_{k \in \mathbb{N}_0}$, while the sequence of renewal times is $(T_i)_{i \in \mathbb{N}} = (k_i)_{i \in \mathbb{N}}$. Each interval $I_i := [k_{i-1}, k_i)$ of integers cuts out a word $\omega_{I_i} := (\omega_{k_{i-1}}, \dots, \omega_{k_i-1})$. Let

$$R_N^\omega = R_N^\omega((k_i)_{i=0}^N) := \frac{1}{N} \sum_{i=0}^{N-1} \delta_{\tilde{\theta}^i(\omega_{I_1}, \dots, \omega_{I_N})}^{\text{per}} \quad (3.8)$$

denote the empirical process of N -tuples of words in ω cut out by the first N renewals. Then we can rewrite $F_N^{\beta,h,\omega}(z)$ as

$$\begin{aligned} F_N^{\beta,h,\omega}(z) &= \mathbf{E} \left(\exp \left[N \int_{\tilde{E}} \{ \tau(y) \log z + (\beta c(y) - h) \} (\pi_1 R_N^\omega)(dy) \right] \right) \\ &= e^{-Nh} \mathbf{E} \left(\exp \left[Nm_{R_N^\omega} \log z + N\beta\Phi(R_N^\omega) \right] \right), \end{aligned} \quad (3.9)$$

where $\tau(y)$ and $c(y)$ are the length, respectively, the first letter of the word y , $\pi_1 R_N^\omega$ is the projection of R_N^ω onto the first word, while $m_{R_N^\omega}$ and $\Phi(R_N^\omega)$ are the average word length, respectively, the average first letter under R_N^ω .

To identify the radius of convergence of the series in the l.h.s. of (3.6), we apply the root test for the series in the r.h.s. of (3.6) using the expression in (3.9). To that end, let

$$S^{\text{que}}(\beta; z) := \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbf{E} \left(\exp \left[Nm_{R_N^\omega} \log z + N\beta\Phi(R_N^\omega) \right] \right). \quad (3.10)$$

Then

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log F_N^{\beta,h,\omega}(z) = -h + S^{\text{que}}(\beta; z). \quad (3.11)$$

We know from (3.4) and the nonnegativity of $f^{\text{que}}(\beta, h)$ that $z^{\text{que}}(\beta, h) \leq 1$, and we are interested in knowing when it is < 1 , respectively, $= 1$ (recall (1.6)). Hence, the sign of the r.h.s. of (3.11) for $z \uparrow 1$ will be important as the next lemma shows.

Lemma 3.1. *For all $\beta \in [0, \infty)$ and $h \in \mathbb{R}$,*

$$\begin{aligned} S^{\text{que}}(\beta; 1-) < h &\implies f(\beta, h) = 0, \\ S^{\text{que}}(\beta; 1-) > h &\implies f(\beta, h) > 0. \end{aligned} \quad (3.12)$$

Proof. The first line holds because, by (3.11), $-h + S^{\text{que}}(\beta; 1-) < 0$ implies that the sums in (3.6) converge for $|z| < 1$, so that $z^{\text{que}}(\beta, h) \geq 1$, which gives $f^{\text{que}}(\beta, h) \leq 0$. The second line holds because if $-h + S^{\text{que}}(\beta; 1-) > 0$, then there exists a $z_0 < 1$ such that $-h + S^{\text{que}}(\beta; z_0) > 0$, which implies that the sums in (3.6) diverge for $z = z_0$, so that $z^{\text{que}}(\beta, h) \leq z_0 < 1$, which gives $f^{\text{que}}(\beta, h) > 0$. ■

Lemma 3.1 implies that

$$h_c^{\text{que}}(\beta) = S^{\text{que}}(\beta; 1-). \quad (3.13)$$

The rest of the proof is devoted to computing $S^{\text{que}}(\beta; 1-)$.




Figure 5: Qualitative plot of $z \mapsto S^{\text{que}}(\beta; z)$.

Step 3: To the expression in (3.10), defining $S^{\text{que}}(\beta; 1)$, we can apply Varadhan's lemma (because μ_0 has finite support), using the LDP of Theorem 2.2, and conclude that

$$S^{\text{que}}(\beta; 1) = \sup_{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})} [\beta \Phi(Q) - I^{\text{que}}(Q)]. \quad (3.14)$$

We would like to do the same for (3.10) with $z < 1$, and subsequently take the limit $z \uparrow 1$, to get (see Fig. 5))

$$S^{\text{que}}(\beta; 1-) = \sup_{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})} [\beta \Phi(Q) - I^{\text{que}}(Q)]. \quad (3.15)$$

However, even though $Q \mapsto \Phi(Q)$ is continuous (because μ_0 has finite support), $Q \mapsto m_Q$ is only lower semicontinuous. Therefore we proceed by first showing that the term $Nm_{R_N^\omega} \log z$ in (3.10) is harmless in the limit as $z \uparrow 1$.

Lemma 3.2. $S^{\text{que}}(\beta; 1-) = S^{\text{que}}(\beta; 1)$ for all $\beta \in [0, \infty)$.

Proof. Since $S^{\text{que}}(\beta; 1-) \leq S^{\text{que}}(\beta; 1)$, we need only prove the reverse inequality. The idea is to show that, for any $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ and in the limit as $N \rightarrow \infty$, R_N^ω can be arbitrarily close to Q with probability $\approx \exp[-NI^{\text{que}}(Q)]$ while $m_{R_N^\omega}$ remains bounded by a large constant. Therefore, letting $N \rightarrow \infty$ followed by $z \uparrow 1$, we can remove the term $Nm_{R_N^\omega} \log z$ in (3.10).

In what follows, we borrow an idea from the proof of Theorem 2.2 in Birkner, Greven and den Hollander [6], Section 4. Fix $A < S^{\text{que}}(\beta; 1)$. By (3.14) and (2.12), there is a $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ with $m_Q < \infty$ such that $\beta \Phi(Q) - I^{\text{que}}(Q) > A$. Because Φ and I^{que} are affine, we may assume without loss of generality that Q is ergodic.

For $\varepsilon > 0$, the set

$$\mathcal{U}_\varepsilon(Q) := \{Q' \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : \Phi(Q') > \Phi(Q) - \varepsilon\} \quad (3.16)$$

is open because Φ is continuous. Proposition 4.1 in [6] gives a lower bound on the probability that $R_N^\omega \in \mathcal{U}_\varepsilon(Q)$. For the case where Q^* is ergodic, as here, the proof constructs a set of renewal paths with N renewals for which $R_N^\omega \in \mathcal{U}_\varepsilon(Q)$. This set is chosen such that, for M large, there are $\lceil N/(M+1) \rceil \ll N$ “long” renewals, which are used to reach stretches in ω of length $\approx Mm_Q$ that look typical for Ψ_Q , and $\lfloor NM/(M+1) \rfloor \approx N$ “short” renewals that look typical for Q , which come in blocks of M consecutive short renewals in between the long renewals and are used to arrange that $R_N^\omega \approx Q$ for large M . The renewal times $0 < j_1 < \dots < j_N < \infty$ are specified after Equation (4.6) in [6], with a minor adaptation: $j_1 = \sigma_1^{(M)}(\omega)$ is the first large renewal time, $j_2 - j_1, \dots, j_{M+1} - j_M$ are the word lengths corresponding to the z_a 's mentioned below Equation (4.2) in [6], $j_{M+2} = \sigma_2^{(M)}(\omega)$ is the second large renewal time, etc. The set of renewal paths is used to show that

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbf{P}(R_N^\omega \in \mathcal{U}_\varepsilon(Q)) \geq -I^{\text{fin}}(Q) - 6\varepsilon, \quad (3.17)$$

which is Equation (4.8) in [6]. Of course, $I^{\text{fin}}(Q) = I^{\text{que}}(Q)$ because $m_Q < \infty$.

Here we also want to control the average length $m_{R_N^\omega}$ of the N renewals. To that end, let $p_B(Q, \varepsilon, M)$ be the probability in the left-hand side of Equation (4.5) in [6]. This quantity depends on Q , ε and M , but not on N . We have

$$Nm_{R_N^\omega} \leq \sigma_{\lceil N/(M+1) \rceil + 1}^{(M)}(\omega) \quad (3.18)$$

and

$$\limsup_{N \rightarrow \infty} \frac{M+1}{N} \sigma_{\lfloor N/(M+1) \rfloor + 1}^{(M)}(\omega) \leq \frac{1}{p_{\mathcal{B}}(Q, \varepsilon, M)} + M[m_Q + \varepsilon] \quad \omega - a.s. \quad (3.19)$$

The latter inequality follows from the definition of the long renewals and the ergodic theorem. Hence, by (3.10), and restricting the expectation to the set of paths described above, we get

$$\begin{aligned} S^{\text{que}}(\beta; z) &= \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbf{E} \left(\exp \left[Nm_{R_N^\omega} \log z + N\beta\Phi(R_N^\omega) \right] \right) \\ &\geq \frac{1}{M+1} \left(\frac{1}{p_{\mathcal{B}}(Q, \varepsilon, M)} + M[m_Q + \varepsilon] \right) \log z + \beta[\Phi(Q) - \varepsilon] - I^{\text{que}}(Q) - 6\varepsilon. \end{aligned} \quad (3.20)$$

Now let $z \uparrow 1$ and $\varepsilon \downarrow 0$, to get $S^{\text{que}}(\beta; 1-) \geq \beta\Phi(Q) - I^{\text{que}}(Q) > A$. Since $A < S^{\text{que}}(\beta; 1)$ was arbitrary, it follows that $S^{\text{que}}(\beta; 1-) \geq S^{\text{que}}(\beta; 1)$, as claimed. \blacksquare

Combining Lemma 3.2 with (3.13) and (3.14), we obtain (1.15). \blacksquare

3.3 Proof of (1.15) for μ_0 satisfying (1.3)

The proof stays the same up to (3.13). Henceforth write $\mathcal{C} = \mathcal{C}(\mu_0)$ to exhibit the fact that the set \mathcal{C} in (1.13) depends on μ_0 via its support E in (1.12), and define

$$A(\beta) := \sup_{Q \in \mathcal{C}(\mu_0)} [\beta\Phi(Q) - I^{\text{que}}(Q)], \quad (3.21)$$

which replaces the right-hand side of (3.15). We will show the following.

Lemma 3.3. $S^{\text{que}}(\beta; 1-) = A(\beta)$ for all $\beta \in (0, \infty)$.

Proof. The proof of the lemma is accomplished in four steps. Along the way we use three technical lemmas, the proof of which is deferred to Section 3.4. Our starting point is the validity of the claim for μ_0 with finite support obtained in Lemma 3.2. (Note that $|E| < \infty$ implies $\mathcal{C} = \mathcal{C}(\mu_0) = \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$.)

Step 1: $S^{\text{que}}(\beta; 1-) \leq A(\beta)$ for all $\beta \in (0, \infty)$ when μ_0 satisfies (1.3).

Proof. We have $S^{\text{que}}(\beta; 1-) \leq S^{\text{que}}(\beta; 1)$. We will show that $S^{\text{que}}(\beta; 1) \leq A(p\beta)/p$ for all $p > 1$. Taking $p \downarrow 1$ and using the continuity of A , proven in Lemma 3.4 below, we get the claim.

For $M > 0$, let

$$\Phi^M(Q) := \int_E (x \wedge M) d(\pi_{1,1}Q)(x). \quad (3.22)$$

Then, for any $p, q > 1$ such that $p^{-1} + q^{-1} = 1$, we have

$$\begin{aligned} \mathbf{E} \left(e^{N\beta\Phi(R_N^\omega)} \right) &= \mathbf{E} \left(e^{\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) \leq M\}}} e^{\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) > M\}}} \right) \\ &\leq \left[\mathbf{E} \left(e^{p\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) \leq M\}}} \right) \right]^{1/p} \left[\mathbf{E} \left(e^{q\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) > M\}}} \right) \right]^{1/q} \\ &\leq \left[\mathbf{E} \left(e^{Np\beta\Phi^M(R_N^\omega)} \right) \right]^{1/p} \left[\mathbf{E} \left(e^{q\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) > M\}}} \right) \right]^{1/q} \end{aligned} \quad (3.23)$$

and hence

$$\frac{1}{N} \log \mathbf{E} \left(e^{N\beta\Phi(R_N^\omega)} \right) \leq \frac{1}{p} \frac{1}{N} \log \mathbf{E} \left(e^{Np\beta\Phi^M(R_N^\omega)} \right) + \frac{1}{q} \frac{1}{N} \log \mathbf{E} \left(e^{q\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) > M\}}} \right). \quad (3.24)$$

Since $Q \mapsto \Phi^M(Q)$ is upper semicontinuous, Varadhan's lemma gives

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbf{E} \left(e^{Np\beta\Phi^M(R_N^\omega)} \right) \leq \sup_{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})} [p\beta\Phi^M(Q) - I^{\text{que}}(Q)]. \quad (3.25)$$

Clearly, Q 's with $\int_E (x \wedge 0) d(\pi_{1,1}Q)(x) = -\infty$ do not contribute to the supremum. Also, Q 's with $\int_E (x \vee 0) d(\pi_{1,1}Q)(x) = \infty$ do not contribute, because for such Q we have $I^{\text{que}}(Q) = \infty$, by Lemma 3.5 below, and $\Phi^M(Q) < \infty$. Since $\Phi^M \leq \Phi$, we therefore have

$$\sup_{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})} [p\beta\Phi^M(Q) - I^{\text{que}}(Q)] \leq \sup_{Q \in \mathcal{C}(\mu_0)} [p\beta\Phi(Q) - I^{\text{que}}(Q)] = A(p\beta). \quad (3.26)$$

Next, we use the following observation. For any sequence $\Theta = (\Theta_N)_{N \in \mathbb{N}}$ of positive random variables we have

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \Theta_N \leq \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{E}(\Theta_N) \quad \Theta - a.s., \quad (3.27)$$

by the first Borel-Cantelli lemma. Applying this to

$$\Theta_N := \mathbf{E} \left(e^{q\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) > M\}}} \right) \quad \text{with} \quad \mathbb{E}(\Theta_N) = \left(\int_E e^{q\beta x 1_{\{x > M\}}} d\mu_0(x) \right)^N =: (c_M)^N, \quad (3.28)$$

we get, after letting $N \rightarrow \infty$ in (3.24),

$$S^{\text{que}}(\beta; 1) \leq \frac{1}{p} A(p\beta) + \frac{1}{q} \log c_M. \quad (3.29)$$

By (1.3), we have $c_M < \infty$ for all $M > 0$ and $\lim_{M \rightarrow \infty} c_M = 1$. Hence $S^{\text{que}}(\beta; 1) \leq A(p\beta)/p$. ■

Step 2: $S^{\text{que}}(\beta; 1-) \geq A(\beta)$ for all $\beta \in (0, \infty)$ when μ_0 has bounded support.

Proof. In the estimates below, we abbreviate

$$L_N^\omega := NmR_N^\omega, \quad (3.30)$$

the sum of the lengths of the first N words. The proof is based on a discretization argument similar to the one used in [6], Section 8. For $\delta > 0$ and $x \in E$, let $\langle x \rangle_\delta := \sup\{k\delta : k \in \mathbb{Z}, k\delta \leq x\}$. The operation $\langle \cdot \rangle$ extends to measures on E , \tilde{E} and $\tilde{E}^{\mathbb{N}}$ in the obvious way. Now, $\langle R_N^\omega \rangle_\delta$ satisfies the quenched LDP with rate function I_δ^{que} , the quenched rate function corresponding to the measure $\langle \mu_0 \rangle_\delta$. Clearly,

$$\mathbf{E} \left(e^{L_N^\omega \log z + N\beta\Phi(R_N^\omega)} \right) \geq \mathbf{E} \left(e^{L_N^\omega \log z + N\beta\Phi(\langle R_N^\omega \rangle_\delta)} \right), \quad (3.31)$$

and so, by the results in Section 3.2, we have

$$S^{\text{que}}(\beta; 1-) \geq \sup_{Q \in \mathcal{C}(\langle \mu_0 \rangle_\delta)} [\beta\Phi(Q) - I_\delta^{\text{que}}(Q)]. \quad (3.32)$$

For every $Q \in \mathcal{C}(\mu_0)$, we have

$$\Phi(Q) = \lim_{\delta \downarrow 0} \Phi(\langle Q \rangle_\delta), \quad I^{\text{que}}(Q) = \lim_{n \rightarrow \infty} I_{\delta_n}^{\text{que}}(\langle Q \rangle_{\delta_n}), \quad (3.33)$$

where $\delta_n = 2^{-n}$. The first relation holds because $\Phi(\langle Q \rangle_\delta) \leq \Phi(Q) \leq \Phi(\langle Q \rangle_\delta) + \delta$, the second relation uses Lemma 3.6(i) below. Hence the claim follows by picking $\delta = \delta_n$ in (3.32) and letting $n \rightarrow \infty$. ■

Step 3: $S^{\text{que}}(\beta; 1-) \geq A(\beta)$ for all $\beta \in (0, \infty)$ when μ_0 satisfies (1.3) with support bounded from below.

Proof. For $M > 0$ and $x \in E$, let $x^M = x \wedge M$. This truncation operation acts on μ_0 by moving the mass in (M, ∞) to M , resulting in a measure μ_0^M with bounded support and with associated quenched rate function $I^{\text{que}, M}$. Let $R_N^{\omega, M}$ be the empirical process of N -tuples of words obtained from R_N^ω defined in (2.4) after replacing each letter $x \in E$ by x^M . We have

$$\mathbf{E} \left(e^{L_N^\omega \log z + N\beta\Phi(R_N^\omega)} \right) \geq \mathbf{E} \left(e^{L_N^\omega \log z + N\beta\Phi(R_N^{\omega, M})} \right). \quad (3.34)$$

Combined with the result in Step 2, this bound implies that

$$S(\beta; 1-) \geq \sup_{Q' \in \mathcal{C}(\mu_0^M)} [\beta\Phi(Q') - I^{\text{que}, M}(Q')]. \quad (3.35)$$

For every $Q \in \mathcal{C}(\mu_0)$, we have

$$\begin{aligned} \Phi(Q) &= \lim_{M \rightarrow \infty} \Phi(Q^M) = \lim_{M \rightarrow \infty} \int_E (x \wedge M) d(\pi_{1,1}Q)(x), \\ I^{\text{que}}(Q) &= \lim_{M \rightarrow \infty} I^{\text{que}, M}(Q^M). \end{aligned} \quad (3.36)$$

The first relation holds by dominated convergence, the second relation uses Lemma 3.6(ii) below. It follows from (3.36) that

$$\limsup_{M \rightarrow \infty} \sup_{Q' \in \mathcal{C}(\mu_0^M)} [\beta\Phi(Q') - I^{\text{que}, M}(Q')] \geq \beta\Phi(Q) - I^{\text{que}}(Q) \quad \forall Q \in \mathcal{C}(\mu_0), \quad (3.37)$$

which combined with (3.35) yields

$$S(\beta; 1-) \geq \beta\Phi(Q) - I^{\text{que}}(Q) \quad \forall Q \in \mathcal{C}(\mu_0). \quad (3.38)$$

Take the supremum over $Q \in \mathcal{C}(\mu_0)$ to get the claim. ■

Step 4: $S^{\text{que}}(\beta; 1-) \geq A(\beta)$ for all $\beta \in (0, \infty)$ when μ_0 satisfies (1.3).

Proof. For $M > 0$ and $x \in E$, let $x^{-M} = x \vee (-M)$. This truncation operation acts on μ_0 by moving the mass in $(-\infty, -M)$ to $-M$, resulting in a measure μ_0^{-M} with support bounded from below and with associated quenched rate function $I^{\text{que}, -M}$. Let $R_N^{\omega, -M}$ be the empirical process of N -tuples of words obtained from R_N^ω defined in (2.4) after replacing each letter $x \in E$ by x^{-M} .

As in Step 1, for any $p, q > 1$ such that $p^{-1} + q^{-1} = 1$, we have

$$\begin{aligned} \mathbf{E} \left(e^{L_N^\omega \log z + N\beta\Phi(R_N^{\omega, -M})} \right) &\leq \mathbf{E} \left(e^{L_N^\omega \log z + N\beta\Phi(R_N^\omega)} e^{-\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) < -M\}}} \right) \\ &\leq \left[\mathbf{E} \left(e^{pL_N^\omega \log z + Np\beta\Phi(R_N^\omega)} \right) \right]^{1/p} \left[\mathbf{E} \left(e^{-q\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) < -M\}}} \right) \right]^{1/q}, \end{aligned} \quad (3.39)$$

and hence

$$\begin{aligned} \frac{1}{N} \log \mathbf{E} \left(e^{L_N^\omega \log z + N\beta\Phi(R_N^{\omega, -M})} \right) \\ \leq \frac{1}{p} \frac{1}{N} \log \mathbf{E} \left(e^{pL_N^\omega \log z + Np\beta\Phi(R_N^\omega)} \right) + \frac{1}{q} \frac{1}{N} \log \mathbf{E} \left(e^{-q\beta \sum_{i=1}^N c(y_i) 1_{\{c(y_i) < -M\}}} \right). \end{aligned} \quad (3.40)$$

Let $N \rightarrow \infty$ followed by $z \uparrow 1$. For the l.h.s. we have the lower bound in Step 3, while the second term in the r.h.s. can be handled as in (3.27–3.29). Therefore, recalling (3.10) and writing $p \log z = \log z^p$, we get

$$\begin{aligned} \sup_{Q \in \mathcal{C}(\mu_0^{-M})} [\beta\Phi(Q) - I^{\text{que}, -M}(Q)] &\leq \frac{1}{p} S^{\text{que}}(p\beta; 1-) + \frac{1}{q} \log C_{-M} \\ \text{with } C_{-M} &:= \int_E e^{-q\beta x 1_{\{x < -M\}}} d\mu_0(x). \end{aligned} \quad (3.41)$$

Letting $M \rightarrow \infty$ and using that $\lim_{M \rightarrow \infty} C_{-M} = 1$ by (1.3), we arrive at

$$\frac{1}{p} S^{\text{que}}(p\beta, 1-) \geq \limsup_{M \rightarrow \infty} \sup_{Q \in \mathcal{C}(\mu_0^{-M})} [\beta\Phi(Q) - I^{\text{que}, -M}(Q)] \geq A(\beta), \quad (3.42)$$

where the last inequality is obtained via arguments similar to those following (3.35), which requires the use of Lemma 3.6(iii) below. Finally, let $p \downarrow 1$ and use the continuity of $\beta \mapsto S(\beta; 1-)$, proven in Lemma 3.4 below. ■

This completes the proof of Lemma 3.3, and hence of Theorem 1.3. ■

3.4 Technical lemmas

In the proof of Lemma 3.3 we used three technical lemmas, which we prove in this section.

Lemma 3.4. $\beta \mapsto A(\beta)$ and $\beta \mapsto S^{\text{que}}(\beta; 1-)$ are finite and convex on $[0, \infty)$ and, consequently, are continuous on $(0, \infty)$.

Proof. For the first function, note that $A(\beta) \leq \sup_{Q \in \mathcal{C}(\mu_0)} [\beta\Phi(Q) - I^{\text{ann}}(Q)] \leq \log M(\beta) < \infty$ by (1.3) and (3.1), and convexity follows from the fact that A is a supremum of linear functions. For the second function, note that $S^{\text{que}}(\beta; 1-) \leq S^{\text{que}}(\beta; 1) = A(\beta)$, and convexity follows from Hölder's inequality. ■

Lemma 3.5. If $\mu, \nu \in P(\mathbb{R})$ satisfy $h(\mu | \nu) < \infty$ and $\int_E e^{\lambda x} d\nu(x) < \infty$ for some $\lambda > 0$, then $\int_E (x \vee 0) d\mu(x) < \infty$.

Proof. The claim follows from the inequality

$$\int_E f \, d\mu \leq h(\mu | \nu) + \log \int_E e^f \, d\nu, \quad (3.43)$$

which is valid for all bounded and measurable f (see Dembo and Zeitouni [7], Lemma 6.2.13) and, by monotone convergence, extends to measurable $f \geq 0$. Pick $f(x) = \lambda(x \vee 0)$, $x \in E$. ■

Lemma 3.6. *For every $Q \in \mathcal{P}^{\text{inv}}(\widetilde{E}^{\mathbb{N}})$,*

- (i) $\lim_{n \rightarrow \infty} I_{\delta_n}^{\text{que}}(\langle Q \rangle_{\delta_n}) = I^{\text{que}}(Q)$ with $\delta_n := 2^{-n}$.
- (ii) $\lim_{M \rightarrow \infty} I^{\text{que}, M}(Q^M) = I^{\text{que}}(Q)$.
- (iii) $\lim_{M \rightarrow \infty} I^{\text{que}, -M}(Q^{-M}) = I^{\text{que}}(Q)$.

Proof. (i) The proof proceeds by choosing an appropriate function $I: [0, 1] \rightarrow \mathbb{R}$ and proving that

$$\begin{aligned} \text{(a)} \quad & I(0) = \lim_{\delta \downarrow 0} I(\delta), \\ \text{(b)} \quad & I(0) \geq I(\delta_1) \geq I(\delta_2) \text{ whenever } \delta_2 = k\delta_1 \in (0, 1) \text{ for some } k \in \mathbb{N}. \end{aligned} \quad (3.44)$$

Recalling (2.10–2.11), we see that we need the following choices for I :

$$\begin{aligned} \text{(1)} \quad I(\delta) &= \begin{cases} N^{-1}h(\langle \pi_N Q \rangle_\delta | \langle \pi_N q_0^{\otimes \mathbb{N}} \rangle_\delta), & \delta > 0, \\ N^{-1}h(\pi_N Q | \pi_N q_0^{\otimes \mathbb{N}}), & \delta = 0, \end{cases} \\ \text{(2)} \quad I(\delta) &= \begin{cases} H(\langle Q \rangle_\delta | \langle q_0^{\otimes \mathbb{N}} \rangle_\delta), & \delta > 0, \\ H(Q | q_0^{\otimes \mathbb{N}}), & \delta = 0, \end{cases} \\ \text{(3)} \quad I(\delta) &= \begin{cases} N^{-1}h(\langle \pi_N \Psi_Q \rangle_\delta | \langle \pi_N \mu_0^{\otimes \mathbb{N}_0} \rangle_\delta), & \delta > 0, \\ N^{-1}h(\pi_N \Psi_Q | \pi_N \mu_0^{\otimes \mathbb{N}_0}), & \delta = 0, \end{cases} \\ \text{(4)} \quad I(\delta) &= \begin{cases} H(\langle \Psi_Q \rangle_\delta | \langle \mu_0^{\otimes \mathbb{N}_0} \rangle_\delta), & \delta > 0, \\ H(\Psi_Q | \mu_0^{\otimes \mathbb{N}_0}), & \delta = 0, \end{cases} \end{aligned} \quad (3.45)$$

with $N \in \mathbb{N}$. It is clear from the definition of specific relative entropy (recall 2.5) that if (a) and (b) hold for the choices (1) and (3), then they also hold for the choices (2) and (4), respectively. We will not actually prove (a) and (b) for the choices (1) and (3), but for the simpler choice

$$I(\delta) = \begin{cases} h(\langle \mu \rangle_\delta | \langle \mu_0 \rangle_\delta), & \delta > 0, \\ h(\mu | \mu_0), & \delta = 0. \end{cases} \quad (3.46)$$

The proof will make it evident how to properly deal with (1) and (3).

Let $B(\mathbb{R})$ be the set of real-valued, bounded and Borel measurable functions on \mathbb{R} and, for $\phi \in B(\mathbb{R})$ and $\delta > 0$, let ϕ_δ be the function defined by $\phi_\delta(x) := \phi(\langle x \rangle_\delta)$. As shown in Dembo and Zeitouni [7], Lemma 6.2.13, we have

$$\begin{aligned} h(\langle \mu \rangle_\delta | \langle \mu_0 \rangle_\delta) &= \sup_{\phi \in B(\mathbb{R})} \left\{ \int_{\mathbb{R}} \phi \, d\langle \mu \rangle_\delta - \log \int_{\mathbb{R}} e^\phi \, d\langle \mu_0 \rangle_\delta \right\} \\ &= \sup_{\phi \in B(\mathbb{R})} \left\{ \int_{\mathbb{R}} \phi_\delta \, d\mu - \log \int_{\mathbb{R}} e^{\phi_\delta} \, d\mu_0 \right\}. \end{aligned} \quad (3.47)$$

From this representation, property (b) follows for the choice in (3.46). Next, fix any $\varepsilon > 0$ and take a ϕ such that $\int_{\mathbb{R}} \phi \, d\mu - \log \int_{\mathbb{R}} e^\phi \, d\mu_0 \geq h(\mu | \mu_0) - \varepsilon$. Then, since ϕ_δ converges pointwise to ϕ as $\delta \downarrow 0$, the bounded convergence theorem together with (3.47) give

$$\liminf_{\delta \downarrow 0} h(\langle \mu \rangle_\delta | \langle \mu_0 \rangle_\delta) \geq h(\mu | \mu_0) - \varepsilon. \quad (3.48)$$

Hence $\liminf_{\delta \downarrow 0} I(\delta) \geq I(0) - \varepsilon$. Since $I(0) \geq I(\delta)$, property (a) follows after letting $\varepsilon \downarrow 0$.

Having thus convinced ourselves that (3.44–3.45) are true, we now know that for any $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ the sequences

$$H(\langle Q \rangle_{\delta_n} \mid \langle q_0^{\otimes \mathbb{N}} \rangle_{\delta_n}), \quad H(\langle \Psi_Q \rangle_{\delta_n} \mid \langle \mu_0^{\otimes \mathbb{N}_0} \rangle_{\delta_n}), \quad n \in \mathbb{N}, \quad (3.49)$$

are increasing and converge to $H(Q \mid q_0^{\otimes \mathbb{N}})$, respectively, $H(\Psi_Q \mid \mu_0^{\otimes \mathbb{N}_0})$. This implies the claim for Q with $m_Q < \infty$ (recall (2.11)). For Q with $m_Q = \infty$ we use that $I^{\text{que}}(Q) = \sup_{\text{tr} \in \mathbb{N}} I([\![Q]\!]_{\text{tr}})$ (recall (2.12)), to conclude that $I_{\delta_n}^{\text{que}}(\langle Q \rangle_{\delta_n})$ is increasing and converges to $I^{\text{que}}(Q)$.

(ii–iii) The proof is similar as for (i). ■

4 Characterization of disorder relevance: Proof of Theorem 1.4

Proof. We will need the following lemma, the proof of which is postponed.

Lemma 4.1. *The supremum $\sup_{Q \in \mathcal{C}} [\beta \Phi(Q) - I^{\text{que}}(Q)]$ is attained for all $\beta \in (0, \infty)$.*

Let Q^* be a measure achieving the supremum in Lemma 4.1. Suppose that $h_c^{\text{que}}(\beta) = h_c^{\text{ann}}(\beta)$. Then

$$\begin{aligned} h_c^{\text{que}}(\beta) &= \beta \Phi(Q^*) - I^{\text{que}}(Q^*) \leq \beta \Phi(Q^*) - I^{\text{ann}}(Q^*) \\ &\leq \beta \Phi(Q_\beta) - I^{\text{ann}}(Q_\beta) = h_c^{\text{ann}}(\beta) = h_c^{\text{que}}(\beta), \end{aligned} \quad (4.1)$$

where the second equality uses that Q_β is the unique measure achieving the supremum in (1.16) (with $I^{\text{ann}}(Q_\beta) < \infty$), as shown by (3.1). It follows that both inequalities in (4.1) are equalities. Consequently, $Q^* = Q_\beta$ and $I^{\text{que}}(Q_\beta) = I^{\text{ann}}(Q_\beta)$.

Conversely, suppose that $I^{\text{que}}(Q_\beta) = I^{\text{ann}}(Q_\beta)$. Then

$$h_c^{\text{que}}(\beta) \geq [\beta \Phi(Q_\beta) - I^{\text{que}}(Q_\beta)] = [\beta \Phi(Q_\beta) - I^{\text{ann}}(Q_\beta)] = h_c^{\text{ann}}(\beta). \quad (4.2)$$

Since $h_c^{\text{que}}(\beta) \leq h_c^{\text{ann}}(\beta)$, this proves that $h_c^{\text{que}}(\beta) = h_c^{\text{ann}}(\beta)$. ■

We now give the proof of Lemma 4.1.

Proof. The proof is accomplished in three steps. The claims in Steps 1 and 2 are obvious when the support of μ_0 is bounded from above, because then Φ is bounded from above and upper semicontinuous. Thus, for these steps we may assume that the support of μ_0 is unbounded from above.

Step 1: The supremum can be restricted to the set $\mathcal{C} \cap \{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : I^{\text{que}}(Q) \leq \gamma\}$ for some $\gamma < \infty$.

Proof. We first prove that

$$\lim_{a \rightarrow \infty} \sup_{\substack{Q \in \mathcal{C} \\ \Phi(Q) = a}} [\beta \Phi(Q) - I^{\text{que}}(Q)] = -\infty. \quad (4.3)$$

To that end we estimate, for $a \in (0, \infty)$,

$$\sup_{\substack{Q \in \mathcal{C} \\ \Phi(Q) = a}} [\beta \Phi(Q) - I^{\text{que}}(Q)] \leq \sup_{\substack{Q \in \mathcal{C} \\ \Phi(Q) = a}} [\beta a - h(\pi_{1,1} Q \mid \mu_0)] = \sup_{\substack{\mu \in \mathcal{P}(E) \\ \int_E |x| d\mu(x) < \infty, \int_E x d\mu(x) = a}} [\beta a - h(\mu \mid \mu_0)], \quad (4.4)$$

where we use that $I^{\text{que}}(Q) \geq I^{\text{ann}}(Q) = H(Q|Q_0) \geq h(\pi_{1,1}Q|\mu_0)$. The last supremum is achieved by a measure μ_λ of the form $d\mu_\lambda(x) = M(\lambda)^{-1}e^{\lambda x}d\mu_0(x)$, $x \in E$, with λ such that $\int_E x d\mu_\lambda(x) = a$ (recall (1.17)). To see why, first note that such a $\lambda = \lambda(a)$ exists because $(\lambda \mapsto \int_E x d\mu_\lambda(x))$ is continuous with value 0 at $\lambda = 0$ and $\lim_{\lambda \rightarrow \infty} \int_E x d\mu_\lambda(x) = \sup[\text{supp}(\mu_0)] = w$, where $w = \infty$ by assumption. Next note that, for any other measure μ with $\int_E x d\mu(x) = a$, we have

$$h(\mu|\mu_\lambda) = h(\mu|\mu_0) - \lambda a + \log M(\lambda) = h(\mu|\mu_0) - h(\mu_\lambda|\mu_0), \quad (4.5)$$

which shows that $h(\mu|\mu_0) \geq h(\mu_\lambda|\mu_0)$ with equality if and only if $\mu = \mu_\lambda$. Consequently,

$$\sup_{\substack{\mu \in \mathcal{P}(E) \\ \int_E |x| d\mu(x) < \infty, \int_E x d\mu(x) = a}} [\beta a - h(\mu|\mu_0)] = \beta \int_E x d\mu_\lambda(x) - h(\mu_\lambda|\mu_0) =: g(\lambda). \quad (4.6)$$

Clearly, $a \rightarrow \infty$ implies $\lambda = \lambda(a) \rightarrow \infty$, and so to prove (4.3) we must show that $\lim_{\lambda \rightarrow \infty} g(\lambda) = -\infty$.

To achieve the latter, note that a lower bound on $h(\mu_\lambda|\mu_0)$ is obtained by applying (3.43) to $f(x) := \bar{\beta}(x \vee 0)$ for some $\bar{\beta} > \beta$. This yields

$$g(\lambda) \leq -(\bar{\beta} - \beta) \int_E x d\mu_\lambda(x) + \log [M(\bar{\beta}) + 1]. \quad (4.7)$$

The integral in the right-hand side tends to infinity as $\lambda \rightarrow \infty$, and so (4.3) indeed follows.

Finally, recall the definition of $A(\beta)$ in (3.21), which is finite because of Lemma 3.4. Then, by (4.3), there is an $a_0 < \infty$ such that

$$\sup_{\substack{Q \in \mathcal{C} \\ \Phi(Q) = a}} [\beta \Phi(Q) - I^{\text{que}}(Q)] \leq A(\beta) - 1 \quad \forall a \geq a_0, \quad (4.8)$$

and so all $Q \in \mathcal{C}$ with $\beta \Phi(Q) - I^{\text{que}}(Q) > A(\beta) - 1$ must satisfy $\Phi(Q) < a_0$ and $I^{\text{que}}(Q) < \beta \Phi(Q) + 1 - A(\beta) \leq \beta a_0 + 1 - A(\beta) =: \gamma$. Consequently, the supremum can be restricted to the set $\mathcal{C} \cap \{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : I^{\text{que}}(Q) \leq \gamma\}$. \blacksquare

Step 2: Φ is upper semicontinuous on $\{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : I^{\text{que}}(Q) \leq \gamma\}$ for every $\gamma > 0$.

Proof. From the definition of Φ and the inequality $h(\pi_{1,1}Q|\mu_0) \leq I^{\text{que}}(Q) \leq \gamma$, it follows that it is enough to show that the map $\mu \mapsto \Psi(\mu) := \int_E (x \vee 0) d\mu(x)$ is upper semicontinuous on $K_\gamma := \{\mu \in \mathcal{P}(E) : h(\mu|\mu_0) \leq \gamma\}$. To do so, let $(\mu^M)_{M \in \mathbb{N}}$ be a sequence in K_γ converging to μ weakly as $M \rightarrow \infty$. Then

$$\Psi(\mu^M) = \int_E [(x \vee 0) \wedge n] d\mu^M(x) + \int_E x 1_{\{x > n\}} d\mu^M(x), \quad (4.9)$$

and so

$$\limsup_{M \rightarrow \infty} \Psi(\mu^M) \leq \int_E [(x \vee 0) \wedge n] d\mu(x) + \sup_{M \in \mathbb{N}} \int_E x 1_{\{x > n\}} d\mu^M(x) \quad \forall n \in \mathbb{N}. \quad (4.10)$$

By the inequality in (3.43), we have

$$\lambda \int_E x 1_{\{x > n\}} d\mu^M(x) \leq h(\mu^M|\mu_0) + \log \int_E e^{\lambda x 1_{\{x > n\}}} d\mu_0(x) \quad \forall M, n \in \mathbb{N}, \lambda > 0, \quad (4.11)$$

and so

$$\sup_{M \in \mathbb{N}} \int_E x 1_{\{x > n\}} d\mu^M(x) \leq \frac{\gamma}{\lambda} + \frac{1}{\lambda} \log \int_E e^{\lambda x 1_{\{x > n\}}} d\mu_0(x). \quad (4.12)$$

By (1.3), the limit as $n \rightarrow \infty$ of the r.h.s. is γ/λ . Since $\lambda > 0$ is arbitrary, we conclude that the limit as $n \rightarrow \infty$ of the left-hand side is zero. Letting $n \rightarrow \infty$ in (4.10) and using monotone convergence, we therefore get $\limsup_{M \rightarrow \infty} \Psi(\mu^M) \leq \Psi(\mu)$, as required. \blacksquare

Step 3: Let $\Gamma(Q) := \beta\Phi(Q) - I^{\text{que}}(Q)$. Then

$$\sup_{Q \in \mathcal{C}} \Gamma(Q) = \sup_{\substack{Q \in \mathcal{C} \\ I^{\text{que}}(Q) \leq \gamma}} \Gamma(Q) \leq \sup_{\substack{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) \\ I^{\text{que}}(Q) \leq \gamma}} \Gamma(Q). \quad (4.13)$$

By Theorem 2.2, I^{que} is lower semicontinuous. Hence, by Step 2, $\beta\Phi - I^{\text{que}}$ is upper semicontinuous on the compact set $\{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : I^{\text{que}}(Q) \leq \gamma\}$, achieving its supremum at some Q^* . Let $\mu^* := \pi_{1,1}Q^*$. Then, by (1.3), the inequality in (3.43) gives

$$\int_E (x \vee 0) d\mu^*(x) \leq \gamma + \log \int_E e^x d\mu_0(x) < \infty \quad (4.14)$$

and, since $\Phi(Q^*) > -\infty$, we also have $\int_E (x \wedge 0) d\mu^*(x) > -\infty$, so that $Q^* \in \mathcal{C}$. Hence

$$\sup_{Q \in \mathcal{C}} \Gamma(Q) = \sup_{\substack{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) \\ I^{\text{que}}(Q) \leq \gamma}} \Gamma(Q) = \Gamma(Q^*), \quad (4.15)$$

which concludes the proof. \blacksquare

5 Reformulation of the criterion for disorder relevance

Note that, by (2.10–2.12), for $\alpha > 0$, the necessary and sufficient condition for relevance, $I^{\text{que}}(Q_\beta) > I^{\text{ann}}(Q_\beta)$, in Theorem 1.4 translates into

$$\lim_{\text{tr} \rightarrow \infty} m_{[Q_\beta]_{\text{tr}}} H(\Psi_{[Q_\beta]_{\text{tr}}} | \mu_0^{\otimes \mathbb{N}_0}) > 0. \quad (5.1)$$

In Lemma 5.3 below, we give two alternative expressions for the specific relative entropy appearing in (5.1). These expressions will be needed in Sections 6 and 7.

I. Asymptotic mean stationarity. In what follows we will make use of the notion of *asymptotic mean stationary* (see Gray [16], Section 1.7). Let A be a topological space and equip $A^{\mathbb{N}_0}$ with the product topology. A measure \mathcal{P} on $A^{\mathbb{N}_0}$ is called *asymptotically mean stationary* if for every Borel measurable $G \subset A^{\mathbb{N}_0}$,

$$\overline{\mathcal{P}}(G) := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathcal{P}(\theta^k G) \quad \text{exists.} \quad (5.2)$$

As in Section 2, θ denotes the left-shift acting on $A^{\mathbb{N}_0}$. If \mathcal{P} is asymptotically mean stationary, then $\overline{\mathcal{P}}$ is a stationary measure, called the *stationary mean* of \mathcal{P} .

For $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, recall from Section 2.3 that $\kappa(Q) \in \mathcal{P}(E^{\mathbb{N}_0})$ is the probability measure induced by the concatenation map $\kappa: \tilde{E}^{\mathbb{N}} \rightarrow E^{\mathbb{N}_0}$ that glues a sequence of words into a sequence of letters, i.e., $\kappa(Q) = Q \circ \kappa^{-1}$. Our aim is to replace Ψ_Q in (5.1) by $\kappa(Q)$, which is not stationary but more convenient to work with. These two probability measures are related in the following way.

Lemma 5.1. *If $m_Q < \infty$, then $\kappa(Q)$ is asymptotically mean stationary with stationary mean $\overline{\kappa(Q)} = \Psi_Q$.*

Proof. Let $X := \kappa(Y) \in E^{\mathbb{N}_0}$, where Y is distributed according to Q . Let I denote the set of indices $i \in \mathbb{N}_0$ where a new word starts ($0 \in I$). For $i \in \mathbb{N}_0$, let $r_i := \inf\{j \in \mathbb{N} : i - j \in I\}$, i.e., the distance from i to the beginning of the word it belongs to. For $j \in I$, let L^j denote the length of the word that starts at j . Then, for any $G \subset E^{\mathbb{N}_0}$ Borel measurable, we have

$$\sum_{i=0}^{n-1} \kappa(Q)(\theta^i X \in G) = \sum_{i=0}^{n-1} \sum_{k=0}^i Q(\theta^i X \in G, r_i = k) = \sum_{k=0}^{n-1} \sum_{i=k}^{n-1} Q(\theta^i X \in G, r_i = k). \quad (5.3)$$

Next, note that

$$\begin{aligned} Q(\theta^i X \in G, r_i = k) &= Q(\theta^i X \in G, i - k \in I, L^{i-k} > k) \\ &= Q(\theta^i X \in G, L^{i-k} > k \mid i - k \in I) Q(i - k \in I) \\ &= Q(\theta^k X \in G, L^0 > k) Q(i - k \in I). \end{aligned} \quad (5.4)$$

Hence, dividing the sum in (5.3) by n , we get

$$\frac{1}{n} \sum_{i=0}^{n-1} \kappa(Q)(\theta^i X \in G) = \sum_{k=0}^{n-1} Q(\theta^k X \in G, L^0 > k) f_{k,n}, \quad (5.5)$$

where we abbreviate $f_{k,n} := n^{-1} \sum_{j=0}^{n-k-1} Q(j \in I)$. By the renewal theorem, $\lim_{n \rightarrow \infty} f_{k,n} = 1/m_Q$ for k fixed. Since

$$\sum_{k=0}^{\infty} Q(L^0 > k) = m_Q < \infty, \quad (5.6)$$

we can apply the bounded convergence theorem, and conclude that

$$\begin{aligned} \overline{\kappa(Q)}(G) &= \frac{1}{m_Q} \sum_{k=0}^{\infty} Q(\theta^k X \in G, L^0 > k) = \frac{1}{m_Q} \sum_{k=0}^{\infty} \sum_{j=k+1}^{\infty} Q(\theta^k X \in G, L^0 = j) \\ &= \frac{1}{m_Q} \sum_{j=1}^{\infty} \sum_{k=0}^{j-1} Q(\theta^k X \in G, L^0 = j) = \Psi_Q(G). \end{aligned} \quad (5.7)$$

The last equality is simply the definition of Ψ_Q in (2.7). ■

To complement Lemma 5.1, we need the following fact stated in Birkner [5], Remark 5, where ergodicity refers to the left-shifts acting on $\tilde{E}^{\mathbb{N}}$ and $E^{\mathbb{N}}$.

Lemma 5.2. *If $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ is ergodic and $m_Q < \infty$, then $\Psi_Q \in \mathcal{P}^{\text{inv}}(E^{\mathbb{N}})$ is ergodic.*

An asymptotic mean stationary measure can be interchanged with its stationary mean in several situations (see Gray [15], Chapter 6), for example in relative entropy computations, as in Lemma 5.3 below. Before stating this lemma, we use an extension of the notion of specific relative entropy to measures that are not necessarily stationary. More precisely, for two measures \mathcal{P} and \mathcal{Q} on a product space $A^{\mathbb{N}}$, we define the specific relative entropy of \mathcal{P} w.r.t. \mathcal{Q} as

$$\overline{H}(\mathcal{P} \mid \mathcal{Q}) := \limsup_{n \rightarrow \infty} \frac{1}{n} h(\pi_n \mathcal{P} \mid \pi_n \mathcal{Q}), \quad (5.8)$$

where π_n is the projection onto the first n coordinates. For $Q \in \mathcal{P}^{\text{inv}}(\widetilde{E}^{\mathbb{N}})$, we introduce the following Radon-Nikodym derivative:

$$f_n(x) := \frac{d\pi_n \kappa(Q)}{d\mu_0^{\otimes n}}(x), \quad x \in E^{\mathbb{N}_0}. \quad (5.9)$$

With this notation, the main result of this section is the following.

Lemma 5.3. *For $Q \in \mathcal{P}^{\text{inv}}(\widetilde{E}^{\mathbb{N}})$ ergodic with $m_Q < \infty$,*

$$H(\Psi_Q | \mu_0^{\otimes \mathbb{N}_0}) = \overline{H}(\kappa(Q) | \mu_0^{\otimes \mathbb{N}_0}), \quad (5.10)$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \log f_n(x) \quad \text{for } \kappa(Q)\text{-a.s. all } x \in E^{\mathbb{N}_0}. \quad (5.11)$$

The first equality holds also without the assumption of ergodicity.

Proof. The first equality follows from Gray [16], Corollary 7.5.1, last equality in Eq. (7.32), which does not need the assumption of ergodicity. For the proof of the other equality, define

$$\bar{f}_n(x) := \frac{d\pi_n \Psi_Q}{d\mu_0^{\otimes n}}(x). \quad (5.12)$$

Since Ψ_Q is stationary and ergodic (Lemma 5.2), Gray [16], Theorem 8.2.1, applied to the pair $\Psi_Q, \mu_0^{\otimes \mathbb{N}_0}$ gives that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \bar{f}_n(x) = H(\Psi_Q | \mu_0^{\otimes \mathbb{N}_0}) \quad (5.13)$$

for Ψ_Q almost all x . But Ψ_Q is the stationary mean of $\kappa(Q)$ (Lemma 5.1), so that Gray [16], Theorem 8.4.1, combined with (5.13) gives

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log f_n(x) = H(\Psi_Q | \mu_0^{\otimes \mathbb{N}_0}) \quad (5.14)$$

for $\kappa(Q)$ almost all x . ■

II. Alternative formulation. We will apply Lemma 5.3 to the measure $[Q_\beta]_{\text{tr}}$, which is ergodic, being a product measure. The word length distribution of it is

$$K^{\text{tr}}(n) := \begin{cases} K(n) & \text{if } 1 \leq n \leq \text{tr} - 1, \\ \sum_{m=\text{tr}}^{\infty} K(m) & \text{if } n = \text{tr}, \\ 0 & \text{if } n > \text{tr}. \end{cases} \quad (5.15)$$

For $[Q_\beta]_{\text{tr}}$, the function f_n in (5.9) becomes

$$f_n(x) = \mathbf{E}_{K^{\text{tr}}} \left(\prod_{k=0}^{n-1} \left(\frac{e^{\beta x_k}}{M(\beta)} \right)^{1_{\{S_k=0\}}} \right) = \mathbf{E}_{K^{\text{tr}}} \left(e^{\sum_{k=0}^{n-1} \{\beta x_k - \log M(\beta)\} 1_{\{S_k=0\}}} \right). \quad (5.16)$$

where $\mathbf{E}_{K^{\text{tr}}}$ denotes expectation with respect to law of the Markov chain S with renewal times distribution K^{tr} . This follows from the definition of Q_β and (1.17). To emphasize the fact that in the last expression the sequence $x \in E^{\mathbb{N}_0}$ is picked from $\kappa([Q_\beta]_{\text{tr}})$, we take two independent sequences

$$(x_k)_{k \in \mathbb{N}_0}, (\hat{x}_k)_{k \in \mathbb{N}_0} \text{ drawn from } \mu_0^{\otimes \mathbb{N}_0} \text{ and } \mu_\beta^{\otimes \mathbb{N}_0}, \text{ respectively,} \quad (5.17)$$

and an independent copy S' of S . Let $I := \{i \geq 0 : S_i = 0\}$, $I' := \{i \geq 0 : S'_i = 0\}$. Then

$$H(\Psi_{[Q_\beta]_{\text{tr}}} | \mu_0^{\otimes \mathbb{N}_0}) = \lim_{N \rightarrow \infty} \frac{1}{n} \log \mathbf{E}_{K^{\text{tr}}} \left[e^{\sum_{k=0}^{n-1} [\beta x_k 1_{\{k \notin I'\}} + \beta \hat{x}_k 1_{\{k \in I'\}} - \log M(\beta)] 1_{\{k \in I\}}} \right]. \quad (5.18)$$

This is the key expression for proving relevance or irrelevance. Note the appearance of two renewals set I, I' , which are the key to understanding the issue of relevant vs. irrelevant disorder (recall Remark 1.17).

6 Monotonicity of disorder relevance: Proof of Theorem 1.5

Proof. In view of (5.10) in Lemma 5.3, the condition for relevance in (5.1) becomes

$$\lim_{\text{tr} \rightarrow \infty} m_{[Q_\beta]_{\text{tr}}} \overline{H}(\kappa([Q_\beta]_{\text{tr}}) | \mu_0^{\otimes \mathbb{N}_0}) > 0. \quad (6.1)$$

We will show that $\beta \mapsto \overline{H}(\kappa([Q_\beta]_{\text{tr}}) | \mu_0^{\otimes \mathbb{N}_0})$ is non-decreasing for every $\text{tr} \in \mathbb{N}$, which will imply the claim because $m_{[Q_\beta]_{\text{tr}}} = m_{K^{\text{tr}}}$ does not depend on β . It will be enough to show that $\beta \mapsto h(\pi_n \kappa([Q_\beta]_{\text{tr}}) | \mu_0^{\otimes n})$ is non-decreasing for all $\text{tr}, n \in \mathbb{N}$.

Fix $\text{tr}, n \in \mathbb{N}$. For $\beta \in [0, \infty)$ and $\bar{x} = (x_0, x_1, \dots, x_{n-1}) \in E^n$, let

$$k(\beta, \bar{x}) := \frac{d\pi_n \kappa([Q_\beta]_{\text{tr}})}{d\mu_0^n}(\bar{x}) = \mathbf{E}_{K^{\text{tr}}} \left(\prod_{k \in J_n} \frac{e^{\beta x_k}}{M(\beta)} \right), \quad (6.2)$$

with $J_n := \{0 \leq k < n : S_k = 0\}$ the set of renewal times prior to time n for the chain S that has renewal time distribution K^{tr} , to which we add 0 for convenience. Our goal is to prove that

$$\beta \mapsto f(\beta) := \int_{\mathbb{R}^n} [k(\beta, \bar{x}) \log k(\beta, \bar{x})] d\mu_0^{\otimes n}(\bar{x}) = h(\pi_n \kappa([Q_\beta]_{\text{tr}}) | \mu_0^{\otimes n}) \quad (6.3)$$

is non-decreasing on $[0, \infty)$. We will do this by proving a stronger property. Namely, for $\bar{\beta} = (\beta_0, \beta_1, \dots, \beta_{n-1}) \in [0, \infty)^n$ and $\bar{x} \in E^n$, let

$$k(\bar{\beta}, \bar{x}) := \mathbf{E}_{K^{\text{tr}}} \left(\prod_{k \in J_n} \frac{e^{\beta_k x_k}}{M(\beta_k)} \right). \quad (6.4)$$

We will show that

$$\bar{\beta} \mapsto f(\bar{\beta}) := \int_{\mathbb{R}^n} [k(\bar{\beta}, \bar{x}) \log k(\bar{\beta}, \bar{x})] d\mu_0^{\otimes n}(\bar{x}) \quad (6.5)$$

is non-decreasing on $[0, \infty)^n$ in each of its arguments.

We will prove monotonicity w.r.t. β_1 only. The argument is the same for the other variables, with one simplification for β_0 , namely, we may drop the corresponding indicator $1_{\{0 \in J_n\}}$ in the third line of (6.6) and in (6.8). First, using that $\int k(\bar{\beta}, \bar{x}) d\mu_0^{\otimes n}(\bar{x}) = 1$ for all $\bar{\beta}$, we compute

$$\begin{aligned} \partial_{\beta_1} f(\bar{\beta}) &= \int_{\mathbb{R}^n} \partial_{\beta_1} [k(\bar{\beta}, \bar{x}) \log k(\bar{\beta}, \bar{x})] d\mu_0^{\otimes n}(\bar{x}) \\ &= \int_{\mathbb{R}^n} \partial_{\beta_1} [k(\bar{\beta}, \bar{x})] \log k(\bar{\beta}, \bar{x}) d\mu_0^{\otimes n}(\bar{x}) \\ &= \int_{\mathbb{R}^n} \partial_{\beta_1} \left(\frac{e^{\beta_1 x_1}}{M(\beta_1)} \right) \mathbf{E}_{K^{\text{tr}}} \left(1_{\{1 \in J_n\}} \prod_{k \in J_n \setminus \{1\}} \frac{e^{\beta_k x_k}}{M(\beta_k)} \right) \log k(\bar{\beta}, \bar{x}) d\mu_0^{\otimes n}(\bar{x}). \end{aligned} \quad (6.6)$$

Next, we note that

$$\begin{aligned} \partial_{\beta_1} \left(\frac{e^{\beta_1 x_1}}{M(\beta_1)} \right) d\mu_0(x_1) &= \frac{e^{\beta_1 x_1} x_1 M(\beta_1) - e^{\beta_1 x_1} M'(\beta_1)}{M(\beta_1)^2} d\mu_0(x_1) \\ &= \left(x_1 - \frac{M'(\beta_1)}{M(\beta_1)} \right) \frac{e^{\beta_1 x_1}}{M(\beta_1)} d\mu_0(x_1) = (x_1 - E_{\beta_1}) d\mu_{\beta_1}(x_1), \end{aligned} \quad (6.7)$$

where $E_{\beta_1} := M'(\beta_1)/M(\beta_1) = \int x_1 d\mu_{\beta_1}(x_1)$. Now, let \bar{x}^1 be \bar{x} without x_1 , and abbreviate

$$A(x_1; \bar{x}^1) := \mathbf{E}_{K^{\text{tr}}} \left(\prod_{k \in J_n \setminus \{1\}} \frac{e^{\beta_k x_k}}{M(\beta_k)} 1_{\{1 \in J_n\}} \right) \log k(\bar{\beta}, \bar{x}). \quad (6.8)$$

Then, for fixed \bar{x}^1 , the integral over x_1 in (6.6) equals

$$\begin{aligned} &\int_{\mathbb{R}^n} (x_1 - E_{\beta_1}) A(x_1; \bar{x}^1) d\mu_{\beta_1}(x_1) \\ &\geq \int_{\mathbb{R}^n} (x_1 - E_{\beta_1}) d\mu_{\beta_1}(x_1) \int_{\mathbb{R}^n} A(x_1; \bar{x}^1) d\mu_{\beta_1}(x_1) = 0, \end{aligned} \quad (6.9)$$

where the inequality holds because both $x_1 \mapsto x_1 - E_{\beta_1}$ and $x_1 \mapsto A(x_1; \bar{x}^1)$ are non-decreasing (for the latter we need that $\beta_1 \in [0, \infty)$). It therefore follows from (6.6), after integrating over \bar{x}^1 as well, that $\partial_{\beta_1} f(\bar{\beta}) \geq 0$. \blacksquare

7 Disorder irrelevance: Proof of Corollaries 1.6 and 1.7(i)

7.1 Proof of Corollary 1.6

Proof. This is immediate from Theorem 1.4 and the fact that $I^{\text{que}} = I^{\text{ann}}$ when $\alpha = 0$. The latter was already noted at the end of Section 2. \blacksquare

7.2 Proof of Corollary 1.7(i)

Proof. The claim follows from an annealed bound on $H(\Psi_{[Q_\beta]_{\text{tr}}} | \mu_0^{\otimes \mathbb{N}_0})$. Indeed, recalling (5.11), (5.16), and putting

$$\Theta_n := \mathbf{E}_{K^{\text{tr}}} \left(\prod_{k=0}^{n-1} \left(\frac{e^{\beta x_k}}{M(\beta)} \right)^{1_{\{S_k=0\}}} \right), \quad (7.1)$$

we have

$$\begin{aligned} \mathbf{E}_{\kappa([Q_\beta]_{\text{tr}})}(\Theta_n) &= \mathbf{E}_{K^{\text{tr}}} \left(E_{\kappa([Q_\beta]_{\text{tr}})} \left(\prod_{k=0}^{n-1} \left(\frac{e^{\beta x_k}}{M(\beta)} \right)^{1_{\{S_k=0\}}} \right) \right) \\ &= \mathbf{E}_{S, S'} \left(E_{\mu_0^{\otimes n}} \left(\prod_{k=0}^{n-1} \left(\frac{e^{\beta x_k}}{M(\beta)} \right)^{1_{\{S_k=0\}}} \prod_{l=0}^{n-1} \left(\frac{e^{\beta x_l}}{M(\beta)} \right)^{1_{\{S'_l=0\}}} \right) \right) \\ &= \mathbf{E}_{S, S'} \left(\Xi(\beta)^{\sum_{k=0}^{n-1} 1_{\{S_k=S'_k=0\}}} \right), \end{aligned} \quad (7.2)$$

where $\mathbf{E}_{S,S'}$ is the expectation with respect to two independent copies S, S' of the Markov chain with renewal time distribution K^{tr} , and

$$\Xi(\beta) := \frac{M(2\beta)}{M(\beta)^2}. \quad (7.3)$$

If we now let

$$f_2^{\text{tr}}(\lambda) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbf{E}_{S,S'} \left(e^{\lambda \sum_{k=0}^{n-1} 1_{\{S_k=S'_k=0\}}} \right), \quad (7.4)$$

then (3.27), (5.11), (5.16) and (7.2) imply that

$$H(\Psi_{[Q_\beta]_{\text{tr}}} | \mu_0^{\otimes N_0}) \leq f_2^{\text{tr}}(\log \Xi(\beta)), \quad \beta \in [0, \infty), \text{tr} \in \mathbb{N}. \quad (7.5)$$

Combining the condition for relevance in (5.1) with the bound in (7.5), we see that to prove irrelevance it suffices to show that, under the conditions in Corollary 1.8,

$$\lim_{\text{tr} \rightarrow \infty} m_{[Q_\beta]_{\text{tr}}} f_2^{\text{tr}}(\log \Xi(\beta)) = 0. \quad (7.6)$$

By (A.2) in Appendix A, we have

$$f_2(\lambda) = 0 \iff \lambda \leq \lambda_0 := -\log \mathbf{P}(I \cap I' \neq \emptyset), \quad (7.7)$$

where I, I' are the sets of renewal times for S, S' without truncation, and $f_2(\lambda)$ as defined in Appendix A. By Lemma A.1, if $\lambda < \lambda_0$, then $\sup_{\text{tr} \in \mathbb{N}} \text{tr} f_2^{\text{tr}}(\lambda) < \infty$. Since $\lim_{\text{tr} \rightarrow \infty} m_{[Q_\beta]_{\text{tr}}}/\text{tr} = 0$ always, (7.6) holds as soon as $\log \Xi(\beta) < \lambda_0$, i.e., $\Xi(\beta) < 1/\mathbf{P}(I \cap I' \neq \emptyset)$. Now the claim follows because $\mathbf{P}(I \cap I' \neq \emptyset) = \chi/(\chi + 1)$ (see Spitzer [20], Section 1), with χ as defined in (1.21), and with the convention that the last ratio is 1 if $\chi = \infty$. ■

8 Disorder relevance: Proof of Corollary 1.7(ii)

Proof. We restrict the expectation in (5.18) to the set

$$A_n := \{(S_k)_{k=0}^n : I \cap \{1, \dots, n\} = I' \cap \{1, \dots, n\}\}, \quad (8.1)$$

i.e., S follows I' and collects only the tilted charges \hat{x}_k defined in (5.17). This gives the lower bound

$$\exp \left[\sum_{k=0}^{n-1} [\beta \hat{x}_k - \log M(\beta)] 1_{\{k \in I'\}} \right] \mathbf{P}(A_n). \quad (8.2)$$

Let $k_n := |I \cap \{1, \dots, n\}|$, $\tau'_0 = 0$ and $\tau'_1 < \dots < \tau'_{k_n}$ the elements of $I' \cap \{1, \dots, n\}$. By the renewal theorem, we have $k_n/n \rightarrow 1/m_{\text{tr}}$ as $n \rightarrow \infty$. Moreover,

$$\mathbf{P}(A_n) = \mathbf{P}(\tau_1 > n - \tau'_{k_n}) \prod_{i=1}^{k_n} K^{\text{tr}}(\tau'_i - \tau'_{i-1}), \quad (8.3)$$

so that

$$\frac{1}{n} \log \mathbf{P}(A_n) = \frac{1}{n} \log \mathbf{P}(\tau_1 > n - \tau_{k_n}) + \frac{k_n}{n} \frac{1}{k_n} \sum_{i=1}^{k_n} \log K^{\text{tr}}(\tau'_i - \tau'_{i-1}) \rightarrow \frac{1}{m_{\text{tr}}} \sum_{k=1}^{\text{tr}} K^{\text{tr}}(k) \log K^{\text{tr}}(k), \quad (8.4)$$

while

$$\frac{1}{n} \sum_{k=0}^{n-1} \{\beta \hat{x}_k - \log M(\beta)\} 1_{\{k \in I'\}} \rightarrow \frac{1}{m_{\text{tr}}} c(\beta) \quad (8.5)$$

with

$$c(\beta) := \beta E_{\mu_\beta}(\hat{x}_1) - \log M(\beta) = \beta [\log M(\beta)]' - \log M(\beta) = h(\mu_\beta | \mu_0). \quad (8.6)$$

Hence

$$m_{\text{tr}} H(\Psi_{[Q_\beta]_{\text{tr}}} | \mu_0^{\otimes N_0}) \geq h(\mu_\beta | \mu_0) + \sum_{k=0}^{\text{tr}} K^{\text{tr}}(k) \log K^{\text{tr}}(k), \quad (8.7)$$

and

$$\liminf_{\text{tr} \rightarrow \infty} m_{[Q_\beta]_{\text{tr}}} H(\kappa([Q_\beta]_{\text{tr}}) | \mu_0^{\otimes N_0}) \geq h(\mu_\beta | \mu_0) - H(K). \quad (8.8)$$

Consequently, $h(\mu_\beta | \mu_0) > H(K)$ is sufficient for disorder relevance. \blacksquare

We close by proving the second line of (1.26):

$$\lim_{\beta \rightarrow \infty} h(\mu_\beta | \mu_0) = \log [1/\mu_0(\{w\})]. \quad (8.9)$$

We distinguish three different cases.

(1) $w = \infty$. Apply (3.43) with $\mu = \mu_\beta$, $\nu = \mu_0$ and $f(x) = x \vee 0$, to get

$$h(\mu_\beta | \mu_0) \geq \int_E (x \vee 0) d\mu_\beta(x) - \log [M(1) + 1]. \quad (8.10)$$

The integral diverges as $\beta \rightarrow \infty$, and so (8.9) follows.

(2) $\mu_0(\{w\}) = 0$ with $w < \infty$. Now μ_β converges weakly as $\beta \rightarrow \infty$ to δ_w , the point measure at w . Hence (8.9) follows by using the lower semicontinuity of $\mu \mapsto h(\mu | \mu_0)$ and the fact that $h(\delta_w | \mu_0) = \infty$ because δ_w is not absolutely continuous w.r.t. μ_0 .

(3) $\mu_0(\{w\}) > 0$ with $w < \infty$. Define

$$f_\beta(x) := \frac{d\mu_\beta}{d\mu_0}(x) = \frac{e^{\beta x}}{M(\beta)}, \quad x \in E. \quad (8.11)$$

This function satisfies

$$\begin{aligned} \lim_{\beta \rightarrow \infty} f_\beta(x) &= 0 \text{ for } x < w, \\ \lim_{\beta \rightarrow \infty} f_\beta(w) &= 1/\mu_0(\{w\}), \\ f_\beta(x) &\leq 1/\mu_0(\{w\}) < \infty \text{ for } x \leq w. \end{aligned} \quad (8.12)$$

Since $t \mapsto t \log t$ is increasing on $[1, \infty)$ and on $(0, 1]$ takes values in $[-e^{-1}, 0]$, we can apply the bounded convergence theorem to the integral

$$h(\mu_\beta | \mu_0) = \int_E f_\beta(x) \log f_\beta(x) d\mu_0(x), \quad (8.13)$$

to get (8.9).

A Appendix

In this appendix we recall a few facts about the homopolymer. For proofs we refer to Giacomin [11], Chapter 2, and den Hollander [18], Chapter 7.

The homopolymer has a path measure as in (1.4), but with exponent $\lambda \sum_{k=0}^{n-1} 1_{\{S_k=0\}}$, $\lambda \in [0, \infty)$. For a given renewal time distribution K , the free energy $f(\lambda)$ is the unique solution of the equation

$$e^{-\lambda} = \sum_{n \in \mathbb{N}} K(n) e^{-nf(\lambda)} \quad (\text{A.1})$$

whenever a solution exists, otherwise $f(\lambda) = 0$. Clearly

$$f(\lambda) = 0 \iff \lambda \leq -\log \mathbf{P}(I \neq \emptyset), \quad (\text{A.2})$$

where $I = \{k \in \mathbb{N} : S_k = 0\}$ is the set of renewal times of S .

Let S, S' be two independent copies of the Markov chain with renewal time distribution K , with sets of renewal times I, I' . Transience of the joint renewal process $I \cap I'$ is equivalent to $\mathbf{P}(I \cap I' \neq \emptyset) < 1$. In that case, let

$$\lambda_0 := -\log \mathbf{P}(I \cap I' \neq \emptyset) > 0, \quad (\text{A.3})$$

and denote by $f_2(\lambda)$ and $f_2^{\text{tr}}(\lambda)$ the free energy when the renewal times of S, S' are drawn from K , respectively, K^{tr} defined in (5.15). Then $\lim_{\text{tr} \rightarrow \infty} f_2^{\text{tr}}(\lambda) = f_2(\lambda)$. Note that $f_2(\lambda) = 0$ iff $\lambda \leq \lambda_0$. This property does not hold for $f_2^{\text{tr}}(\lambda)$, but the following lemma shows that $f_2^{\text{tr}}(\lambda)$ tends to zero fast as $\text{tr} \rightarrow \infty$ when $\lambda < \lambda_0$.

Lemma A.1. *Suppose that $\mathbf{P}(I \cap I' \neq \emptyset) < 1$. Then $\sup_{\text{tr} \in \mathbb{N}} \text{tr} f_2^{\text{tr}}(\lambda) < \infty$ for all $\lambda < \lambda_0$.*

Proof. As in the paragraph preceding the lemma, define $I^{\text{tr}}, I'^{\text{tr}}$, where now the Markov chains S, S' have renewal time distribution K^{tr} . Let K_2, K_2^{tr} be the renewal time distributions generating the sets $I \cap I', I^{\text{tr}} \cap I'^{\text{tr}}$ respectively. Put $L_2(n) := \sum_{k=1}^n K_2(k)$ and $L_2^{\text{tr}}(n) := \sum_{k=1}^n K_2^{\text{tr}}(k)$. Then $L_2(\infty) = e^{-\lambda_0}$ and $L_2^{\text{tr}}(\infty) = 1$ because the renewal process $I^{\text{tr}} \cap I'^{\text{tr}}$ is recurrent. Since $K_2^{\text{tr}}(n) = K_2(n)$ for $1 \leq n < \text{tr}$, it follows from (A.1) that

$$\begin{aligned} e^{-\lambda} &= \sum_{n=1}^{\text{tr}-1} K_2(n) e^{-nf_2^{\text{tr}}(\lambda)} + \sum_{n=\text{tr}}^{\infty} K_2^{\text{tr}}(n) e^{-nf_2^{\text{tr}}(\lambda)} \\ &\leq L_2(\text{tr}-1) + e^{-\text{tr} f_2^{\text{tr}}(\lambda)} [1 - L_2(\text{tr}-1)], \end{aligned} \quad (\text{A.4})$$

where the equality holds because $f_2^{\text{tr}}(\lambda) > 0$ for $\lambda > 0$. Hence

$$\text{tr} f_2^{\text{tr}}(\lambda) \leq \log \left[\frac{1 - L_2(\text{tr}-1)}{e^{-\lambda} - L_2(\text{tr}-1)} \right]. \quad (\text{A.5})$$

The term between brackets tends to $(1 - e^{-\lambda_0}) / (e^{-\lambda} - e^{-\lambda_0})$ as $\text{tr} \rightarrow \infty$, which is finite for $\lambda < \lambda_0$. ■

The order of the phase transition for the homopolymer depends on the tail of K . If K satisfies (1.24), then

$$f(\lambda) \sim \lambda^{1/(1 \wedge \alpha)} L^*(1/\lambda), \quad \lambda \downarrow 0, \quad (\text{A.6})$$

for some L^* that is strictly positive and slowly varying at infinity. Hence, the phase transition is order 1 when $\alpha \in [1, \infty)$ and order $m \in \mathbb{N} \setminus \{1\}$ when $\alpha \in [\frac{1}{m}, \frac{1}{m-1})$. This shows that the value $\alpha = \frac{1}{2}$ is critical in view of the Harris criterion mentioned in Remark 1.13.

References

- [1] K.S. Alexander, The effect of disorder on polymer depinning transitions, *Commun. Math. Phys.* 279 (2008) 117–146.
- [2] K.S. Alexander and V. Sidoravicius, Pinning of polymers and interfaces by random potentials, *Ann. Appl. Probab.* 16 (2006) 636–669.
- [3] K.S. Alexander and N. Zygouras, Quenched and annealed critical points in polymer pinning models, *Commun. Math. Phys.* 291 (2009) 659–689.
- [4] K.S. Alexander and N. Zygouras, Equality of critical points for polymer depinning transitions with loop exponent one, *Ann. Appl. Probab.* 20 (2010) 356–366.
- [5] M. Birkner, Conditional large deviations for a sequence of words, *Stoch. Proc. Appl.* 118 (2008) 703–729.
- [6] M. Birkner, A. Greven and F. den Hollander, Quenched large deviation principle for words in a letter sequence, *Probab. Theory Relat. Fields* 147 (2010), available online.
- [7] A. Dembo and O. Zeitouni, *Large Deviations Techniques and Applications* (2nd. ed.), Springer, New York, 1998.
- [8] B. Derrida, G. Giacomin, H. Lacoïn and F.L. Toninelli, Fractional moment bounds and disorder relevance for pinning models, to appear in *Commun. Math. Phys.* 287 (2009) 867–887.
- [9] R.A. Doney, One-sided local large deviation and renewal theorems in the case of infinite mean, *Probab. Theory Relat. Fields* 107 (1997) 451–465.
- [10] H. O. Georgii, *Gibbs measures and phase transitions*. de Gruyter Studies in Mathematics, 9. Walter de Gruyter & Co., Berlin, 1988.
- [11] G. Giacomin, *Random Polymer Models*, Imperial College Press, London, 2007.
- [12] G. Giacomin, H. Lacoïn and F.L. Toninelli, Marginal relevance of disorder for pinning models, to appear in *Comm. Pure Appl. Math.*
- [13] G. Giacomin, H. Lacoïn and F.L. Toninelli, Disorder relevance at marginality and critical point shift, preprint 2009.
- [14] G. Giacomin and F.L. Toninelli, On the irrelevant disorder regime of pinning models, *Ann. Probab.* 37 (2007) 1841–1875.
- [15] R.M. Gray, *Probability, Random Processes, and Ergodic Properties*, Springer, New York, 1988. Available at <http://ee.stanford.edu/~gray/>
- [16] R.M. Gray, *Entropy and Information Theory*, Springer, New York, 1990. Available at <http://ee.stanford.edu/~gray/>
- [17] A.B. Harris, Effect of random defects on the critical behaviour of Ising models, *J. Phys. C* 7 (1974) 1671–1692.

- [18] F. den Hollander, *Random Polymers*, Lectures Notes in Mathematics 1974, Springer, Berlin, 2009.
- [19] H. Lacoïn, The martingale approach to disorder irrelevance for pinning models, preprint 2010.
- [20] F. Spitzer, *Principles of Random Walk* (2nd. ed.), Springer, Berlin, 1976.
- [21] F.L. Toninelli, A replica-coupling approach to disordered pinning models, *Commun. Math. Phys.* 280 (2008) 389–401.
- [22] F.L. Toninelli, Disordered pinning models and copolymers: beyond annealed bounds, *Ann. Appl. Probab.* 18 (2008) 1569–1587.
- [23] P. Walters, *An Introduction to Ergodic Theory*, Graduate Texts in Mathematics 79, Springer, New York, 1982.