

Existence and exponential mixing of infinite white α -stable systems with unbounded interactions

Citation for published version (APA):

Xu, L., & Zegarliński, B. (2011). *Existence and exponential mixing of infinite white α -stable systems with unbounded interactions*. (Report Eurandom; Vol. 2011001). Eurandom.

Document status and date:

Published: 01/01/2011

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EURANDOM PREPRINT SERIES
2011-001

**Existence and Exponential mixing of infinite white α -stable
Systems with unbounded interactions**

Lihu Xu and Boguslaw Zegarlinski
ISSN 1389-2355

Existence and Exponential mixing of infinite white α -stable Systems with unbounded interactions*

Lihu Xu

PO Box 513, EURANDOM, 5600 MB Eindhoven, The Netherlands.

Email: xu@math.tu-berlin.de.

Boguslaw Zegarlinski

Mathematics Department, Imperial College London, SW7 2AZ, United Kingdom.

Email: b.zegarlinski@imperial.ac.uk.

Abstract

We study an infinite white α -stable systems with unbounded interactions, and prove the existence of a solution by Galerkin approximation and an exponential mixing property by an α -stable version of gradient bounds.

Key words: Exponential mixing, White symmetric α -stable processes, Lie bracket, Finite speed of propagation of information, Gradient bounds.

AMS 2000 Subject Classification: Primary 37L55, 60H10, 60H15.

Submitted to EJP on January 19, 2010, final version accepted November 9, 2010.

*We would like to thank Jerzy Zabczyk for pointing out an error in Proposition 2.7 of the original version. The first author would like to thank the hospitality of Mathematics department of Université Paul Sabatier of Toulouse, part of his work was done during visiting Toulouse.

1 Introduction

The SPDEs driven by Lévy noises were intensively studied in the past several decades ([24], [3],[25], [28], [7], [5], [22], [21], ...). The noises can be Wiener([11],[12]) Poisson ([5]), α -stable types ([27],[33]) and so on. To our knowledge, many of these results in these articles are in the frame of Hilbert space, and thus one usually needs to assume that the Lévy noises are square integrable. This assumption rules out a family of important Lévy noises – α -stable noises. On the other hand, the ergodicity of SPDEs has also been intensively studied recently ([12],[18], [30], [33], [15]), most of these known results are about the SPDEs driven by Wiener type noises. There exist few results on the ergodicity of the SPDEs driven by the jump noises ([33], [24]).

In this paper, we shall study an interacting spin system driven by *white* symmetric α -stable noises ($1 < \alpha \leq 2$). More precisely, our system is described by the following infinite dimensional SDEs: for each $i \in \mathbb{Z}^d$,

$$\begin{cases} dX_i(t) = [J_i(X_i(t)) + I_i(X(t))]dt + dZ_i(t) \\ X_i(0) = x_i \end{cases} \quad (1.1)$$

where $X_i, x_i \in \mathbb{R}$, $\{Z_i; i \in \mathbb{Z}^d\}$ are a sequence of i.i.d. symmetric α -stable processes with $1 < \alpha \leq 2$, and the assumptions for the I and J are specified in Assumption 2.2. Equation (1.1) can be considered as a SPDEs in some Banach space, we shall study the existence of the dynamics, Markov property and the exponential mixing property. When $Z(t)$ is Wiener noise, the equation (1.1) has been intensively studied in modeling quantum spin systems in the 90s of last century (see e.g. [1], [2], [12], ...). Besides this, we have the other two motivations to study (1.1) as follows.

The first motivation is to extend the known existence and ergodic results about the interacting system in Chapter 17 of [24]. In that book, some interacting systems similar to (1.1) were studied under the framework of SPDEs ([11], [12]). In order to prove the existence and ergodicity, one needs to assume that the noises are square integrable and that the interactions are linear and finite range. Comparing with the systems in [24], the white α -stable noises in (1.1) are *not* square integrable, the interactions I_i are not linear but *Lipschitz* and have *infinite range*. Moreover, we shall not work on Hilbert space but on some considerably large subspace \mathbb{B} of $\mathbb{R}^{\mathbb{Z}^d}$, which seems more *natural* (see Remark 2.1). The advantage of using this subspace is that we can split it into compact balls (under product topology) and control some important quantities in these balls (see Proposition 3.1 for instance). Besides the techniques in SPDEs, we shall also use those in interacting particle systems such as finite speed of propagation of information property.

The second motivation is from the work by [35] on interacting unbounded spin systems driven by Wiener noise. The system studied there is also similar to (1.1), but has two essential differences. [35] studied a *gradient* system perturbed by *Wiener* noises, it is not hard to show the stochastic systems is reversible and admits a unique invariant measure μ . Under the framework of $L^2(\mu)$, the generator of the system is self-adjoint and thus we can construct dynamics by the spectral decomposition technique. However, the deterministic part in (1.1) is *not* necessarily a gradient type and the noises are more general. This means that our system is possibly not reversible, so we have to construct the dynamics by some other method. More precisely, we shall prove the existence of the dynamics by studying some Galerkin approximation, and passing to its limit by the finite speed

of propagation and some uniform bounds of the approximate dynamics. On the other hand, [35] proved the following pointwise ergodicity $|P_t f(x) - \mu(f)| \leq C(f, x)e^{-mt}$, where P_t is the semigroup generated by a reversible generator. The main tool for proving this ergodicity is by a logarithmic Sobolev inequality (LSI). Unfortunately, the LSI is not available in our setting, however, we can use the spirit of *Bakry-Emery criterion* in LSI to obtain a *gradient* bounds, from which we show the same ergodicity result as in [35]. We remark that although such strategy could be in principle applied to models considered in [35], unlike the method based on LSI (where only asymptotic mixing is relevant), in the present level of technology it can only cover the weak interaction regime far from the ‘critical point’.

Let us give two concrete examples for our system (1.1). The first one is by setting $I_i(x) = \sum_{j \in \mathbb{Z}^d} a_{ij} x_j$ and $J_i(x_i) = -(1 + \varepsilon)x_i - cx_i^{2n+1}$ with any $\varepsilon > 0$, $c \geq 0$ and $n \in \mathbb{N}$ for all $i \in \mathbb{Z}^d$, where (a_{ij}) is a transition probability of random walk on \mathbb{Z}^d . If we take $c = 0$ and $Z_i(t) = B_i(t)$ in (1.1) with $(B_i(t))_{i \in \mathbb{Z}^d}$ i.i.d. standard Brownian motions, then this example is similar to the neutral stepping stone model (see [13], or see a more simple introduction in [32]) and the interacting diffusions ([16], [19]) in stochastic population dynamics. We should point out that there are some essential differences between these models and this example, but it is interesting to try our method to prove the results in [19].

The organization of the paper is as follows. Section 2 introduces some notations and assumptions which will be used throughout the paper, and gives two key estimates. In third and fourth sections, we shall prove the main theorems – Theorem 2.3 and Theorem 2.4 respectively.

2 Notations, assumptions, main results and two key estimates

2.1 Notations, assumptions and main results

We shall first introduce the definition of symmetric α -stable processes ($0 < \alpha \leq 2$), and then give more detailed description for the system (1.1).

Let $Z(t)$ be one dimensional α -stable process ($0 < \alpha \leq 2$), as $0 < \alpha < 2$, it has infinitesimal generator ∂_x^α ([4]) defined by

$$\partial_x^\alpha f(x) = \frac{1}{C_\alpha} \int_{\mathbb{R} \setminus \{0\}} \frac{f(y+x) - f(x)}{|y|^{\alpha+1}} dy \quad (2.1)$$

with $C_\alpha = - \int_{\mathbb{R} \setminus \{0\}} (\cos y - 1) \frac{dy}{|y|^{1+\alpha}}$. As $\alpha = 2$, its generator is $\frac{1}{2}\Delta$. One can also define $Z(t)$ by Poisson point processes or by Fourier transform ([8]). The α -stable property means

$$Z(t) \stackrel{d}{=} t^{1/\alpha} Z(1). \quad (2.2)$$

Note that we have use the symmetric property of ∂_x^α in the easy identity $[\partial_x^\alpha, \partial_x] = 0$ where $[\cdot, \cdot]$ is the Lie bracket. The *white* symmetric α -stable processes are defined by

$$\{Z_i(t)\}_{i \in \mathbb{Z}^d}$$

where $\{Z_i(t)\}_{i \in \mathbb{Z}^d}$ are a sequence of i.i.d. symmetric α -stable process defined as the above.

We shall study the system (1.1) on $\mathbb{B} \subset \mathbb{R}^{\mathbb{Z}^d}$ defined by

$$\mathbb{B} = \bigcup_{R>0, \rho>0} B_{R, \rho}$$

where for any $R, \rho > 0$

$$B_{R, \rho} = \{x = (x_i)_{i \in \mathbb{Z}^d}; |x_i| \leq R(|i| + 1)^\rho\} \text{ with } |i| = \sum_{k=1}^d |i_k|.$$

Remark 2.1. The above \mathbb{B} is a considerably large subspace of $\mathbb{R}^{\mathbb{Z}^d}$. Define the subspace $l_{-\rho} := \{x \in \mathbb{R}^{\mathbb{Z}^d}; \sum_{k \in \mathbb{Z}^d} |k|^{-\rho} |x_k| < \infty\}$, it is easy to see that $l_{-\rho} \subset \mathbb{B}$ for all $\rho > 0$. Moreover, one can also check that the distributions of the white α -stable processes $(Z_i(t))_{i \in \mathbb{Z}^d}$ at any fixed time t are supported on \mathbb{B} . From the form of the equation (1.1), one can expect that the distributions of the system at any fixed time t is similar to those of white α -stable processes but with some (complicated) shifts. Hence, it is *natural* to study (1.1) on \mathbb{B} .

Assumption 2.2 (Assumptions for I and J). *The I and J in (1.1) satisfies the following conditions:*

1. For all $i \in \mathbb{Z}^d$, $I_i : \mathbb{B} \rightarrow \mathbb{R}$ is a continuous function under the product topology on \mathbb{B} such that

$$|I_i(x) - I_i(y)| \leq \sum_{j \in \mathbb{Z}^d} a_{ji} |x_j - y_j|$$

where $a_{ij} \geq 0$ satisfies the conditions: \exists some constants $K > 0$ and $\gamma > 0$ such that

$$a_{ij} \leq K e^{-|i-j|^\gamma}.$$

2. For all $i \in \mathbb{Z}^d$, $J_i : \mathbb{R} \rightarrow \mathbb{R}$ is a differentiable function such that

$$\frac{d}{dx} J_i(x) \leq 0 \quad \forall x \in \mathbb{R};$$

and for some $\kappa, \kappa' > 0$

$$|J_i(x)| \leq \kappa' (|x|^\kappa + 1) \quad \forall x \in \mathbb{R}.$$

3. $\eta := \left(\sup_{j \in \mathbb{Z}^d} \sum_{i \in \mathbb{Z}^d} a_{ij} \right) \vee \left(\sup_{i \in \mathbb{Z}^d} \sum_{j \in \mathbb{Z}^d} a_{ij} \right) < \infty$, $c := \inf_{i \in \mathbb{Z}^d, y \in \mathbb{R}} \left(-\frac{d}{dy} J_i(y) \right)$.

Without loss of generality, we assume that $I_i(0) = 0$ for all $i \in \mathbb{Z}^d$ and that $K' = 0$, $K = 1$ and $\gamma = 1$ in Assumption 2.2 from now on, i.e.

$$a_{ij} \leq e^{-|i-j|} \quad \forall i, j \in \mathbb{Z}^d. \quad (2.3)$$

Without loss of generality, we also assume from now on

$$J_i(0) = 0 \quad \forall i \in \mathbb{Z}^d. \quad (2.4)$$

Let us now list some notations to be frequently used in the paper, and then give the main results, i.e. Theorems 2.3 and 2.4.

- Define $|i - j| = \sum_{1 \leq k \leq d} |i_k - j_k|$ for any $i, j \in \mathbb{Z}^d$, define $|\Lambda|$ the cardinality of any given finite set $\Lambda \subset \mathbb{Z}^d$.
- For the national simplicity, we shall write $\partial_i := \partial_{x_i}$, $\partial_{ij} := \partial_{x_i x_j}^2$ and $\partial_i^\alpha := \partial_{x_i}^\alpha$. It is easy to see that $[\partial_i^\alpha, \partial_j] = 0$ for all $i, j \in \mathbb{Z}^d$.
- For any finite sublattice $\Lambda \subset \subset \mathbb{Z}^d$, let $C_b(\mathbb{R}^\Lambda, \mathbb{R})$ be the bounded continuous function space from \mathbb{R}^Λ to \mathbb{R} , denote $\mathcal{D} = \bigcup_{\Lambda \subset \subset \mathbb{Z}^d} C_b(\mathbb{R}^\Lambda, \mathbb{R})$ and

$$\mathcal{D}^k = \{f \in \mathcal{D}; f \text{ has bounded } 0, \dots, k \text{th order derivatives}\}.$$

- For any $f \in \mathcal{D}$, denote $\Lambda(f)$ the localization set of f , i.e. $\Lambda(f)$ is the smallest set $\Lambda \subset \mathbb{Z}^d$ such that $f \in C_b(\mathbb{R}^\Lambda, \mathbb{R})$.
- For any $f \in C_b(\mathbb{B}, \mathbb{R})$, define $\|f\| = \sup_{x \in \mathbb{B}} |f(x)|$. For any $f \in \mathcal{D}^1$, define $|\nabla f(x)|^2 = \sum_{i \in \mathbb{Z}^d} |\partial_i f(x)|^2$ and

$$\|f\| = \sum_{i \in \mathbb{Z}^d} \|\partial_i f\|.$$

Theorem 2.3. *There exists a Markov semigroup P_t on the space $\mathcal{B}_b(\mathbb{B}, \mathbb{R})$ generated by the system (1.1).*

Theorem 2.4. *If $c \geq \eta + \delta$ with any $\delta > 0$ and c, η defined in (3) of Assumption 2.2, then there exists some probability measure μ supported on \mathbb{B} such that for all $x \in \mathbb{B}$,*

$$\lim_{t \rightarrow \infty} P_t^* \delta_x = \mu \text{ weakly.}$$

Moreover, for any $x \in \mathbb{B}$ and $f \in \mathcal{D}^2$, there exists some $C = C(\Lambda(f), \eta, c, x) > 0$ such that we have

$$\left| \int_{\mathbb{B}} f(y) dP_t^* \delta_x - \mu(f) \right| \leq C e^{-\frac{1}{8} \wedge \frac{\delta}{2} t} \|f\|. \quad (2.5)$$

2.2 Two key estimates

In this subsection, we shall give an estimate for the operator a and $a + \delta$, where a is defined in Assumption 2.2 and δ is the Krockner's function, and also an estimate for a generalized 1 dimensional Ornstein-Uhlenbeck α -stable process governed by (2.8).

2.2.1 Estimates for a and $a + c\delta$

The lemma below will play an important role in several places such as proving (3.18). If $(a_{ij})_{i,j \in \mathbb{Z}^d}$ is the transition probability of a random walk on \mathbb{Z}^d , then (2.6) with $c = 0$ gives an estimate for the transition probability of the n steps walk.

Lemma 2.5. *Let a_{ij} be as in Assumption 2.2 and satisfy (2.3). Define*

$$[(c\delta + a)^n]_{ij} := \sum_{i_1, \dots, i_{n-1} \in \mathbb{Z}^d} (c\delta + a)_{ii_1} \cdots (c\delta + a)_{i_{n-1}j}$$

where $c \geq 0$ is some constant and δ is some Krockner's function, we have

$$[(c\delta + a)^n]_{ij} \leq (c + \eta)^n \sum_{k \geq |j-i|} (2k)^{nd} e^{-k} \quad (2.6)$$

Remark 2.6. Without the additional assumption (2.3), one can also have the similar estimates as above, for instance, $(a^n)_{ij} \leq \eta^n \sum_{k \geq |j-i|} (Ck)^{nd} \exp\{-k^{\gamma/2}\}$. The $C > 0$ is some constant depending on K, K' and γ , and will not play any essential roles in the later arguments.

Proof. Denote the collection of the $(n+1)$ -vortices paths connecting i and j by $\gamma_{i \sim j}^n$, i.e.

$$\gamma_{i \sim j}^n = \{(\gamma(i))_{i=1}^{n+1} : \gamma(1) = i, \gamma(2) \in \mathbb{Z}^d, \dots, \gamma(n) \in \mathbb{Z}^d, \gamma(n+1) = j\},$$

for any $\gamma \in \gamma_{i \sim j}^n$, define its length by

$$|\gamma| = \sum_{k=1}^n |\gamma(k+1) - \gamma(k)|.$$

We have

$$\begin{aligned} [(a + c\delta)^n]_{ij} &= \sum_{\gamma \in \gamma_{i \sim j}^n} (a + \delta)_{\gamma(1), \gamma(2)} \cdots (a + c\delta)_{\gamma(n), \gamma(n+1)} \\ &\leq \sum_{|\gamma|=|i-j|}^{\infty} (2|\gamma|)^{dn} (c + \eta)^n e^{-|\gamma|} \end{aligned} \quad (2.7)$$

where the inequality is obtained by the following observations:

- $\min_{\gamma \in \gamma_{i \sim j}^n} |\gamma| \geq |i - j|$.
- the number of the pathes in $\gamma_{i \sim j}^n$ with length $|\gamma|$ is bounded by $[(2|\gamma|)^d]^n$
- $(a + c\delta)_{\gamma(1), \gamma(2)} \cdots (a + c\delta)_{\gamma(n), \gamma(n+1)} = \prod_{\{k; \gamma(k+1) = \gamma(k)\}} (a + c\delta)_{\gamma(k), \gamma(k+1)} \times \prod_{\{k; \gamma(k+1) \neq \gamma(k)\}} a_{\gamma(k), \gamma(k+1)} \leq (c + \eta)^n e^{-|\gamma|}$.

□

2.2.2 1d generalized Ornstein-Uhlenbeck α -stable processes

Our generalized α -stable processes satisfies the following SDE

$$\begin{cases} dX(t) = J(X(t))dt + dZ(t) \\ X(0) = x \end{cases} \quad (2.8)$$

where $X(t), x \in \mathbb{R}$, $J : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable function with polynomial growth, $J(0) = 0$ and $\frac{d}{dx}J(x) \leq 0$, and $Z(t)$ is a one dimensional symmetric α -stable process with $1 < \alpha \leq 2$. One can write $J(x) = \frac{J(x)}{x}x$, clearly $\frac{J(x)}{x} \leq 0$ with the above assumptions (it is natural to define $\frac{J(0)}{0} = J'(0)$). $J(x) = -cx$ ($c > 0$) is a special case of the above J , this is the motivation to call (2.8) the generalized Ornstein-Uhlenbeck α -stable processes. The following uniform bound is important for proving (2) of Proposition 3.1.

Proposition 2.7. *Let $X(t)$ be the dynamics governed by (2.8) and denote $\mathcal{E}(s, t) = \exp\{\int_s^t \frac{J(X(r))}{X(r)} dr\}$. If $\sup_{x \in \mathbb{R}} \frac{J(x)}{x} \leq -\varepsilon$ with any $\varepsilon > 0$, then*

$$\mathbb{E}_x \left| \int_0^t \mathcal{E}(s, t) dZ_s \right| < C(\alpha, \varepsilon) \quad (2.9)$$

where $C(\alpha, \varepsilon) > 0$ only depends on α, ε .

Proof. From (1) of Proposition 3.1, we have

$$X(t) = \mathcal{E}(0, t)x + \int_0^t \mathcal{E}(s, t) dZ(s). \quad (2.10)$$

By integration by parts formula ([9]),

$$\begin{aligned} & \mathbb{E} \left| \int_0^t \mathcal{E}(s, t) dZ(s) \right| \\ &= \mathbb{E} \left| Z(t) - \int_0^t Z(s) d\mathcal{E}(s, t) \right| \\ &\leq \mathbb{E} |Z(t)\mathcal{E}(0, t)| + \mathbb{E} \left| \int_0^t (Z(t) - Z(s)) d\mathcal{E}(s, t) \right|. \end{aligned}$$

By (2.2), the first term on the r.h.s. of the last line is bounded by

$$\mathbb{E} |Z(t)\mathcal{E}(0, t)| \leq e^{-\varepsilon t} \mathbb{E} |Z(t)| \leq Ce^{-\varepsilon t} t^{1/\alpha} \rightarrow 0 \quad (t \rightarrow \infty).$$

As for the second term, one has

$$\begin{aligned} & \mathbb{E} \left| \int_0^t \frac{Z(t) - Z(s)}{(t-s)^{1/\gamma} \vee 1} [(t-s)^{1/\gamma} \vee 1] d\mathcal{E}(s, t) \right| \\ &\leq \mathbb{E} \left(\sup_{0 \leq s \leq t} \left| \frac{Z(t) - Z(s)}{(t-s)^{1/\gamma} \vee 1} \right| \left| \int_0^t [(t-s)^{1/\gamma} \vee 1] d\mathcal{E}(s, t) \right| \right) \end{aligned}$$

where $1 < \gamma < \alpha$. It is easy to see that $\frac{d\mathcal{E}(s,t)}{1-\mathcal{E}(0,t)}$ is a probability measure on $[0, t]$, by Jessen's inequality, we have

$$\begin{aligned} & \left(\int_0^t (t-s)^{1/\gamma} \vee 1 d\mathcal{E}(s,t) \right) \\ &= \left(\int_0^t (t-s) \vee 1 \frac{d\mathcal{E}(s,t)}{1-\mathcal{E}(0,t)} \right)^{1/\gamma} (1-\mathcal{E}(0,t)) \\ &\leq \left(\int_0^t (t-s) \vee 1 d\mathcal{E}(s,t) \right)^{1/\gamma} \leq \left(\int_0^t \mathcal{E}(s,t) ds \right)^{1/\gamma} + t\mathcal{E}(0,t) \leq C(\varepsilon, \gamma). \end{aligned}$$

On the other hand, by Doob's martingale inequality and α -stable property (2.2), for all $N \in \mathbb{N}$, we have

$$\begin{aligned} \mathbb{E} \sup_{1 \leq t \leq 2^N} \left| \frac{Z(t)}{t^{1/\gamma}} \right| &\leq \mathbb{E} \sum_{i=1}^N \sup_{2^{i-1} \leq t \leq 2^i} \left| \frac{Z(t)}{t^{1/\gamma}} \right| \leq \sum_{i=1}^N \frac{\mathbb{E} \sup_{2^{i-1} \leq t \leq 2^i} |Z(t)|}{2^{(i-1)/\gamma}} \\ &\leq C \sum_{i=1}^N \frac{2^{i/\alpha}}{2^{(i-1)/\gamma}} \leq C(\alpha, \gamma). \end{aligned}$$

From the above three inequalities, we immediately have

$$\mathbb{E} \left| \int_0^t (Z(t) - Z(s)) d\mathcal{E}(s,t) \right| \leq C(\alpha, \gamma, \varepsilon).$$

Collecting all the above estimates, we conclude the proof of (2.9). \square

3 Existence of Infinite Dimensional Interacting α -stable Systems

In order to prove the existence theorem of the equation (1.1), we shall first study its Galerkin approximation, and uniformly bound some approximate quantities. To pass to the Galerkin approximation limit, we need to apply a well known estimate in interacting particle systems – finite speed of propagation of information property.

3.1 Galerkin Approximation

Denote $\Gamma_N := [-N, N]^d$, which is a cube in \mathbb{Z}^d centered at origin. We approximate the infinite dimensional system by

$$\begin{cases} dX_i^N(t) = [J_i(X_i^N(t)) + I_i^N(X^N(t))]dt + dZ_i(t), \\ X_i^N(0) = x_i, \end{cases} \quad (3.1)$$

for all $i \in \Gamma_N$, where $x^N = (x_i)_{i \in \Gamma_N}$ and $I_i^N(x^N) = I_i(x^N, 0)$. It is easy to see that (3.1) can be written in the following vector form

$$\begin{cases} dX^N(t) = [J^N(X^N(t)) + I^N(X^N(t))]dt + dZ^N(t), \\ X^N(0) = x^N \end{cases} \quad (3.2)$$

The infinitesimal generator of (3.2) ([4], [33]) is

$$\mathcal{L}_N = \sum_{i \in \Gamma_N} \partial_i^\alpha + \sum_{i \in \Gamma_N} [J_i(x_i^N) + I_i^N(x^N)] \partial_i,$$

it is easy to see that

$$[\partial_k, \mathcal{L}_N] = (\partial_k J_k(x_k^N)) \partial_k + \sum_{i \in \Gamma_N} (\partial_k I_i^N(x^N)) \partial_i. \quad (3.3)$$

We shall study the mild solution of Eq. (3.2) in the sense that for each $i \in \Gamma_N$,

$$X_i(t) = \mathcal{E}_i(0, t)x_i + \int_0^t \mathcal{E}_i(s, t)I_i^N(X^N(s))ds + \int_0^t \mathcal{E}_i(s, t)dZ_i(s), \quad (3.4)$$

where $\mathcal{E}_i(s, t) = \exp\{\int_s^t \frac{J_i(X_i^N(r))}{X_i^N(r)} dr\}$ with $\frac{J_i(0)}{0} := J_i'(0)$.

The following proposition is important for proving the main theorems. (3) is the key estimates for obtaining the limiting semigroup of (1.1), while (2) plays the crucial role in proving the ergodicity.

Proposition 3.1. *Let I_i, J_i satisfy Assumption 2.2, together with (2.3) and (2.4), then*

1. Eq. (3.2) has a unique mild solution $X^N(t)$ in the sense of (3.4).
2. For all $x \in B_{R, \rho}$, if $c > \eta$ with c, η defined in (3) of Assumption 2.2, we have

$$\mathbb{E}_x[|X_i^N(t)|] \leq C(\rho, R, d, \eta, c)(1 + |i|^\rho).$$

3. For all $x \in B_{R, \rho}$, we have

$$\mathbb{E}_x[|X_i^N(t)|] \leq C(\rho, R, d)(1 + |i|^\rho)(1 + t)e^{(1+\eta)t}.$$

4. For any $f \in C_b^2(\mathbb{R}^{\Gamma_N}, \mathbb{R})$, define $P_t^N f(x) = \mathbb{E}_x[f(X^N(t))]$, we have $P_t^N f(x) \in C_b^2(\mathbb{R}^{\Gamma_N}, \mathbb{R})$.

Proof. To show (1), we first formally write down the mild solution as in (1), then apply the classical Picard iteration ([9], Section 5.3). We can also prove (1) by some other method as in the appendix of [34].

For the notational simplicity, we shall drop the index N of the quantities if no confusions arise. By (1), we have

$$X_i(t) = \mathcal{E}_i(0, t)x_i + \int_0^t \mathcal{E}_i(s, t)I_i(X^N(s))ds + \int_0^t \mathcal{E}_i(s, t)dZ_i(s). \quad (3.5)$$

By (1) of Assumption 2.2 (w.l.o.g. we assume $I_i(0) = 0$ for all i),

$$\begin{aligned} |X_i(t)| &\leq \sum_{j \in \Gamma_N} \delta_{ji} \left(|x_j| + \left| \int_0^t \mathcal{E}_j(s, t)dZ_j(s) \right| \right) \\ &\quad + \int_0^t e^{-c(t-s)} \sum_{j \in \Gamma_N} a_{ji} |X_j(s)| ds. \end{aligned} \quad (3.6)$$

We shall iterate the the above inequality in two ways, i.e. the following Way 1 and Way 2, which are the methods to show (2) and (3) respectively. The first way is under the condition $c > \eta$, which is crucial for obtaining a upper bound of $\mathbb{E}|X_i(t)|$ uniformly for $t \in [0, \infty)$, while the second one is without any restriction, i.e. $c \geq 0$, but one has to pay a price of an exponential growth in t .

Way 1: The case of $c > \eta$. By the definition of c, η in (3) of Assumption 2.2, (3.6) and Proposition 2.7,

$$\mathbb{E}|X_i(t)| \leq \sum_{j \in \mathbb{Z}^d} \delta_{ji}(|x_j| + C(c)) + \int_0^t e^{-c(t-s)} \sum_{j \in \mathbb{Z}^d} a_{ji} \mathbb{E}|X_j(s)| ds. \quad (3.7)$$

Iterating (3.7) once, one has

$$\begin{aligned} \mathbb{E}|X_i(t)| &\leq \sum_{j \in \mathbb{Z}^d} \delta_{ji}(|x_j| + C(c)) + \sum_{j \in \mathbb{Z}^d} \frac{a_{ji}}{c} (|x_j| + C(c)) \\ &\quad + \int_0^t e^{-c(t-s)} \int_0^s e^{-c(s-r)} \sum_{j \in \mathbb{Z}^d} (a^2)_{ji} \mathbb{E}|X_j(r)| dr ds, \end{aligned} \quad (3.8)$$

where $C(c) > 0$ is some constant only depending on c and α (but we omit α since it does not play any crucial role here). Iterating (3.7) infinitely many times, we have

$$\begin{aligned} \mathbb{E}|X_i(t)| &\leq \sum_{n=0}^M \frac{1}{c^n} \sum_{j \in \mathbb{Z}^d} (a^n)_{ji} (|x_j| + C(c)) + R_M \\ &\leq \sum_{n=0}^{\infty} \frac{1}{c^n} \sum_{j \in \mathbb{Z}^d} (a^n)_{ji} |x_j| + \frac{C(c)}{1 - \eta/c} \end{aligned} \quad (3.9)$$

where R_M is an M -tuple integral (see the double integral in (3.8)) and $\lim_{M \rightarrow \infty} R_M = 0$. To estimate the double summation in the last line, we split the sum ' $\sum_{j \in \mathbb{Z}^d} \dots$ ' into two pieces, and control them by (2.6) and $\frac{1}{c^n}$ respectively. More precisely, let $\Lambda(i, n) \subset \mathbb{Z}^d$ be a cube centered at i such that $\text{dist}(i, \Lambda^c(i, n)) = n^2$ (up to some $O(1)$ correction), one has

$$\sum_{n=1}^{\infty} \frac{1}{c^n} \sum_{j \in \mathbb{Z}^d} (a^n)_{ji} |x_j| = \sum_{n=1}^{\infty} \frac{1}{c^n} \left(\sum_{j \in \Lambda(i, n)} + \sum_{j \in \Lambda^c(i, n)} \right) (a^n)_{ji} |x_j|. \quad (3.10)$$

Since $x \in B_{R,\rho}$, we have by (2.6) with $c = 0$ therein

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{1}{c^n} \sum_{j \in \Lambda^c(i,n)} (a^n)_{ji} |x_j| \\
& \leq R \sum_{n=0}^{\infty} \frac{1}{c^n} \sum_{j \in \Lambda^c(i,n)} (a^n)_{ji} (|j|^\rho + 1) \\
& \leq C(R, \rho) \sum_{n=0}^{\infty} \frac{1}{c^n} \sum_{j \in \Lambda^c(i,n)} (a^n)_{ji} (|j - i|^\rho + |i|^\rho + 1) \\
& \leq C(R, \rho) \sum_{n=0}^{\infty} \frac{\eta^n}{c^n} \sum_{j \in \Lambda^c(i,n)} \sum_{k \geq |j-i|} (2k)^{nd} e^{-\frac{1}{2}k} e^{-\frac{1}{2}k} (|j - i|^\rho + |i|^\rho + 1) \\
& \leq C(R, \rho) \sum_{n=1}^{\infty} \frac{\eta^n}{c^n} \sum_{k \geq n^2} (2k)^{nd} e^{-\frac{1}{2}k} \sum_{j \in \Lambda^c(i,n)} e^{-\frac{1}{2}|j-i|} (|j - i|^\rho + |i|^\rho + 1) \\
& \leq C(\rho, R, d)(1 + |i|^\rho)
\end{aligned} \tag{3.11}$$

where the last inequality is by the fact $\sum_{k \geq n^2} (2k)^{nd} e^{-\frac{1}{2}k} \leq \sum_{k \geq 1} e^{-\frac{1}{2}k + nd \log(2k)} < \infty$ and the fact $\sum_{j \in \Lambda^c(i,n)} e^{-\frac{1}{2}|j-i|} |j - i|^\rho \leq \sum_{j \in \mathbb{Z}^d} e^{-\frac{1}{2}|j-i|} |j - i|^\rho < \infty$. For the other piece, one has

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{1}{c^n} \sum_{j \in \Lambda(i,n)} (a^n)_{ji} |x_j| \\
& \leq C(R, \rho) \sum_{n=0}^{\infty} \frac{1}{c^n} \sum_{j \in \Lambda(i,n)} (a^n)_{ji} (|j - i|^\rho + |i|^\rho + 1) \\
& \leq C(R, \rho) \sum_{n=0}^{\infty} \frac{\eta^n}{c^n} |\Lambda(i, n)| (n^{2\rho} + |i|^\rho + 1) \\
& \leq C(\rho, R) \sum_{n=0}^{\infty} \frac{\eta^n}{c^n} n^{2d} (n^{2\rho} + |i|^\rho + 1) \\
& \leq C(R, \rho, \eta, c)(1 + |i|^\rho).
\end{aligned} \tag{3.12}$$

Collecting (3.9), (3.11) and (3.12), we immediately obtain (2).

Way 2: The general case of $c \geq 0$. By the integration by parts, Doob's martingale inequality and the

easy relation $d\mathcal{E}_j(s, t) = \mathcal{E}_j(s, t)[-L_j(X(s))]ds$ where $L_j(x) = \frac{J_j(x)}{x}$, we have

$$\begin{aligned}
& \mathbb{E} \left| \int_0^t \mathcal{E}_j(s, t) dZ_j(s) \right| \\
& \leq \mathbb{E}|Z_j(t)| + \mathbb{E} \left| \int_0^t \mathcal{E}_j(s, t) L_j(X(s)) Z_j(s) ds \right| \\
& \leq Ct^{1/\alpha} + \mathbb{E} \left[\sup_{0 \leq s \leq t} |Z_j(s)| \left| \int_0^t \mathcal{E}_j(s, t) (-L_j(X(s))) ds \right| \right] \\
& \leq Ct^{1/\alpha} + \mathbb{E} \sup_{0 \leq s \leq t} |Z_j(s)| \\
& \leq Ct^{1/\alpha}.
\end{aligned} \tag{3.13}$$

By (3.6) and (3.13), one has

$$\mathbb{E}|X_i(t)| \leq \sum_{j \in \mathbb{Z}^d} \delta_{ji} (|x_j| + Ct^{\frac{1}{\alpha}}) + \int_0^t \sum_{j \in \mathbb{Z}^d} (\delta + a)_{ji} \mathbb{E}|X_j(s)| ds \tag{3.14}$$

Iterating the above inequality infinitely many times,

$$\mathbb{E}|X_i(t)| \leq \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{j \in \mathbb{Z}^d} [(\delta + a)^n]_{ji} |x_j| + Ce^{(1+\eta)t} t^{\frac{1}{\alpha}}, \tag{3.15}$$

By estimating the double summation in the last line by the same method as in Way 1, we finally obtain (3).

(4) immediately follows from Proposition 5.6.10 and Corollary 5.6.11 in [9]. \square

3.2 Finite speed of propagation of information property

The following relation (3.18) is usually called finite speed of propagation of information property ([17]), which roughly means that the effects of the initial condition (i.e. f in our case) need a long time to be propagated (by interactions) far away. The main reason for this phenomenon is that the interactions are finite range or sufficiently weak at long range.

From the view point of PDEs, (3.18) implies equicontinuity of $P_t^N f(x)$ under product topology on any $B_{\rho, R}$, combining this with the fact that $P_t^N f(x)$ are uniformly bounded, we can find some subsequence $P_t^{N_k} f(x)$ uniformly converge to a limit $P_t f(x)$ on $B_{\rho, R}$ by Ascoli-Arzela Theorem (notice that $B_{\rho, R}$ is compact under product topology). This is also another motivation of establishing the estimates (3.18).

Lemma 3.2.

1. For any $f \in \mathcal{D}^2$, we have

$$\sum_{k \in \mathbb{Z}^d} \|\partial_k P_t^N f\|^2 \leq e^{2\eta t} \|f\|^2. \tag{3.16}$$

and

$$\|P_t^N f\| \leq C(I, t) \|f\|. \quad (3.17)$$

where $C(I, t) > 0$, depending on the interaction I and t , is an increasing function of t .

2. (Finite speed of propagation of information property) Given any $f \in \mathcal{D}^2$ and $k \notin \Lambda(f)$, for any $0 < A \leq 1/4$, there exists some $B \geq 8$ such that when $n_k > Bt$, we have

$$\|\partial_k P_t^N f\|^2 \leq 2e^{-At - An_k} \|f\|^2 \quad (3.18)$$

where $n_k = \lceil \sqrt{\text{dist}(k, \Lambda(f))} \rceil$.

Proof. For the notational simplicity, we shall drop the parameter N of P_t^N in the proof. By the fact $\lim_{t \rightarrow 0^+} \frac{P_t F^2 - F^2}{t} \geq \lim_{t \rightarrow 0^+} \frac{(P_t F)^2 - F^2}{t}$, one has $\mathcal{L}_N F^2 - 2F \mathcal{L}_N F \geq 0$. Hence, for any $f \in \mathcal{D}^2$, by (3.3) and the fact $\partial_k J_k \leq 0$, we have the following calculation

$$\begin{aligned} \frac{d}{ds} P_{t-s} (\partial_k P_s f)^2 &= -P_{t-s} \left[\mathcal{L}_N (\partial_k P_s f)^2 - 2(\partial_k P_s f) \partial_k (\mathcal{L}_N P_s f) \right] \\ &= -P_{t-s} \left[\mathcal{L}_N (\partial_k P_s f)^2 - 2(\partial_k P_s f) \mathcal{L}_N (\partial_k P_s f) \right] \\ &\quad + 2P_{t-s} ((\partial_k P_s f) [\partial_k, \mathcal{L}_N] P_s f) \\ &\leq 2P_{t-s} ((\partial_k P_s f) [\partial_k, \mathcal{L}_N] P_s f) \\ &= 2P_{t-s} \left((\partial_k P_s f) \sum_{i \in \Gamma_N} (\partial_k I_i) \partial_i P_s f \right) \\ &\quad + 2P_{t-s} ((\partial_k P_s f) (\partial_k J_k) \partial_k P_s f) \\ &\leq 2P_{t-s} \left((\partial_k P_s f) \sum_{i \in \Gamma_N} (\partial_k I_i) \partial_i P_s f \right). \end{aligned} \quad (3.19)$$

Moreover, by the above inequality, Assumption 2.2, and the inequality of arithmetic and geometric means in order,

$$\begin{aligned} |\partial_k P_t f|^2 &\leq \|\partial_k f\|^2 + 2 \int_0^t P_{t-s} \left(|\partial_k P_s f| \sum_{i \in \Gamma_N} |\partial_k I_i| |\partial_i P_s f| \right) ds \\ &\leq \|\partial_k f\|^2 + \eta \int_0^t P_{t-s} (|\partial_k P_s f|^2) ds + \int_0^t P_{t-s} \left(\sum_{i \in \Gamma_N} a_{ki} |\partial_i P_s f|^2 \right) ds \\ &\leq \|\partial_k f\|^2 + \int_0^t P_{t-s} \left(\sum_{i \in \mathbb{Z}^d} (a_{ki} + \eta \delta_{ki}) |\partial_i P_s f|^2 \right) ds. \end{aligned}$$

where η is defined in (3) of Assumption 2.2. Iterating the above inequality, we have

$$\begin{aligned} \|\partial_k P_t f\|^2 &\leq \|\partial_k f\|^2 + t \sum_{i \in \mathbb{Z}^d} (a_{ki} + \eta \delta_{ki}) \|\partial_i f\|^2 \\ &\quad + \int_0^t P_{t-s_1} \int_0^{s_1} P_{s_1-s_2} \sum_{i \in \mathbb{Z}^d} [(a + \eta \delta)^2]_{ki} \|\partial_i P_{s_2} f\|^2 ds_2 ds_1 \\ &\leq \dots \leq \sum_{n=0}^N \frac{t^n}{n!} \sum_{i \in \mathbb{Z}^d} [(a + \eta \delta)^n]_{ki} \|\partial_i f\|^2 + Re(N) \end{aligned}$$

where $Re(N) \rightarrow 0$ as $N \rightarrow \infty$. Hence,

$$\|\partial_k P_t f\|^2 \leq \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{i \in \mathbb{Z}^d} [(a + \eta \delta)^n]_{ki} \|\partial_i f\|^2. \quad (3.20)$$

Summing k over \mathbb{Z}^d in the above inequality, one has

$$\begin{aligned} \sum_{k \in \mathbb{Z}^d} \|\partial_k P_t f\|^2 &\leq \sum_{k \in \mathbb{Z}^d} \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{i \in \mathbb{Z}^d} [(a + \eta \delta)^n]_{ki} \|\partial_i f\|^2 \\ &\leq \sum_{n=0}^{\infty} \frac{t^n}{n!} \sup_i \sum_{k \in \mathbb{Z}^d} [(a + \eta \delta)^n]_{ki} \sum_{i \in \mathbb{Z}^d} \|\partial_i f\|^2 \\ &\leq e^{2\eta t} \sum_{i \in \mathbb{Z}^d} \|\partial_i f\|^2 \leq e^{2\eta t} \|f\|^2 \end{aligned}$$

As for (3.17), one can also easily obtain from (3.20) that $\sum_{k \in \mathbb{Z}^d} \|\partial_k P_t^N f\| \leq C(I, t) \sqrt{\sum_{i \in \mathbb{Z}^d} \|\partial_k f\|^2} \leq C(I, t) \|f\|$ and that $C(I, t) > 0$ is an increasing function related to t .

In order to prove 2, one needs to estimate the double sum of (3.20) in a more delicate way. We shall split the sum ' $\sum_{n=0}^{\infty}$ ' into two pieces ' $\sum_{n=0}^{n_k}$ ' and ' $\sum_{n=n_k}^{\infty}$ ', with $n_k = \lceil \sqrt{\text{dist}(k, \Lambda(f))} \rceil$, and control them by (2.6) and some basic calculation respectively. More precisely, for the piece ' $\sum_{n=0}^{n_k}$ ', by (2.6) and the definition of $n_k = \lceil \sqrt{\text{dist}(k, \Lambda(f))} \rceil$, we have

$$\begin{aligned} &\sum_{n=0}^{n_k} \frac{t^n}{n!} \sum_{i \in \mathbb{Z}^d} [(a + \eta \delta)^n]_{ki} \|\partial_i f\|^2 \\ &\leq \sum_{n=0}^{n_k} \frac{t^n}{n!} \sum_{i \in \Lambda(f)} \sum_{j \geq |k-i|} (2\eta)^n 2^{nd} (j + \Lambda(f))^{dn} e^{-j} \|\partial_i f\|^2 \\ &\leq e^t \sum_{i \in \Lambda(f)} \sum_{j \geq |k-i|} \exp \left\{ dn_k \log[2(2\eta)^{1/d} (j + \Lambda(f))] - \frac{1}{4} n_k^2 - \frac{j}{4} \right\} e^{-\frac{j}{2}} \|\partial_i f\|^2 \\ &\leq C(d, \Lambda(f), \eta) e^t \sum_{i \in \Lambda(f)} \sum_{j \geq n_k^2} e^{-\frac{j}{2}} \|\partial_i f\|^2 \\ &\leq C(d, \Lambda(f), \eta) e^t e^{-\frac{1}{2} n_k^2} \|f\|^2. \end{aligned}$$

For the other piece, it is easy to see

$$\begin{aligned} & \sum_{n \geq n_k} \frac{t^n}{n!} \sum_{i \in \mathbb{Z}^d} [(a + \eta\delta)^n]_{ki} \|\partial_i f\|^2 \\ &= \sum_{n \geq n_k} \frac{t^n}{n!} \sum_{i \in \Lambda(f)} [(a + \eta\delta)^n]_{ki} \|\partial_i f\|^2 \leq \frac{t^{n_k}}{n_k!} e^{2\eta t} \|f\|^2. \end{aligned}$$

Combining (3.20) and the above two estimates, we immediately have

$$\|\partial_k P_t f\|^2 \leq \{C e^t e^{-\frac{1}{2}n_k^2} + \frac{t^{n_k}}{n_k!} e^{2\eta t}\} \|f\|^2.$$

For any $A > 0$, choosing $B \geq 1$ such that

$$2 - \log B + \log(2\eta) + \frac{2\eta}{B} \leq -2A,$$

as $n > Bt$, one has

$$\begin{aligned} \frac{t^n (2\eta)^n}{n!} e^{2\eta t} &\leq \exp\left\{n \log \frac{2\eta}{B} + 2n + (2\eta) \frac{n}{B}\right\} \\ &\leq \exp\{-2An\} \leq \exp\{-An - At\}. \end{aligned}$$

Now take $0 < A \leq 1/4$, $B \geq 8$ and n as the above, we can easily check that

$$e^t e^{-\frac{1}{2}n^2} \leq e^{-\frac{1}{4}n^2} e^{-\frac{1}{4}nBt+t} \leq e^{-An-At}.$$

Replacing n by n_k , we conclude the proof of (3.18). □

3.3 Proof of Theorem 2.3

As mentioned in the previous subsection, by (3.18) and the fact that $P_t^N f(x)$ are uniformly bounded, we can find some subsequence $P_t^{N_k} f(x)$ uniformly converges to a limit $P_t f(x)$ on $B_{\rho, R}$ by Ascoli-Arzela Theorem. However, this method cannot give more detailed description of P_t such as Markov property. Hence, we need to analyze $P_t^N f$ in a more delicate way.

Proof of Theorem 2.3. We shall prove the theorem by the following two steps:

1. $P_t f(x) := \lim_{N \rightarrow \infty} P_t^N f(x)$ exists pointwisely on $x \in \mathbb{B}$ for any $f \in \mathcal{D}^2$ and $t > 0$.
2. Extending the domain of P_t to $\mathcal{B}_b(\mathbb{B})$ and proving that P_t is Markov on $\mathcal{B}_b(\mathbb{B})$.

Step 1: To prove (1), it suffices to show that $\{P_t^N f(x)\}_N$ is a cauchy sequence for $x \in B_{R, \rho}$ with any fixed R and ρ .

Given any $M > N$ such that $\Gamma_M \supset \Gamma_N \supset \Lambda(f)$, we have by a similar calculus as in (3.19)

$$\begin{aligned} & \frac{d}{ds} P_{t-s}^M (P_s^M f - P_s^N f)^2 \\ &= -P_{t-s}^M \left[\mathcal{L}_M (P_s^M f - P_s^N f)^2 - 2 (P_s^M f - P_s^N f) \mathcal{L}_M (P_s^M f - P_s^N f) \right] \\ & \quad + 2P_{t-s}^M \left[(P_s^M f - P_s^N f) (\mathcal{L}_M - \mathcal{L}_N) P_s^N f \right] \\ & \leq 2P_{t-s}^M \left[(P_s^M f - P_s^N f) (\mathcal{L}_M - \mathcal{L}_N) P_s^N f \right], \end{aligned}$$

moreover, by the facts $\Lambda(P_s^N f) = \Gamma_N$, $\Gamma_M \supset \Gamma_N$ and $\Lambda(J_k) = k$,

$$(\mathcal{L}_M - \mathcal{L}_N) P_s^N f = \sum_{i \in \Gamma_N} (I_i^M(x^M) - I_i^N(x^N)) \partial_i P_s^N f.$$

Therefore, by Markov property of P_t^M , the following easy fact (by fundamental theorem of calculus, definition of I^M , and (1) of Assumption 2.2)

$$|I^M(x^M) - I^N(x^N)| \leq \sum_{j \in \Gamma_M \setminus \Gamma_N} a_{ji} |x_j|,$$

the assumption (2.3) (i.e. $a_{ij} \leq e^{-|i-j|}$), and (3) of Proposition 3.1 in order, we have for any $x \in B_{R,\rho}$,

$$\begin{aligned} & (P_t^M f(x) - P_t^N f(x))^2 \\ & \leq 2 \|f\|_\infty \int_0^t P_{t-s}^M \left(\sum_{i \in \Gamma_N} \sum_{j \in \Gamma_M \setminus \Gamma_N} a_{ji} |x_j| \|\partial_i P_s^N f\| \right) (x) ds \\ & \leq 2 \|f\|_\infty \sum_{i \in \Gamma_N} \sum_{j \in \Gamma_M \setminus \Gamma_N} e^{-|i-j|} \int_0^t \mathbb{E}_x[|X_j^M(t-s)|] \|\partial_i P_s^N f\| ds \\ & \leq C(t, \rho, R, d) \|f\|_\infty \sum_{i \in \Gamma_N} \sum_{j \in \Gamma_M \setminus \Gamma_N} e^{-|i-j|} (|j|^\rho + 1) \int_0^t \|\partial_i P_s^N f\| ds. \end{aligned} \tag{3.21}$$

Now let us estimate the double sum in the last line of (3.21), the idea is to split the first sum $\sum_{i \in \Gamma_N}$ into two pieces $\sum_{i \in \Lambda}$ and $\sum_{\Gamma_N \setminus \Lambda}$, and control them by $e^{-|i-j|}$ and (3.18) respectively. More precisely, take a cube $\Lambda \supset \Lambda(f)$ (to be determined later) inside Γ_N , we have by (3.17)

$$\begin{aligned} & \sum_{i \in \Lambda} \sum_{j \in \Gamma_M \setminus \Gamma_N} e^{-|i-j|} (|j|^\rho + 1) \int_0^t \|\partial_i P_s^N f\| ds \\ & \leq 2^\rho \sum_{i \in \Lambda} \sum_{j \in \Gamma_M \setminus \Gamma_N} e^{-|i-j|} (|j-i|^\rho + |i|^\rho + 1) \int_0^t \|\partial_i P_s^N f\| ds \\ & \leq 2^\rho \int_0^t \sum_{i \in \Lambda} \|\partial_i P_s^N f\| ds \sum_{k \geq \text{dist}(\Lambda, \Gamma_M \setminus \Gamma_N)} \sum_{j: |j-i|=k} e^{-k} (k^\rho + |\Lambda|^\rho + 1) \\ & \leq 2^\rho t C(I, t) \sum_{i \in \mathbb{Z}^d} \|\partial_i f\| \sum_{k \geq \text{dist}(\Lambda, \Gamma_M \setminus \Gamma_N)} (|\Lambda| + k)^d e^{-k} (k^\rho + |\Lambda|^\rho + 1) \\ & \leq \epsilon \end{aligned}$$

for arbitrary $\epsilon > 0$ as long as Γ_N, Γ_M (which depend on Λ , the interaction I , t) are both sufficiently large.

For the piece ' $\sum_{\Gamma_N \setminus \Lambda}$ ', one has by (3.18)

$$\begin{aligned}
& \sum_{i \in \Gamma_N \setminus \Lambda} \sum_{j \in \Gamma_M \setminus \Gamma_N} e^{-|i-j|} (|j|^\rho + 1) \int_0^t e^{t-s} \|\partial_i P_s^N f\| ds \\
& \leq 2^\rho e^t \sum_{i \in \Gamma_N \setminus \Lambda} \sum_{j \in \Gamma_M \setminus \Gamma_N} e^{-|i-j|} (|j - i|^\rho + |i|^\rho + 1) \int_0^t e^{-As - An_i} ds \\
& \leq C(t, \rho, A) \sum_{i \in \Gamma_N \setminus \Lambda} (1 + |i|^\rho) e^{-A[\text{dist}(i, \Lambda(f))]}^{1/2} \\
& \leq \epsilon
\end{aligned}$$

as we choose Λ big enough so that $\text{dist}(\Gamma_N \setminus \Lambda, \Lambda(f))$ is sufficiently large. Combing all the above, we immediately conclude step 1. We denote

$$P_t f(x) = \lim_{N \rightarrow \infty} P_t^N f(x).$$

Step 2: Proving that P_t is a Markov semigroup on $\mathcal{B}_b(\mathbb{B})$. We first extend P_t to be an operator on $\overline{\mathcal{B}_b(\mathbb{B})}$, then prove this new P_t satisfies semigroup and Markov property.

It is easy to see from step 1, for any fixed $x \in \mathbb{B}$, P_t is a linear functional on \mathcal{D}^2 . Since \mathbb{B} is locally compact (under product topology), by Riesz representation theorem for linear functional ([14], pp 223), we have a Radon measure on \mathbb{B} , denoted by $P_t^* \delta_x$, so that

$$P_t f(x) = P_t^* \delta_x(f). \quad (3.22)$$

By (3) of Proposition 3.1, take any $x \in \mathbb{B}$, it is clear that the approximate process $X^N(t, x^N) \in \mathbb{B}$ a.s. for all $t > 0$. Hence, for all $N > 0$, we have

$$P_t^N(1_{\mathbb{B}})(x) = \mathbb{E}[1_{\mathbb{B}}(X^N(t, x^N))] = 1 \quad \forall x \in \mathbb{B}.$$

Let $N \rightarrow \infty$, by step 1 (noticing $1_{\mathbb{B}} \in \mathcal{D}^2$), we have for all $x \in \mathbb{B}$

$$P_t 1_{\mathbb{B}}(x) = 1,$$

which immediately implies that $P_t^* \delta_x$ is a probability measure supported on \mathbb{B} . With the measure $P_t^* \delta_x$, one can easily extend the operator P_t from \mathcal{D}^2 to $\mathcal{B}_b(\mathbb{B})$ by bounded convergence theorem since \mathcal{D}^2 is dense in $\mathcal{B}_b(\mathbb{B})$ under product topology.

Now we prove the semigroup property of P_t , by bounded convergence theorem and the dense property of \mathcal{D}^2 in $\mathcal{B}_b(\mathbb{B})$, it suffices to prove this property on \mathcal{D}^2 . More precisely, for any $f \in \mathcal{D}^2$, we shall prove that for all $x \in \mathbb{B}$

$$P_{t_2+t_1} f(x) = P_{t_2} P_{t_1} f(x). \quad (3.23)$$

To this end, it suffices to show (3.23) for all $x \in B_{R,\rho}$.

On the one hand, from the first step, one has

$$\lim_{N \rightarrow \infty} P_{t_2+t_1}^N f(x) = P_{t_2+t_1} f(x) \quad \forall x \in B_{R,\rho}. \quad (3.24)$$

On the other hand, we have

$$\begin{aligned} |P_{t_2} P_{t_1} f(x) - P_{t_2}^N P_{t_1}^N f(x)| &\leq |P_{t_2} P_{t_1} f(x) - P_{t_2} P_{t_1}^N f(x)| \\ &+ |P_{t_2}^M P_{t_1}^N f(x) - P_{t_2} P_{t_1}^N f(x)| + |P_{t_2}^M P_{t_1}^N f(x) - P_{t_2}^N P_{t_1}^N f(x)|, \end{aligned} \quad (3.25)$$

with $M > N$ to be determined later according to N . It is easy to have by step 1 and bounded convergence theorem

$$|P_{t_2} P_{t_1} f(x) - P_{t_2} P_{t_1}^N f(x)| = |P_{t_2}^* \delta_x(P_{t_1} f - P_{t_1}^N f)| \rightarrow 0 \quad (3.26)$$

as $N \rightarrow \infty$. Moreover, by the first step, one has

$$|P_{t_2}^M P_{t_1}^N f(x) - P_{t_2} P_{t_1}^N f(x)| < \varepsilon \quad (3.27)$$

for arbitrary $\varepsilon > 0$ as long as $M \in \mathbb{N}$ (depending on Λ_N) is sufficiently large. As for the last term on the r.h.s. of (3.25), by the same arguments as in (3.21) and those immediately after (3.21), we have

$$\begin{aligned} &\left(P_{t_2}^M P_{t_1}^N f(x) - P_{t_2}^N P_{t_1}^N f(x) \right)^2 \\ &\leq C(t_1, t_2, \rho, R, d) \|f\|_\infty \sum_{i \in \Gamma_N} \sum_{j \in \Gamma_M \setminus \Gamma_N} e^{-|i-j|} (|j|^\rho + 1) \int_0^{t_2} \|\partial_i P_{t_1+s}^N f\| ds \\ &< \varepsilon \end{aligned} \quad (3.28)$$

for arbitrary $\varepsilon > 0$ if Γ_M and Γ_N are both sufficiently large.

Collecting (3.25)-(3.28), we have

$$\lim_{N \rightarrow \infty} P_{t_2}^N P_{t_1}^N f(x) = P_{t_2} P_{t_1} f(x),$$

which, with (3.24) and the fact $P_{t_2+t_1}^N = P_{t_2}^N P_{t_1}^N$, implies (3.23) for $x \in B_{R,\rho}$.

Since $P_t(\mathbf{1}) = 1$ and $P_t(f) \geq 0$ for any $f \geq 0$, P_t is a Markov semigroup ([17]). □

4 Proof of Ergodicity Result

The main ingredient of the proof follows the spirit of Bakry-Emery criterion for logarithmic Sobolev inequality ([6], [17]). In [6], the authors first studied the logarithmic Sobolev inequalities of some

diffusion generator by differentiating its first order square field $\Gamma_1(\cdot)$ (see the definition of Γ_1 and Γ_2 in chapter 4 of [17]) and obtained the following relations

$$\frac{d}{dt}P_{t-s}\Gamma_1(P_s f) \leq -cP_{t-s}\Gamma_2(P_s f) \quad (4.1)$$

where P_t is the semigroup generated by the diffusion generator, and $\Gamma_2(\cdot)$ is the second order square field. If $\Gamma_2(\cdot) \geq C\Gamma_1(\cdot)$, then one can obtain logarithmic Sobolev inequality. The relation $\Gamma_2(\cdot) \geq C\Gamma_1(\cdot)$ is called *Bakry-Emery criterion*.

In our case, one can also compute $\Gamma_1(\cdot), \Gamma_2(\cdot)$ of P_t^N , which have the similar relation as (4.1). It is interesting to apply this relation to prove some regularity of the semigroup P_t^N , but seems hard to obtain the gradient bounds by it. Alternatively, we replace $\Gamma_1(f)$ by $|\nabla f|^2$, which is actually not the first order square field of our case but the one of the diffusion generators, and differentiate $P_{t-s}|\nabla P_s f|^2$. We shall see that the following relation (4.4) plays the same role as the Bakry-Emery criterion.

Lemma 4.1. *If $c \geq \eta + \delta$ with any $\delta > 0$ and c, η defined in (3) of Assumption 2.2, we have*

$$|\nabla P_t^N f|^2 \leq e^{-2\delta t} P_t^N |\nabla f|^2 \quad \forall f \in \mathcal{D}^2 \quad (4.2)$$

Proof. For the notational simplicity, we drop the index N of the quantities. By a similar calculus as in (3.19), we have

$$\begin{aligned} \frac{d}{ds}P_{t-s}|\nabla P_s f|^2 &= -P_{t-s} \left(\mathcal{L}_N |\nabla P_s f|^2 - 2\nabla P_s f \cdot \mathcal{L}_N \nabla P_s f \right) \\ &\quad + 2P_{t-s} (\nabla P_s f \cdot [\nabla, \mathcal{L}_N] P_s f) \\ &\leq 2P_{t-s} (\nabla P_s f \cdot [\nabla, \mathcal{L}_N] P_s f) \\ &= 2P_{t-s} \left(\sum_{i,j \in \Gamma_N} \partial_j I_i(x) \partial_i P_s f \partial_j P_s f \right) \\ &\quad + 2P_{t-s} \left(\sum_{i \in \Gamma_N} \partial_i J_i(x_i) (\partial_i P_s f)^2 \right), \end{aligned} \quad (4.3)$$

where \cdot is the inner product of vectors in \mathbb{R}^{Γ_N} . Denote the quadratic form by

$$Q(\xi, \xi) = \sum_{i,j \in \Gamma_N} [\partial_i J_i(x_i) \delta_{ij} + \partial_j I_i(x)] \xi_i \xi_j \quad \forall \xi \in \mathbb{R}^{\Gamma_N},$$

it is easy to see by the assumption that

$$-Q(\xi, \xi) \geq \delta |\xi|^2. \quad (4.4)$$

This, combining with (4.3), immediately implies

$$\frac{d}{ds}P_{t-s}|\nabla P_s f|^2 \leq -2\delta P_{t-s} (|\nabla P_s f|^2), \quad (4.5)$$

from which we conclude the proof. \square

Let us now combining Lemma 4.1 and the finite speed of propagation of information property (3.18) to prove the ergodic result.

Proof of Theorem 2.4. We split the proof into the following three steps:

Step 1: For all $f \in \mathcal{D}^2$, $\lim_{t \rightarrow \infty} P_t f(0) = \ell(f)$ where $\ell(f)$ is some constant depending on f .

For any $\forall t_2 > t_1 > 0$, we have by triangle inequality

$$\begin{aligned} |P_{t_2} f(0) - P_{t_1} f(0)| &\leq |P_{t_2} f(0) - P_{t_2}^N f(0)| + |P_{t_2}^N f(0) - P_{t_1}^N f(0)| \\ &\quad + |P_{t_1}^N f(0) - P_{t_1} f(0)|. \end{aligned}$$

By Theorem 2.3, there exists some $N(t_1, t_2) \in \mathbb{N}$ such that as $N > N(t_1, t_2)$

$$|P_{t_2} f(0) - P_{t_2}^N f(0)| + |P_{t_1} f(0) - P_{t_1}^N f(0)| < e^{-\frac{\delta \Lambda A}{2} t_1} \|f\|. \quad (4.6)$$

Next, we show that for all $N \in \mathbb{N}$,

$$|P_{t_2}^N f(0) - P_{t_1}^N f(0)| \leq C(A, \delta, \Lambda(f)) e^{-\frac{\delta \Lambda A}{2} t_1} \|f\|. \quad (4.7)$$

By the semigroup property of P_t^N and fundamental theorem of calculus, one has

$$\begin{aligned} |P_{t_2}^N f(0) - P_{t_1}^N f(0)| &= \left| \mathbb{E}_0 \left[P_{t_1}^N f(X^N(t_2 - t_1)) - P_{t_1}^N f(0) \right] \right| \\ &= \left| \int_0^1 \mathbb{E}_0 \left[\frac{d}{d\lambda} P_{t_1}^N f(\lambda X^N(t_2 - t_1)) \right] d\lambda \right| \\ &\leq \int_0^1 \sum_{i \in \Gamma_N} \mathbb{E}_0 \left[|\partial_i P_{t_1}^N f(\lambda X^N(t_2 - t_1))| |X_i^N(t_2 - t_1)| \right] d\lambda. \end{aligned} \quad (4.8)$$

To estimate the sum $\sum_{i \in \Gamma_N}$ in the last line, we split it into two pieces $\sum_{i \in \Lambda}$ and $\sum_{i \in \Gamma_N \setminus \Lambda}$, and control them by Lemma 4.1 and the finite speed of propagation of information property in Lemma 3.2. Let us show the more details as follows.

Take $0 < A \leq 1/4$, and let $B = B(A, \eta) \geq 8$ be chosen as in Lemma 3.2. We choose a cube $\Lambda \supset \Lambda(f)$ inside Γ_N so that $\text{dist}(\Lambda^c, \Lambda(f)) = B^2 t_1^2$ (up to some order $O(1)$ correction). On the one hand, by (4.2), we clearly have $\|\partial_i P_t f\| \leq e^{-\delta t} \|f\|$ for all $i \in \Gamma_N$. Therefore, by (2) of Proposition 3.1,

$$\begin{aligned} &\sum_{i \in \Lambda} \mathbb{E}_0 \left[|\partial_i P_{t_1}^N f(\lambda X^N(t_2 - t_1))| |X_i^N(t_2 - t_1)| \right] \\ &\leq \sum_{i \in \Lambda} \|\partial_i P_{t_1}^N f\| \mathbb{E}_0 \left[|X_i^N(t_2 - t_1)| \right] \\ &\leq C \sum_{i \in \Lambda} e^{-\delta t_1} \|f\| (1 + |i|^\rho) \end{aligned} \quad (4.9)$$

As for the piece ' $\sum_{i \in \Gamma_N \setminus \Lambda}$ ', it is clear to see $n_i = \sqrt{\text{dist}(i, \Lambda(f))} \geq Bt_1$ for $i \in \Gamma_N \setminus \Lambda$, by Lemma 3.2 and (2) of Proposition 3.1, one has

$$\begin{aligned}
& \sum_{i \in \Gamma_N \setminus \Lambda} \mathbb{E}_0 \left[|\partial_i P_{t_1}^N f(\lambda X^N(t_2 - t_1))| |X_i^N(t_2 - t_1)| \right] \\
& \leq \sum_{i \in \Gamma_N \setminus \Lambda} \|\partial_i P_{t_1}^N f\| \mathbb{E}_0 \left[|X_i^N(t_2 - t_1)| \right] \\
& \leq C \sum_{i \in \Gamma_N \setminus \Lambda} e^{-An_i - At_1} (1 + |i|^\rho) \|f\|
\end{aligned} \tag{4.10}$$

Since $0 \in B_{R, \rho}$ with any $R, \rho > 0$, we take $\rho = 1$ and $R = 1$ in the previous inequalities. Combining (4.8), (4.9) and (4.10), we immediately have

$$\begin{aligned}
& |P_{t_2}^N f(0) - P_{t_1}^N f(0)| \\
& \leq C \left[\sum_{i \in \Gamma_N \setminus \Lambda} e^{-An_i - \frac{A}{2}t_1} (1 + |i|) + (B^2 t_1^2 + 1 + \Lambda(f))^{1+d} e^{-\frac{\delta}{2}t_1} \right] e^{-\frac{A\Lambda\delta}{2}t_1} \|f\|.
\end{aligned} \tag{4.11}$$

and $\sum_{i \in \Gamma_N \setminus \Lambda} e^{-An_i} (1 + |i|) \leq \sum_{i \in \mathbb{Z}^d \setminus \Lambda} e^{-An_i} (1 + |i|) < \infty$, whence (4.7) follows. Combining (4.11) and (4.6), one has

$$|P_{t_2} f(0) - P_{t_1} f(0)| \leq C(A, \delta, \Lambda(f)) e^{-\frac{\delta\Lambda}{2}t_1} \|f\|. \tag{4.12}$$

Step 2: Proving that $\lim_{t \rightarrow \infty} P_t f(x) = \ell(f)$ for all $x \in \mathbb{B}$.

It suffices to prove that the above limit is true for every x in one ball $B_{R, \rho}$. By triangle inequality, one has

$$\begin{aligned}
|P_t f(x) - \ell(f)| & \leq |P_t f(x) - P_t^N f(x)| + |P_t^N f(x) - P_t^N f(0)| \\
& \quad + |P_t^N f(0) - P_t f(0)| + |P_t f(0) - \ell(f)|
\end{aligned} \tag{4.13}$$

By (4.12),

$$|P_t f(0) - \ell(f)| < C e^{-\frac{A\Lambda\delta}{2}t} \|f\|, \tag{4.14}$$

where $C = C(A, \delta, \Lambda(f)) > 0$. By Theorem 2.3, $\forall t > 0, \exists N(t, R, \rho) \in \mathbb{N}$ such that as $N > N(t, R, \rho)$

$$\begin{aligned}
|P_t f(x) - P_t^N f(x)| & < e^{-\frac{A\Lambda\delta}{2}t} \|f\|, \\
|P_t^N f(0) - P_t f(0)| & < e^{-\frac{A\Lambda\delta}{2}t} \|f\|.
\end{aligned} \tag{4.15}$$

By an argument similar as in (4.8)-(4.10), we have

$$\begin{aligned}
|P_t^N f(x) - P_t^N f(0)| &\leq \sum_{i \in \mathbb{Z}^d} \|\partial_i P_t^N f\| |x_i| \\
&\leq C \left[(B^2 t_1^2 + 1 + \Lambda(f))^{\rho+d} e^{-\delta t} + \sum_{i \in \Gamma_N \setminus \Lambda} e^{-An_i - At} (1 + |i|^\rho) \right] \|f\| \\
&\leq C \left[(B^2 t^2 + 1 + \Lambda(f))^{\rho+d} e^{-\frac{\delta}{2} t} + \sum_{i \in \Gamma_N \setminus \Lambda} e^{-An_i - \frac{A}{2} t} (1 + |i|^\rho) \right] e^{-\frac{A\Lambda\delta}{2} t} \|f\|.
\end{aligned} \tag{4.16}$$

Collecting (4.13)-(4.16), we immediately conclude Step 2.

Step 3: Proof of the existence of ergodic measure μ and (2.5).

From step 2, for each $f \in \mathcal{D}^2$, there exists a constant $\ell(f)$ such that

$$\lim_{t \rightarrow \infty} P_t f(x) = \ell(f)$$

for all $x \in \mathbb{B}$. It is easy to see that ℓ is a linear functional on \mathcal{D}^2 , since \mathbb{B} is locally compact (under the product topology), there exists some unsigned Radon measure μ supported on \mathbb{B} such that $\mu(f) = \ell(f)$ for all $f \in \mathcal{D}^2$. By the fact that $P_t \mathbf{1}(x) = 1$ for all $x \in \mathbb{B}$ and $t > 0$, μ is a probability measure.

On the other hand, since $P_t f(x) = P_t^* \delta_x(f)$ and $\lim_{t \rightarrow \infty} P_t f = \mu(f)$, we have $P_t^* \delta_x \rightarrow \mu$ weakly and μ is strongly mixing. Moreover, by (4.13)-(4.16), we immediately have

$$|P_t f(x) - \mu(f)| \leq C(A, \delta, x, \Lambda(f)) e^{-\frac{A\Lambda\delta}{2} t} \|f\|,$$

recall that $0 < A \leq 1/4$ in 2 of Lemma 3.2 and take $A = 1/4$ in the above inequality, we immediately conclude the proof of (2.5). \square

References

- [1] S. Albeverio, Yu. G. Kondratiev, and M. Röckner, *Uniqueness of the stochastic dynamics for continuous spin systems on a lattice*, J. Funct. Anal. **133** (1995), no. 1, 10–20. MR1351639
- [2] S. Albeverio, Yu. G. Kondratiev, and T. V. Tsikalenko, *Stochastic dynamics for quantum lattice systems and stochastic quantization. I. Ergodicity*, Random Oper. Stochastic Equations **2** (1994), no. 2, 103–139, Translated by the authors. MR1293068
- [3] S. Albeverio, V. Mandrekar, and B. Rüdiger, *Existence of mild solutions for stochastic differential equations and semilinear equations with non-Gaussian Lévy noise*, Stochastic Process. Appl. **119** (2009), no. 3, 835–863. MR2499860

- [4] Sergio Albeverio, Barbara Rüdiger, and Jiang-Lun Wu, *Invariant measures and symmetry property of Lévy type operators*, Potential Anal. **13** (2000), no. 2, 147–168. MR1782254
- [5] Sergio Albeverio, Jiang-Lun Wu, and Tu-Sheng Zhang, *Parabolic SPDEs driven by Poisson white noise*, Stochastic Process. Appl. **74** (1998), no. 1, 21–36. MR1624076
- [6] Dominique Bakry and Michel Émery, *Inégalités de Sobolev pour un semi-groupe symétrique*, C. R. Acad. Sci. Paris Sér. I Math. **301** (1985), no. 8, 411–413. MR808640
- [7] Richard F. Bass and Zhen-Qing Chen, *Systems of equations driven by stable processes*, Probab. Theory Related Fields **134** (2006), no. 2, 175–214. MR2222382
- [8] Jean Bertoin, *Lévy processes*, Cambridge Tracts in Mathematics, vol. 121, Cambridge University Press, Cambridge, 1996. MR1406564
- [9] Klaus Bichteler, *Stochastic integration with jumps*, Encyclopedia of Mathematics and its Applications, vol. 89, Cambridge University Press, Cambridge, 2002. MR1906715
- [10] Anton Bovier, *An introduction to aging*: <http://www-wt.iam.uni-bonn.de/~bovier/files/bonn.pdf>, 2007, preprint.
- [11] Giuseppe Da Prato and Jerzy Zabczyk, *Stochastic equations in infinite dimensions*, Encyclopedia of Mathematics and its Applications, vol. 44, Cambridge University Press, Cambridge, 1992.
- [12] Giuseppe Da Prato and Jerzy Zabczyk, *Ergodicity for infinite-dimensional systems*, London Mathematical Society Lecture Note Series, vol. 229, Cambridge University Press, Cambridge, 1996.
- [13] Don Dawson, *Stochastic population dynamics*, (2009), <http://www.math.ubc.ca/~db5d/SummerSchool09/lectures-dd/lecture14.pdf>.
- [14] Gerald B. Folland, *Real analysis*, second ed., Pure and Applied Mathematics (New York), John Wiley & Sons Inc., New York, 1999, Modern techniques and their applications, A Wiley-Interscience Publication. MR1681462
- [15] Tadahisa Funaki and Bin Xie, *A stochastic heat equation with the distributions of Lévy processes as its invariant measures*, Stochastic Process. Appl. **119** (2009), no. 2, 307–326. MR2493992
- [16] A. Greven and F. den Hollander, *Phase transitions for the long-time behavior of interacting diffusions*, Ann. Probab. **35** (2007), no. 4, 1250–1306. MR2330971
- [17] A. Guionnet and B. Zegarlinski, *Lectures on logarithmic Sobolev inequalities*, Séminaire de Probabilités, XXXVI, Lecture Notes in Math., vol. 1801, Springer, Berlin, 2003, pp. 1–134. MR1971582
- [18] Martin Hairer and Jonathan C. Mattingly, *Ergodicity of the 2D Navier-Stokes equations with degenerate stochastic forcing*, Ann. of Math. (2) **164** (2006), no. 3, 993–1032. MR2259251
- [19] M. Hutzenthaler and A. Wakolbinger, *Ergodic behavior of locally regulated branching populations*, Ann. Appl. Probab. **17** (2007), no. 2, 474–501. MR2308333

- [20] Nobuyuki Ikeda and Shinzo Watanabe, *Stochastic differential equations and diffusion processes*, North-Holland Mathematical Library, vol. 24, North-Holland Publishing Co., Amsterdam, 1981. MR637061
- [21] Carlo Marinelli and Michael Röckner, *Well-posedness and asymptotic behavior for stochastic reaction-diffusion equations with multiplicative poisson noise*, arXiv:0903.3299v2.
- [22] Bernt Øksendal, *Stochastic partial differential equations driven by multi-parameter white noise of Lévy processes*, Quart. Appl. Math. **66** (2008), no. 3, 521–537. MR2445527
- [23] Robert Olkiewicz, Lihu Xu, and Bogusław Zegarliński, *Nonlinear problems in infinite interacting particle systems*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **11** (2008), no. 2, 179–211. MR2426714
- [24] S. Peszat and J. Zabczyk, *Stochastic partial differential equations with Lévy noise*, Encyclopedia of Mathematics and its Applications, vol. 113, Cambridge University Press, Cambridge, 2007, An evolution equation approach. MR2356959
- [25] Szymon Peszat and Jerzy Zabczyk, *Stochastic heat and wave equations driven by an impulsive noise*, Stochastic partial differential equations and applications—VII, Lect. Notes Pure Appl. Math., vol. 245, Chapman & Hall/CRC, Boca Raton, FL, 2006, pp. 229–242. MR2227232
- [26] Martijn R. Pistorius, *An excursion-theoretical approach to some boundary crossing problems and the Skorokhod embedding for reflected Lévy processes*, Séminaire de Probabilités XL, Lecture Notes in Math., vol. 1899, Springer, Berlin, 2007, pp. 287–307. MR2409012
- [27] E. Priola and J. Zabczyk, *Structural properties of semilinear spdes driven by cylindrical stable processes*, Probab. Theory Related Fields (2009), (to appear).
- [28] Enrico Priola and Jerzy Zabczyk, *Densities for Ornstein-Uhlenbeck processes with jumps*, Bull. Lond. Math. Soc. **41** (2009), no. 1, 41–50. MR2481987
- [29] L. C. G. Rogers and David Williams, *Diffusions, Markov processes, and martingales. Vol. 1*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2000, Foundations, Reprint of the second (1994) edition. MR1796539
- [30] Marco Romito and Lihu Xu, *Ergodicity of the 3d stochastic navier-stokes equations driven by mildly degenerate noise*, 2009, preprint.
- [31] Hermann Rost, *Skorokhod's theorem for general Markov processes*, Transactions of the Sixth Prague Conference on Information Theory, Statistical Decision Functions, Random Processes (Tech. Univ. Prague, Prague, 1971; dedicated to the memory of Antonín Špaček), Academia, Prague, 1973, pp. 755–764. MR0356248
- [32] Lihu Xu, *Stochastic population dynamics*, Lecture notes based on the lectures given by Don Dawson at summer school of UBC in 2009 (pp 44, still being writing in progress).
- [33] Lihu Xu and Bogusław Zegarliński, *Ergodicity of the finite and infinite dimensional α -stable systems*, Stoch. Anal. Appl. **27** (2009), no. 4, 797–824. MR2541378

- [34] Lihu Xu and Boguslaw Zegarlinski, *Existence and ergodicity of infinite white α -stable systems with unbounded interactions*, arXiv:0911.2866, 2009.
- [35] Boguslaw Zegarlinski, *The strong decay to equilibrium for the stochastic dynamics of unbounded spin systems on a lattice*, *Comm. Math. Phys.* **175** (1996), no. 2, 401–432. MR1370101