

Solution to Problem 65-7*: Solution to an integral equation

Citation for published version (APA):

Bouwkamp, C. J. (1966). Solution to Problem 65-7*: Solution to an integral equation. *SIAM Review*, 8(3), 393-395. <https://doi.org/10.1137/1008085>

DOI:

[10.1137/1008085](https://doi.org/10.1137/1008085)

Document status and date:

Published: 01/01/1966

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

*Problem 65-6, A Ratio of Two Definite Multiple Integrals,** by A. J. STRECOK (Argonne National Laboratory).

Determine

$$I(-1, x)/I(1, x),$$

where

$$I(s, x) = \int_0^x \int_0^t t(x^2 - t^2)^{-1/2} e^{s(t^2 - u^2)} du dt$$

for nonzero s and x .

Solution by P. J. SHORT (White Sands Missile Range).

The solution here can be obtained from the solution to *Problem 64-8* (October, 1965). Denoting the integral in *Problem 64-8* by $I^*(x)$, it follows by making the substitutions $v = u\sqrt{s}$ and $w = t\sqrt{s}$ that

$$I(s, x) = \frac{\sqrt{\pi}}{2s} I^*(x\sqrt{s}) = \frac{\pi}{4s} \{e^{sx^2} - 1\}.$$

Whence,

$$\frac{I(-1, x)}{I(1, x)} = e^{-x^2}.$$

Also solved by R. D. ADAMS (University of Kansas), D. BEIN (Fairleigh Dickinson University), C. J. BOUWKAMP (Technological University, Eindhoven, Netherlands), M. P. FRIEDMAN (Smithsonian Institute), D. L. LANSING (NASA—Langley Research Center), C. B. A. PECK (two solutions) (Pennsylvania State College), H. B. ROSENSTOCK (U. S. Naval Research Laboratory), L. RUBENFELD (Courant Institute of Mathematical Sciences), P. A. SCHEINOK (Hahnemann Medical College), C. B. SHAW, JR. (Electro-Optical Systems, Inc.), S. SPITAL (California State Polytechnic College), F. W. STEUTEL (Technische Hogeschool Twente, Enschede, Netherlands) and the proposer.

Problem 65-7, Solution to an Integral Equation,†* by A. J. STRECOK (Argonne National Laboratory).

Show that a solution of the equation

$$(1) \quad 1 = e^{-x^2} + \int_0^x 2te^{-t^2} \psi(x, t) dt$$

is given by

$$(2) \quad \psi(x, t) = \frac{2}{\pi} (x^2 - t^2)^{-1/2} \int_0^t e^{u^2} du,$$

and give conditions on $\psi(x, t)$ which make this a unique solution.

* Work performed under the auspices of the United States Atomic Energy Commission.

† Work performed under the auspices of the United States Atomic Energy Commission.

Solution by C. J. BOUWKAMP (Technological University, Eindhoven, Netherlands).

The first part of the problem is simple in view of the solution given of *Problem 65-6* if applied to $s = -1$, viz.,

$$\int_0^x 2te^{-t^2}\psi(x, t) dt = \frac{4}{\pi} I(-1, x) = 1 - e^{-x^2}, \quad x > 0.$$

The second part of the problem is not well posed. Equation (1) is not a proper integral equation. The functional equation (1) has an abundance of solutions, and conditions on $\psi(x, t)$ that will make the particular solution (2) the only solution of (1) seem to be quite artificial. This is substantiated by noting it is easy to construct solutions of the types $f(t)g(x)$ and $f(t) + g(x)$, where $f(t)$ is wholly arbitrary. Thus,

$$\psi_1(x, t) = f(t)(1 - e^{-x^2}) \left(\int_0^x 2te^{-t^2}f(t) dt \right)^{-1}$$

and

$$\psi_2(x, t) = 1 + f(t) - (1 - e^{-x^2})^{-1} \int_0^x 2te^{-t^2}f(t) dt$$

are solutions of (1) for any $f(t)$ making the right-hand side meaningful. Apparently, the proposer is interested in solutions that are singular at $x = t$. However, there is a whole class of such solutions. This is shown as follows.

Assume that (1) is meant to hold for $x > 0$, and that $\psi(x, t)$ is defined and continuous for $0 \leq t < x$. It is somewhat easier to discuss the problem in transformed variables. To that end, set

$$x^2 = \xi, \quad t^2 = \tau, \quad \chi(\xi, \tau) = \psi(x, t).$$

Then (1) becomes

$$(3) \quad 1 = e^{-\xi} + \int_0^\xi e^{-\tau} \chi(\xi, \tau) d\tau,$$

and the function (2) corresponds to

$$(4) \quad \chi(\xi, \tau) = \frac{1}{\pi} (\xi - \tau)^{-1/2} \int_0^\tau e^u u^{-1/2} du,$$

this being a particular solution of (3). In fact, by invoking the theory of Abel's integral equation, this can be proved independently of the solution to *Problem 65-6*.

Now assume that

$$\chi(\xi, \tau) = (\xi - \tau)^{-\alpha} e^\tau g(\tau),$$

where $g(t)$ is an unknown function and $0 < \alpha < 1$. Then (4) transforms into the generalized Abel equation

$$\int_0^\xi \frac{g(\tau)}{(\xi - \tau)^\alpha} d\tau = 1 - e^{-\xi},$$

which is solved by

$$g(\tau) = \frac{e^{-\tau} \sin(\alpha\pi)}{\pi} \int_0^\tau e^u u^{\alpha-1} du.$$

That is to say, in terms of the original variables, the function

$$(5) \quad \psi_3(x, t) = \frac{2 \sin(\alpha\pi)}{\pi(x^2 - t^2)^\alpha} \int_0^t e^{u^2} u^{2\alpha-1} du$$

is a particular solution of (1) for any α with $0 < \alpha < 1$. The proposer's solution (2) is obtained for $\alpha = 1/2$.

Of course, there exist still other solutions of (1). Thus, if the proposer writes his unknown function $\psi(x, t)$ as a product of the kernel function $(x^2 - t^2)^{-1/2}$ and an unknown function $h(t)$, the solution indicated for $h(t)$ is unique, by invoking the existing theory of Abel's integral equation. But then the kernel function should have been prescribed.

Also solved by R. D. ADAMS (University of Kansas), M. P. FRIEDMAN (Smithsonian Institution), F. W. STEUTEL (Technische Hogeschool Twente, Enschede, Netherlands) and P. P. WANG (Bell Telephone Laboratories).