

Greater male variability in daily energy expenditure develops through puberty

Citation for published version (APA):

International Atomic Energy Agency Doubly Labeled Water database group, Halsey, L. G., & Bouten, C. V. C. (2023). Greater male variability in daily energy expenditure develops through puberty. *Biology Letters*, 19(9), Article 20230152. <https://doi.org/10.1098/rsbl.2023.0152>

Document license:
TAVERNE

DOI:
[10.1098/rsbl.2023.0152](https://doi.org/10.1098/rsbl.2023.0152)

Document status and date:
Published: 01/09/2023

Document Version:
Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Research



Cite this article: Halsey LG *et al.* 2023

Greater male variability in daily energy expenditure develops through puberty. *Biol. Lett.* **19**: 20230152.

<https://doi.org/10.1098/rsbl.2023.0152>

Received: 19 April 2023

Accepted: 31 August 2023

Received: 19 April 2023

Accepted: 31 August 2023

Subject Category:

Evolutionary biology

Subject Areas:

evolution

Keywords:

inter-individual variation, morphometry, age, height, weight

Author for correspondence:

Lewis G. Halsey

e-mail: l.halsey@roehampton.ac.uk

[†]IAEA DLW Database Consortium members are listed in the supplementary materials.

[‡]Co-corresponding authors.

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6832989>.

Greater male variability in daily energy expenditure develops through puberty

Lewis G. Halsey¹, Vincent Careau², Philip N. Ainslie³, Heliodoro Alemán-Mateo⁴, Lene F. Andersen⁵, Liam J. Anderson^{3,6}, Leonore Arab⁷, Issad Baddou⁸, Linda Bandini⁹, Kweku Bedu-Addo¹⁰, Ellen E. Blaak¹¹, Stephane Blanc¹², Alberto G. Bonomi¹³, Carlijn V. C. Bouten¹⁴, Pascal Bovet¹⁵, Soren Brage¹⁶, Maciej S. Buchowski¹⁷, Nancy Butte¹⁸, Stephan G. Camps¹⁹, Regian Casper²⁰, Graeme L. Close³, Lisa H. Colbert²¹, Jamie A. Cooper²², Richard Cooper²³, Prasangi Dabare²⁴, Sai Krupa Das²⁵, Peter S. W. Davies²⁶, Sanjoy Deb^{27,28}, Christine Delisle Nyström²⁹, William Dietz³⁰, Lara R. Dugas^{23,31}, Simon Eaton³², Ulf Ekelund³³, Asmaa El Hamdouchi⁸, Sonja Entringer^{34,35}, Terrence Forrester³⁶, Barry W. Fudge³⁷, Melanie Gillingham³⁸, Annelies H. Goris¹⁹, Michael Gurven³⁹, Hinke Haisma⁴⁰, Catherine Hambly⁴¹, Daniel Hoffman⁴², Marije B. Hoos¹⁹, Sumei Hu⁴³, Noorjehan Joonas⁴⁴, Annemiek Joosen¹⁹, Peter Katzmarzyk⁴⁵, Kitty P. Kempen¹⁹, Misaka Kimura⁴⁶, William E. Kraus⁴⁷, Wantanee Kriengsinyos⁴⁸, Rebecca Kuriyan⁴⁹, Robert F. Kushner⁵⁰, Estelle V. Lambert⁵¹, Pulani Lanerolle⁵², Christel L. Larsson⁵³, Nader Lessan⁵⁴, Marie Lof²⁸, Corby Martin⁴⁵, Eric Matsiko⁵⁵, Gerwin A. Meijer¹¹, James C. Morehen^{3,56}, James P. Morton³, Aviva Must⁵⁷, Marian Neuheuser⁵⁸, Theresa A. Nicklas¹⁸, Robert M. Ojiambo^{59,60}, Kirsi H. Pietilainen⁶¹, Yannis P. Pitsiladis⁶², Jacob Plange-Rhule¹⁰, Guy Plasqui⁶³, Ross L. Prentice⁶³, Roberto Rabinovich⁶⁴, Susan B. Racette⁶⁵, David A. Raichen⁶⁶, Eric Ravussin⁴⁵, Leanne Redman⁴⁵, John J. Reilly⁶⁷, Rebecca Reynolds⁶⁸, Susan Roberts²⁵, Jennifer C. Rood⁴⁵, Dulani Samaranyake⁶⁹, Luís B. Sardinha⁷⁰, Albertine J. Scutt⁷¹, Analiza M. Silva⁷⁰, Srishti Sinha⁷², Anders M. Sjödin⁷³, Eric Stice⁷⁴, Albert Stunkard⁷⁵, Samuel S. Urlacher⁷⁶, Mauro Eduardo Valencia⁴, Giulio Valenti¹⁹, Ludo M. van Etten¹⁹, Edgar A. Van Mil⁷⁷, Jeanine A. Verbunt¹⁹, Jonathan C. K. Wells⁷⁸, George Wilson³, Brian Wood^{41,79,80}, Tsukasa Yoshida⁸¹, Xueying Zhang^{82,83,84}, Alexia Murphy-Alford^{85,‡}, Cornelia Loechl^{85,‡}, Amy Luke^{31,86,‡}, Herman Pontzer^{87,88,‡}, Jennifer Rood^{45,‡}, Hiroyuki Sagayama^{89,‡}, Klaas R. Westerterp^{11,‡}, William W. Wong^{18,‡}, Yosuke Yamada^{46,81,‡}, John R. Speakman^{86,87,‡} and IAEA DLW Database Consortium[†]

¹School of Life and Health Sciences, University of Roehampton, London SW15 4JD, UK

²Department of Biology, University of Ottawa, Ottawa, Ontario, Canada

³Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK


⁴Centro de Investigación en Alimentación y Desarrollo, A.C., Mexico

⁵Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, 0317 Oslo, Norway

⁶School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK

⁷David Geffen School of Medicine, University of California, Los Angeles, CA, USA

- ⁸Unité Mixte de Recherche en Nutrition et Alimentation, CNESTEN-Université Ibn Tofail URAC39, Morocco
- ⁹University of Massachusetts Chan Medical School, Worcester, MA, USA
- ¹⁰Department of Physiology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana
- ¹¹Nutrition and Translational Research in Metabolism (NUTRIM), Maastricht University Medical Centre, Maastricht, Netherlands
- ¹²Institut Pluridisciplinaire Hubert Curien, CNRS Université de Strasbourg, Strasbourg, France
- ¹³Phillips Research, Eindhoven, The Netherlands
- ¹⁴Department of Biomedical Engineering and Institute for Complex Molecular Systems, Eindhoven University of Technology, Eindhoven, The Netherlands
- ¹⁵University Center for primary care and public health (Unisante), Lausanne, Switzerland
- ¹⁶MRC Epidemiology Unit, University of Cambridge, Cambridge, UK
- ¹⁷Division of Gastroenterology, Hepatology and Nutrition, Department of Medicine, Vanderbilt University, Nashville, TN, USA
- ¹⁸Department of Pediatrics, Baylor College of Medicine, USDA/ARS Children's Nutrition Research Center, Houston, TX, USA
- ¹⁹Maastricht University, Maastricht, The Netherlands
- ²⁰Department of Psychiatry, University of Chicago, IL, USA
- ²¹Kinesiology, University of Wisconsin, Madison, WI, USA
- ²²Nutritional Sciences, University of Wisconsin, USA
- ²³Department of Public Health Sciences, Parkinson School of Health Sciences and Public Health, Maywood, IL, USA
- ²⁴Department of Physiotherapy, Faculty of Allied Health Sciences, General Sir John Kotelawala Defence University, Sri Lanka
- ²⁵Jean Mayer USDA Human Nutrition Research Center on Aging, Tufts University, Boston, MA, USA
- ²⁶Child Health Research Centre, Level 6 Centre for Children's Health Research, University of Queensland, 62 Graham Street, South Brisbane, Queensland
- ²⁷Centre for Nutraceuticals, School of Life Sciences, University of Westminster, London, UK
- ²⁸Department of Circulation and Medical Imaging, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway
- ²⁹Department of Biosciences and Nutrition, Karolinska Institute, Stockholm, Sweden
- ³⁰George Washington University, Washington, DC, USA
- ³¹Division of Epidemiology and Biostatistics, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa
- ³²UCL, Great Ormond Street Institute of Child Health, London, UK
- ³³Department of Sport Medicine, Norwegian School of Sport Sciences, Oslo, Norway
- ³⁴Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health (BIH), Institute of Medical Psychology, Berlin, Germany
- ³⁵University of California Irvine, Irvine, CA, USA
- ³⁶Solutions for Developing Countries, University of the West Indies, Mona, Kingston, Jamaica
- ³⁷University of Glasgow, Glasgow, UK
- ³⁸Department of Molecular and Medical Genetics, Oregon Health & Science University, Portland, OR
- ³⁹Department of Anthropology, University of California Santa Barbara, Santa Barbara, CA, USA
- ⁴⁰Population Research Centre, Faculty of Spatial Sciences, University of Groningen, The Netherlands
- ⁴¹Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, UK
- ⁴²Department of Nutritional Sciences, Program in International Nutrition, Rutgers University, New Brunswick, NJ, USA
- ⁴³Institute of Genetics and development Biology, Chinese Academy of Sciences, Beichen Xi lu, Beijing, People's Republic of China
- ⁴⁴Central health Laboratory, Ministry of Health and Wellness, Port Louis, Mauritius
- ⁴⁵Pennington Biomedical Research Center, Baton Rouge, LA, USA
- ⁴⁶Institute for Active Health, Kyoto University of Advanced Science, Kyoto, Japan
- ⁴⁷Department of Medicine, Duke University, Durham, NC, USA
- ⁴⁸Institute of Nutrition, Mahidol University, Salaya, Thailand
- ⁴⁹Division of Nutrition, St John's Research Institute, Bangalore, India
- ⁵⁰Northwestern University, Chicago, IL, USA
- ⁵¹Health through Physical Activity, Lifestyle and Sport Research Centre, Division of Exercise Science and Sports Medicine (ESSM), FIMS International Collaborating Centre of Sports Medicine, Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa
- ⁵²Department of Biochemistry and Molecular Biology, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka
- ⁵³University of Gothenburg, Gothenburg, Sweden
- ⁵⁴Imperial College London Diabetes Centre, Abu Dhabi, United Arab Emirates
- ⁵⁵Department of Human Nutrition and Dietetics, University of Rwanda, Kigali, Rwanda
- ⁵⁶The FA Group, Burton-Upon-Trent, Staffordshire, UK
- ⁵⁷Tufts University School of Medicine, Boston, USA
- ⁵⁸Division of Public Health Sciences, Fred Hutchinson Cancer Research Center and School of Public Health, University of Washington, Seattle, WA, USA
- ⁵⁹Moi University, Eldoret, Kenya
- ⁶⁰University of Global Health Equity, Rwanda
- ⁶¹Helsinki University Central Hospital, Helsinki, Finland
- ⁶²University of Brighton, Eastbourne, UK
- ⁶³Department of Nutrition and Movement Sciences, Maastricht University, Maastricht, The Netherlands
- ⁶⁴University of Edinburgh, Edinburgh, UK
- ⁶⁵Arizona State University, Phoenix, AZ, USA
- ⁶⁶Biological Sciences and Anthropology, University of Southern California, CA, USA
- ⁶⁷Physical activity for Health group, School of Psychological Sciences and Health, University of Strathclyde, Glasgow, UK
- ⁶⁸Centre for Cardiovascular Sciences, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK
- ⁶⁹Department of Community Medicine, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka
- ⁷⁰Exercise and health laboratory, CIPER, Faculdade Motricidade Humana, Universidade de Lisboa, Portugal
- ⁷¹University of Wageningen, Wageningen, The Netherlands
- ⁷²Division of Nutrition, St John's Research Institute, Bangalore, India
- ⁷³Department of Nutrition, Exercise and Sports, Copenhagen University, Copenhagen, Denmark
- ⁷⁴Stanford University, Stanford CA, USA
- ⁷⁵University of Pennsylvania Perelman School of Medicine Department of Psychiatry
- ⁷⁶Department of Anthropology, Baylor University, Waco, TX, USA
- ⁷⁷Maastricht and Lifestyle Medicine Center for Children, Jeroen Bosch Hospital 's-Hertogenbosch, Maastricht University, Maastricht, The Netherlands
- ⁷⁸Population, Policy and Practice Research and Teaching Department, UCL Great Ormond Street Institute of Child Health, London, UK
- ⁷⁹University of California Los Angeles, Los Angeles, USA
- ⁸⁰Department of Human Behavior, Ecology, and Culture, Max Planck Institute for Evolutionary Anthropology
- ⁸¹National Institute of Health and Nutrition, National Institutes of Biomedical Innovation, Health and Nutrition, Tokyo, Japan
- ⁸²Center for Energy Metabolism and Reproduction, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, People's Republic of China
- ⁸³China medical University, Shenyang, Liaoning, People's Republic of China
- ⁸⁴CAS Center of Excellence in Animal Evolution and Genetics, Kunming, People's Republic of China
- ⁸⁵Nutritional and Health Related Environmental Studies Section, Division of Human Health, International Atomic Energy Agency, Vienna, Austria
- ⁸⁶Division of Epidemiology, Department of Public Health Sciences, Loyola University School of Medicine, Maywood IL, USA
- ⁸⁷Department of Evolutionary Anthropology, Duke University, Durham, NC, USA
- ⁸⁸Duke Global Health Institute, Duke University, Durham, NC, USA
- ⁸⁹Faculty of Health and Sport Sciences, University of Tsukuba, Ibaraki, Japan

 LGH, 0000-0002-0786-7585; VC, 0000-0002-2826-7837; EEB, 0000-0002-2496-3464; JRS, 0000-0002-2457-1823

There is considerably greater variation in metabolic rates between men than between women, in terms of basal, activity and total (daily) energy expenditure (EE). One possible explanation is that EE is associated with male sexual characteristics (which are known to vary more

than other traits) such as musculature and athletic capacity. Such traits might be predicted to be most prominent during periods of adolescence and young adulthood, when sexual behaviour develops and peaks. We tested this hypothesis on a large dataset by comparing the amount of male variation and female variation in total EE, activity EE and basal EE, at different life stages, along with several morphological traits: height, fat free mass and fat mass. Total EE, and to some degree also activity EE, exhibit considerable greater male variation (GMV) in young adults, and then a decreasing GMV in progressively older individuals. Arguably, basal EE, and also morphometrics, do not exhibit this pattern. These findings suggest that single male sexual characteristics may not exhibit peak GMV in young adulthood, however total and perhaps also activity EE, associated with many morphological and physiological traits combined, do exhibit GMV most prominently during the reproductive life stages.

1. Introduction

Individuals of a sexually reproducing species vary in terms of nearly every measurable characteristic, morphological, physiological and cognitive. In mammals, often the magnitude of this inter-individual variability, at least in terms of body morphometrics and, in humans, also cognition, has been reported as greater between males than between females [1,2]. This phenomenon is termed ‘greater male variability’ (GMV) [3]. For example, it has been reported that human males are more varied in their physical performance than are human females [4,5], and male chimpanzees have more varied brain structures than do female chimpanzees [6]. Given that the metabolism of an animal is arguably an emergent property influenced by the culmination of GMV in various bodily traits, in a previous study we postulated that GMV in energy expenditure could be particularly large [7]. Supporting this suggestion, we found that in adult humans, energy expenditure exhibits considerable GMV in terms of basal energy expenditure (BEE) and total (daily) expenditures (TEE), even after controlling for key morphological traits such as height, fat free mass and fat mass [7].

We also discovered that with ageing, variation between people in their TEE decreases, but this happens more rapidly in men than in women which results in the magnitude of GMV in TEE attenuating in older age groups [7]. One possible explanation for GMV is that males experiencing stronger sexual selection results in the expression of male sexual traits with greater variance than that of female sexual traits [8]. Energy expenditure is related to various traits of sexual interest to females such as cognitive capacity [9], physical endeavour, strength and muscle mass [10], and perhaps also aerobic fitness [11]. Variability in male sexual characteristics could be most prominent during the period of life associated with sexual reproduction (i.e. late adolescence and young adulthood), explaining the aforementioned decrease in GMV in energy expenditure in older people.

If indeed GMV in energy expenditure is underpinned by sexual selection, we predict there will be no GMV in humans prior to sexual maturity, and that GMV will peak during young adulthood due to an increase in male variation,

which will then decrease more rapidly than female variation into old age. We tested this hypothesis by analysing extensive data on human energy expenditure for individuals of ages spanning the entire human life course.

2. Methods

We analysed data from the International Atomic Energy Agency (IAEA) DLW database v.3.7 [12]. The dataset comprised TEE measurements (MJ d^{-1}) for 4992 females and 2626 males, and BEE measurements (MJ d^{-1}) using indirect calorimetry for 1542 females and 934 males. Estimates of activity energy expenditure (AEE) were calculated by subtracting BEE from $0.9 \times \text{TEE}$ (TEE adjusted to account for the thermic effect to food). Ages of the participants ranged from newborns to 101 years old. We also included in our analyses the following traits: height (cm), fat-free mass (kg), fat mass (kg) and body weight (kg). Further details are provided in Halsey *et al.* [7].

(a) Statistical analyses

All analyses were conducted in R v. 3.5.3 [13]. We quantified variance in male and female energy expenditure using Bayesian general linear models based on Monte-Carlo Markov chain (MCMC) models using the ‘MCMCglmm’ package [14]. Each model included one trait as the response variable (TEE, AEE or BEE) and sex as an independent variable along with age as a categorical variable, and height, fat-free mass and fat mass as continuous variables, with no intercept fitted so that the model returned separate mean estimates for males and females within each age category [15]. Age was recoded as a categorical variable with eight levels for the TEE model, and six levels for the AEE and BEE models (due to a smaller overall BEE sample size, although sample size per age category was still smaller). Using the ‘idh’ function, we allowed the residual variance to be different in males and females within each age category. Further models included height, fat-free mass or fat mass as the response variable (with only sex and age as independent variables). All models also included country as a random effect to account for the unequal sampling distribution across countries. When included as covariates, the three morphometric variables were centred so that model estimates were estimated at the centre of the distribution of the covariates [15]. To make sure that any sex differences in variance were not due to mean differences, we also standardized the variance within each sex and age category by dividing variances with the sex-specific mean estimates to obtain the coefficient of variance (CoV). From each model, we calculated the posterior mode and 95% highest posterior density credible intervals (CIs) for the CoV estimates for each age group in males and females. Treating these as 95% confidence intervals, in cases where the CIs for the relative variances of the two sexes do not overlap, the evidence of a difference in variance between the sexes is strong [16]. Finally, we calculated the male : female ratio of CoV whereby a value greater than 1 indicated GMV.

3. Results

Visual interpretation of graphs of energy expenditure and morphology against age category (figure 1), using the 95% CIs for guidance, indicates that TEE and AEE exhibit increases in male variance from early childhood into young adulthood followed by decreases with further ageing. In TEE, this generates the development of GMV into young adulthood and a decrease in the elderly, while in AEE a pattern of increase and then decrease is apparent but less clear. A pattern in the data for BEE is not apparent although




Figure 1. Top row: male (blue triangles) and female (red circles) coefficients of variance (CoV) in measures of energy expenditure and body morphometrics as a function of age category. Error bars represent 95% credible intervals (95% CI). These data are presented in electronic supplementary material, table S1. Bottom row: the ratio of male to female (M/F) CoV with 95% CI. The dashed lines represent equal variance; values above indicate greater male variance (GMV). TEE = total energy expenditure; AEE = activity energy expenditure; BEE = basal energy expenditure.

GMV does peak in early adulthood. Average height shows no pattern of GMV developing into young adulthood and then fading. Fat free mass also shows no such pattern. Fat mass does exhibit an increase in male variance followed by a decrease, which results in a similar pattern in GMV, however peak GMV occurs well before adulthood.

4. Discussion

In young children, there is little difference between boys and girls in terms of between-person variance in TEE (figure 1). However, by late adolescence (15–19 years) males are exhibiting greater variation and in turn GMV. This GMV increases in magnitude into young adulthood and beyond, and then somewhat decreases by old age due to decreasing inter-male variation. This pattern through the life course bears hallmarks of a sexually selected signal, or at least a trait that is an emergent property of characteristics some or all of which are sexually selected. The qualitative pattern of increasing then decreasing inter-male variation is also apparent although less strong in AEE, and increasing then decreasing GMV is also apparent. Albeit the sample size was lower and inter-study measurement variation probably higher, the above findings together tentatively suggest that changes in AEE variance over the life course might somewhat drive the TEE variance patterns.

Patterns for BEE are less clear and might be due to, again, a lower sample size than for TEE and with greater measurement variation between studies. More data are required.

Given that body morphology is considered to harbour sexual traits, we might predict that at least some morphological measures would exhibit a pattern of GMV similar to that seen in TEE, however this was not the case. While fat mass exhibits an increase then decrease in inter-male variation over the life course, the start of peak GMV occurs before adolescence rather than during the age of peak reproduction. Average height and fat free mass both show a spike in inter-male variation, and in turn substantive GMV, for a short period in early adolescence, an observation also apparent in data for Dutch children [17]. However, this is likely generated by the variability in both magnitude and degree of growth spurt in boys at this age.

About two-thirds of the dataset are for citizens of the USA, a country that has particularly high rates of obesity (<https://www.oecd.org/health/health-data.htm>). Nonetheless, qualitatively the patterns in male variation, female variation and GMV through the life course are generally similar to those present in the full dataset when the USA data are removed (electronic supplementary material, figure S1 and table S2).

In conclusion, inter-individual variance over the life course for height and fat free mass do not exhibit the hypothesized patterns given that they are two morphometric variables considered to be male sexual characteristics (albeit height is a relatively fixed variable come adulthood). Fat mass is the morphometric variable that is closest to exhibiting the hypothesized patterns, despite being considered more of a female sexual characteristic. TEE and to some extent AEE present with GMV developing and peaking in young adulthood and subsequently declining, perhaps indicating that they are either male sexual characteristics or related to such characteristics. Perhaps energy expenditure most clearly presents with the hypothesized patterns of GMV over the life course because it results from the combined effects of many characteristics.

Ethics. This work did not require ethical approval from a human subject or animal welfare committee.

Data accessibility. The data are provided in the electronic supplementary material [18].

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. L.G.H.: conceptualization, formal analysis, investigation, project administration, software, visualization, writing—original draft, writing—review and editing; V.C.: formal analysis, software, visualization, writing—review and editing; J.R.S.: data curation, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. The authors also gratefully acknowledge funding from the Chinese Academy of Sciences (grant no. CAS153E11KYSB20190045) to J.R.S. and the US National Science Foundation (grant no. BCS-1824466) awarded to H.P.

Acknowledgements. Yvonne Schönbeck provided important information about morphometric measurements for Dutch children. A chat over

dinner with Karsten Koehler, Eimear Dolan and Danny Longman brought up a number of thoughts that influenced this manuscript. The DLW database, which can be found at <https://doublylabelled-waterdatabase.iaea.org/home>, is hosted by the IAEA and generously

supported by Taiyo Nippon Sanso and SERCON. We are grateful to the IAEA and these companies for their support and especially to Takashi Oono for his tremendous efforts at fundraising on our behalf.

References

- Reinhold K, Engqvist L. 2013 The variability is in the sex chromosomes. *Evolution* **67**, 3662–3668. (doi:10.1111/evo.12224)
- Baye A, Monseur C. 2016 Gender differences in variability and extreme scores in an international context. *Large-Scale Assess. Educ.* **4**, 1. (doi:10.1186/s40536-015-0015-x)
- Darwin, C. 1871 *The descent of man, and selection in relation to sex*. London, UK: John Murray.
- Lehre AC, Lehre KP, Laake P, Danbolt NC. 2009 Greater intrasex phenotype variability in males than in females is a fundamental aspect of the gender differences in humans. *Dev. Psychobiol.* **51**, 198–206. (doi:10.1002/dev.20358)
- Wilmore JH. 1974 Alterations in strength, body composition and anthropometric measurements consequent to a 10-week weight training program. *Med. Sci. Sports Exerc.* **6**, 133–138.
- DeCasien AR, Sherwood CC, Schapiro SJ, Higham JP. 2020 Greater variability in chimpanzee (*Pan troglodytes*) brain structure among males. *Proc. R. Soc. B* **287**, 20192858. (doi:10.1098/rspb.2019.2858)
- Halsey LG *et al.* 2022 Variability in energy expenditure is much greater in males than females. *J. Hum. Evol.* **171**, 103229. (doi:10.1016/j.jhevol.2022.103229)
- Harrison LM, Noble DW, Jennions MD. 2022 A meta-analysis of sex differences in animal personality: no evidence for the greater male variability hypothesis. *Biol. Rev.* **97**, 679–707. (doi:10.1111/brv.12818)
- Gonczewicz A, Górkiewicz T, Dzik JM, Jędrzejewska-Szmek J, Knapska E, Konarzewski M. 2022 Brain size, gut size and cognitive abilities: the energy trade-offs tested in artificial selection experiment. *Proc. R. Soc. B* **289**, 20212747. (doi:10.1098/rspb.2021.2747)
- Lidborg LH, Cross CP, Boothroyd LG. 2022 A meta-analysis of the association between male dimorphism and fitness outcomes in humans. *Elife* **11**, e65031. (doi:10.7554/eLife.65031)
- Poehlman ET, Melby CL, Badylak SF, Calles J. 1989 Aerobic fitness and resting energy expenditure in young adult males. *Metabolism* **38**, 85–90. (doi:10.1016/0026-0495(89)90185-6)
- Speakman JR *et al.* 2019 The International Atomic Energy Agency international Doubly Labelled Water Database: aims, scope and procedures. *Ann. Nutr. Metab.* **75**, 114–118. (doi:10.1159/000503668)
- R Development Core Team. 2011 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Hadfield JD. 2010 MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *J. Stat. Softw.* **33**, 1–22. (doi:10.18637/jss.v033.i02)
- Schielzeth H. 2010 Simple means to improve the interpretability of regression coefficients. *Methods Ecol. Evol.* **1**, 103–113. (doi:10.1111/j.2041-210X.2010.00012.x)
- Belia S, Fidler F, Williams J, Cumming G. 2005 Researchers misunderstand confidence intervals and standard error bars. *Psychol. Methods* **10**, 389. (doi:10.1037/1082-989X.10.4.389)
- Schönbeck Y, Talma H, Van Dommelen P, Bakker B, Buitendijk SE, HiraSing RA, Van Buuren S. 2013 The world's tallest nation has stopped growing taller: the height of Dutch children from 1955 to 2009. *Pediatr. Res.* **73**, 371–377. (doi:10.1038/pr.2012.189)
- Halsey LG *et al.* 2023 Greater male variability in daily energy expenditure develops through puberty. Figshare. (doi:10.6084/m9.figshare.c.6832989)