

Thomas Young's research on fluid transients: 200 years on

Citation for published version (APA):
Tijsseling, A. S., & Anderson, A. (2008). Thomas Young's research on fluid transients: 200 years on. (CASAreport; Vol. 0816). Technische Universiteit Eindhoven.

Document status and date:

Published: 01/01/2008

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

Link to publication

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- · Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Download date: 16. Jun. 2025

EINDHOVEN UNIVERSITY OF TECHNOLOGY

Department of Mathematics and Computer Science

CASA-Report 08-16 April 2008

Thomas Young's research on fluid transients: 200 years on

by

A.S. Tijsseling, A. Anderson



Centre for Analysis, Scientific computing and Applications Department of Mathematics and Computer Science Eindhoven University of Technology P.O. Box 513

5600 MB Eindhoven, The Netherlands

ISSN: 0926-4507

Thomas Young's research on fluid transients: 200 years on

Arris S Tijsseling
Department of Mathematics
and Computer Science
Eindhoven University of Technology
P.O. Box 513, 5600 MB Eindhoven
The Netherlands

Alexander Anderson School of Mechanical and Systems Engineering Newcastle University Newcastle upon Tyne NE1 7RU United Kingdom

ABSTRACT

Thomas Young published in 1808 his famous paper (1) in which he derived the pressure wave speed in an incompressible liquid contained in an elastic tube. Unfortunately, Young's analysis was obscure and the wave speed was not explicitly formulated, so his achievement passed unnoticed until it was rediscovered nearly half a century later by the German brothers Weber.

This paper briefly reviews Young's life and work, and concentrates on his achievements in the area of hydraulics and waterhammer. Young's 1808 paper is "translated" into modern terminology. Young's discoveries, though difficult for modern readers to identify, appear to include most if not all of the key elements which would subsequently be combined into the pressure rise equation of Joukowsky.

Keywords

waterhammer, fluid transients, solid transients, wave speed, history, Thomas Young

NOTATION

c	sonic wave speed, m/s	p	fluid pressure, Pa
D	internal tube diameter, m	R	internal tube radius, m
E	Young's modulus, Pa	t	time, s
e	tube wall thickness, m	v	velocity, m/s
f	elastic limit, Pa	$\boldsymbol{\mathcal{X}}$	length, m
g	gravitational acceleration, m/s ²	δ	change, jump
h	height, pressure head, m	ε	longitudinal strain
K	fluid bulk modulus, Pa	ho	mass density, kg/m ³
K^*	effective bulk modulus, Pa	σ	longitudinal stress, Pa
k	elasticity coefficient, m/Pa		_

INTRODUCTION

By the end of the 19th century, the three key elements for the development of modern waterhammer theory were in place in the seminal works of Joukowsky and Allievi (2):

- an expression for the waterhammer disturbance wave celerity depending on both fluid and pipe wall elasticity (Eq 2 below),
- the Joukowsky formula for the waterhammer pressure rise (Eq 3 below), and
- the functional form of waterhammer solutions depending on the characteristics along which the pressure waves propagated.

The 19th century saw the gradual emergence of these key elements through the work of three principal groups of investigators:

- physiologists interested in haemodynamics (3, 4), e.g. Kries (5) and Galabin (6),
- acousticians interested in the propagation of sound (7), and
- hydrodynamicists, hydraulicians and engineers engaging with practical pipe systems and devices, including the hydraulic ram (from which the term for waterhammer in many languages arises, e.g. the French "coup de bélier" or Italian "colpo d'ariete"), of whom Ménabréa (8) is an early example.

The aim of this paper is to draw attention to the contributions of Thomas Young at the start of the 19th century.

To the modern scientific reader Young's published works can be difficult to follow. He stands at the end of an era when the style of presentation of science in England remained in the tradition exemplified by Newton's Principia, a style with a strong base in Euclidean geometry for its demonstrations and verbal (rather than algebraic) statements of physical laws and mathematical results. As will be shown, Young was one of the last proponents of this style and he became as aware as his European contemporaries of its limitations. Nevertheless, his immediate 19th century successors who built on his achievements seem to have been convinced of these, possibly because they were closer to and thus more familiar with this older style. Notwithstanding his attachment to this archaic mode of presentation, though, it will be argued that Young was a truly innovative scientist in first developing key concepts for what would become, about ninety years later, a recognisable theory of waterhammer (though he himself does not appear to have drawn them all together).

Exactly 200 years ago Thomas Young (1773-1829) published a paper entitled "Hydraulic Investigations, subservient to an intended Croonian Lecture on the Motion of the Blood" (1). In this paper he can be seen to have arrived at the celerity (c) of a pressure wave propagating in an incompressible liquid of mass density ρ contained in an elastic tube with Young's modulus E as

$$c = \sqrt{\frac{E \ e}{\rho \ D}} \tag{1}$$

where e/D is the ratio of wall thickness to tube diameter. This formula is valid for waterhammer in flexible hoses and for the pulse in haemodynamics. It represents "half" of the classical waterhammer wave speed derived by the Dutch mathematician Diederik Korteweg (9) seventy years later as

$$c = \sqrt{\frac{E \ e}{\rho \ D}} / \sqrt{1 + \frac{E \ e}{K \ D}} \tag{2}$$

which takes into account the elasticity K of the liquid within the tube. Unfortunately Young's analysis is obscure to present-day readers and the actual Eq 1 was not written explicitly in his paper, so this achievement (like many others) passed unnoticed until it was rediscovered nearly half a century later by the German brothers Ernst-Heinrich and Wilhelm Weber (10, 11). It has been noted in medical historical reviews, e.g. (3, 4, 12) but overlooked in histories of waterhammer, e.g. (13, 14).

In this respect it is typical that Young had also implicitly derived (but for elastic solids rather than fluids) an equivalent of the Joukowsky equation

$$p = \rho c v \tag{3}$$

that relates pressure (p) to velocity (v) in sound and vibration, in his encyclopaedic book "A Course of Lectures on Natural Philosophy and the Mechanical Arts" (15-17). In addition, he was also an early commentator on the hydraulic ram.

This paper concentrates on Young's achievements in the field of hydraulics and waterhammer. Young's implicit discovery of the Joukowsky equation for solids is discussed. Young's 1808 paper (1) is difficult to read and therefore, following the example of Boulanger (3), his derivation of Eq 1 is "translated" into modern terminology. Finally, the work of Young's immediate successors, who first expressed his Eq 1 in its modern form, is briefly summarised.



Figure 1 Thomas Young in the 1820s.

YOUNG'S LIFE AND WORK

Thomas Young (Figs. 1 and 2) was an intriguing person and scientist. He has inspired many people to write biographical papers in all sorts and sizes (see App. A), mostly of hagiographic nature and none of them highlighting Young's under-appreciated contribution to the theory of fluid transients. Some of the most revealing comments about his style were written by himself (characteristically in third person) in his own "Autobiographical Fragment", rediscovered and published by Hilts (18), e.g. "...and for about two years he was the colleague of Sir Humphrey Davy as a lecturer, though his

language was never either very popular or very fluent, his compressed and laconic style and manner being more adapted for the study of a man of science than for the amusement of a lady of fashion." In view of what follows below, his own comment on one of his papers also deserves notice: "The mathematical reasoning, for want of mathematical symbols, was not understood; from a dislike of the affectation of algebraic calculation which he had observed in the French, the author was led into something like an affection of simplicity, which was equally objectionable." (18).




Figure 2 "Mr Thomas Young, of Little Queen Street, Westminster, a gentleman conversant with various branches of literature and science, and author of a paper on vision published in the Philosophical Transactions." So reads the citation on Young's Royal Society certificate for election.

Of his own work, he similarly wrote (18): "His pursuits, diversified as they were, had all originated in the first instance from the study of physic (i.e. medicine): the eye and the ear led him to the consideration of sound and of light...". The range of his achievements is too extensive to cover in a single paper. A short and incomplete list of his achievements includes, inter alia:

- Young advocated Huygens' wave theory of light as opposed to Newton's particle model; he discovered the principle of (light) wave interference and he invented the double-slit experiment. He made vast progress in the field of optics, an area later fully developed by Fresnel.
- Young discovered that the three primary colours are not a property of light but of the structure of the human eye. His theory of colours was rediscovered fifty years later by Helmholtz and further developed by Maxwell. He discovered the phenomenon of astigmatism.
- Young estimated the size of molecules and blood corpuscles, fifty years before anyone else.
- Young and Laplace independently derived the fundamental equation of surface tension, and Young calculated the contact angle between an adhesive liquid and a solid, an idea elaborated sixty years later by Dupré. He studied the tensile strength of liquids.

- Other investigations by Young include: sound waves and harmonics, tides, visualization techniques (shadows of water waves; wave superposition, foreshadowing Fourier analysis).
- Young led the basis for the deciphering of Egyptian hieroglyphics, a task later accomplished by Champollion.

He was the first to use the term (kinetic) "energy" in the modern sense and he introduced the term "Indo-European" for a large family of related languages. Named after him are: Young's modulus (of elasticity), Young's fringes (of interference patterns), Young's rule (for the dose of medicine), Young's temperament (for keyboard tuning), and Young's mode (of wave propagation). Helmholtz (himself a figure in the development of waterhammer theory) wrote (19): "Young was one of the most acute men who ever lived, but had the misfortune to be too far in advance of his contemporaries. They looked on him with astonishment, but could not follow his bold speculations, and thus a mass of his important thoughts remained buried and forgotten in the Transactions of the Royal Society until a later generation by slow degrees arrives at the rediscovery of his discoveries, and came to appreciate the force of his arguments and the accuracy of his conclusions."

YOUNG'S WORK ON SOLID AND FLUID TRANSIENTS

The theory of impact

In the years 1801-1803 Young interrupted his medical career at the newly founded Royal Institution in London, where he held the chair of Natural Philosophy in 1802-1803. For his lectures he prepared in very short time a syllabus (20) consisting of an amazing five hundred articles on the subjects: 1. Mechanics, 2. Hydrodynamics, 3. Physics, and 4. Mathematical demonstration. These "Lectures" were published in 1807 (15), and reprinted in 1845 (16) and 2002 (17). It is remarkable that Young never received the promised remuneration of 1000 pounds owing to the bankruptcy of the publishers.

Young (15, pp. 143-145) found that the strain ε produced by the impact of elastic solid bodies equals v/c. With Hooke's law stating that $\varepsilon = -\sigma/E$, where σ is stress and E is Young's modulus of elasticity, this gives $\sigma = -Ev/c$. Assuming that $c = \sqrt{(E/\rho)}$, one obtains for the solids equivalent of the Joukowsky Eq 3:

$$\sigma = -\rho c v \tag{4}$$

where (in contrast to pressure) the stress is defined as negative when the material is compressed. Young (1) was the first to find the pressure wave speed for incompressible liquids contained in elastic tubes, and the authors think, and Beal (21, p. 31) states, that Young was also aware of the speed of sound in solid bars,

$$c = \sqrt{\frac{E}{\rho}} \tag{5}$$

As ever, Young's work is difficult to read, but Timoshenko (22, pp. 93-94) gives a neat summary of the above expressed in modern terminology. It is noted that the strain ε in liquids contained in pipes equals p/K^* , where K^* is the effective bulk modulus representing

fluid compressibility and pipe wall elasticity. According to Saint Venant (23), Babinet independently arrived at Eq 5 in 1829 (the year of Young's death).

It is typical for Young (24, 25, 15, 1) that he had found all the ingredients to arrive at the "Joukowsky equation" for solids *and* fluids, but that his achievements were not picked up by his contemporaries. For example, Young (26, p. 23) mentions that "the magnitude of the pulse ... is proportional to the velocity of the transmission ... ". Young also showed that his *E* modulus applied both to compression and to extension of rods, and also extended its application to liquids (21, p. 31).

The waterhammer wave speed

In 1808 Young delivered the prestigious and still existing Croonian Lecture of the Royal Society. In preparation for this lecture he wrote Ref (1), which is the key paper for his work on the propagation of pressure waves in tubes. It included a new formula for the steady flow of fluids in pipes, the resistance to flow caused by bends, and the propagation of a disturbance through an elastic tube. The Croonian Lecture itself was on the functions of hearts and arteries (26). The prevailing view of the time was that contraction of the walls of arteries was an important cause of the circulation of blood in the human body, but Young's paper conclusively disproved this idea. Young's paper (1) is of fundamental importance to the history of waterhammer, because he derived for the first time the now standard Eq 1 for wave velocity for an incompressible fluid in an elastic tube.

Young's argument proceeded as follows. "The same reasoning, that is employed for determining the velocity of an impulse, transmitted through an elastic *solid* or *fluid* body, is also applicable to the case of an incompressible fluid contained in an elastic pipe" (this clearly suggests that Young had obtained the speed of sound in a solid bar). The problem is then to determine the apparent modulus of elasticity conferred on the incompressible fluid by the elasticity of pipe walls, or, in Young's terminology, to discover "the height of the modulus" to be substituted into Newton's basic formula (24, 25)

$$c = \sqrt{gh} \tag{6}$$

for the speed of sound, this formula giving a velocity half as great as that of a body falling freely from a height $2h \ [2h = g \ t^2/2 \ \text{gives} \ t = \sqrt{(4h/g)}$, and therefore $gt = 2\sqrt{(gh)} \]$. Note that Young first introduced his modulus with the dimension of height rather than the modern dimension of stress (22, p. 92; 27, p. 82; 28, p. 155) which is due to Navier (29), a custom that is continued by contemporary hydraulicians who use head to denote pressure in liquids. Note that $h = p/(\rho g)$ in Eq (6) gives the sonic speed in gas, $\sqrt{(p/\rho)}$.

Continuing the argument, if the pipe is such that the increase in tension force varies as the increase in circumference or diameter from the natural state (i.e., the pipe is elastic and obeys Hooke's law) up to the limit (at which the pressure in the fluid must balance the tension in the pipe by Newton's first law) where an infinite increase in diameter occurs (i.e., plastic deformation at elastic limit), then the height of a column of liquid equivalent to the pressure causing failure is designated "the modular column of the pipe". This is an application of the maximum stress theory that was favoured by English writers over the maximum strain theory, which was favoured on the Continent (22, p. 89). The relationship is readily demonstrated since, from the stress/strain curve up to

the elastic limit $f = \sigma \varepsilon/2 = \sigma^2/(2E)$ (for $\sigma = E\varepsilon$) or, replacing the stresses with their equivalent "heights", $2h = (2h)^2 \rho g/(2E)$, i.e., $h = E/(\rho g)$.

For the equivalent elasticity conferred on the incompressible fluid Young used the continuity principle. If a short length of pipe of diameter D and length x is compressed in length by a pressure pulsation to $(x-\delta x)$, then if the fluid is incompressible the diameter D must increase to preserve continuity so that $(2\delta D/D - \delta x/x) = 0$. But the increase in hoop strain $(\delta D/D) = (\sigma/E)$ for a pipe in tension, and the hoop stress for an increase in pressure δp is given by $D\delta p/(2e)$, thus

$$\sigma = \frac{D}{2e}\delta p \tag{7}$$

so that $D/(Ee) = \delta p/(\delta x/x)$. Eq 7 is probably the oldest formula for fluid-structure interaction, and analogous to Young's equation for surface tension (30). The right-hand side of this last relationship defines precisely an apparent compressibility for the liquid, which is therefore given conveniently by the expression on the left-hand side. Young terminated his argument at this point, but it is a trivial matter to make the substitution into Eq. 6 to give the classic result of Eq 1 above explicitly.

Young was undoubtedly in a position to obtain the celerity of the waterhammer wave given by Eq 2, if he so desired. The continuity method he used can be extended to take account of compressible fluids (indeed it was the method used by Korteweg (9), Kries (5) and Joukowsky (2), seventy to ninety years later). Nevertheless he did not, though he did go on to consider the reflection and collision of waves, to state that the particle velocity must be less than the wave velocity and to examine the effect of a contraction in a pipe. As indicated in the previous section, he was also in the position to formulate Joukowsky's Eq 3.

The hydraulic ram

Young was acquainted with the hydraulic ram, a pumping device based on the waterhammer principle. In his "Lectures" (15, Vol. 1, pp. 337-338) he writes:

"The momentum of a stream of water, flowing through a long pipe, has also been employed for raising a small quantity of water to a considerable height.

The passage of the pipe being stopped by a valve, which is raised by the stream, as soon as its motion becomes sufficiently rapid, the whole column of fluid must necessarily concentrate its action almost instantaneously on the valve; and in this manner it loses, as we have before observed, the characteristic property of hydraulic pressure, and *acts as if it were a single solid*; so that, supposing the pipe to be perfectly elastic and inextensible, the impulse must overcome any pressure, however great, that might be opposed to it, and if the valve open into a pipe leading to an air vessel, a certain quantity of the water will be forced in, so as to condense the air, more or less rapidly, to the degree that may be required, for raising a portion of the water in it, to any given height. Mr. Whitehurst (31) appears to have been the first that employed this method: it was afterwards improved by Mr. Boulton (32); and the same machine has lately attracted much attention in France under the denomination of the hydraulic ram of Mr. Montgolfier (33). (Fig. 3.)" (references added by the present authors.) This is Joseph Michel Montgolfier (1740-1810), one of the brothers who built the first manned balloon (in 1783).




Figure 3 The hydraulic ram of Montgolfier (15, Vol. 2, Fig. 323). When the water in the pipe AB has acquired a sufficient velocity, it raises the valve B, which stops its passage, so that a part of it is forced through the valve C, into the air vessel D, whence it rises through the pipe E.

YOUNG'S WORK ON PIPE FRICTION

Prior to addressing transient flow, Young studied steady flow in pipes (34-36; 15, p. 166). Also, the first part of paper (1) concerns steady pressure losses in pipes. "From own and others' experimental data" Young concluded that "the friction could not be represented by any single power of the velocity, although it frequently approached to the proportion of that power, of which the exponent is 1.8; but that it appeared to consist of two parts, the one varying simply as the velocity, the other as its square. The proportion of these parts to each other must however be considered as different, in pipes of different diameters, the first being less perceptible in very large pipes, or in rivers, but becoming greater than the second in very minute tubes, while the second also becomes greater, for each given portion of the internal surface of the pipe, as the diameter is diminished." With hindsight, Young found here the laminar (linear), fully turbulent (square), and intermediate turbulent (Blasius 1.75) flow regimes. Laird (37) writes on this: "In the 1808 paper (1) Young gives an analysis of the (steady) pressureflow relations in tubes and was well ahead of his time in describing scaling laws of such a flow. The relative importance of the square law vs the linear "Poiseuille like" term are discussed as a function of dimensions, velocity, viscosity, etc. In fact, the essence of scaling with Reynolds' number is clearly enunciated roughly forty years before Osborne Reynolds (38) carried out his crucial experiments." A historical account of the subject is given in Refs (39) and (40).

AFTER YOUNG

In 1850, Ernst-Heinrich Weber published a paper (10) on experiments with blood flow in which he stated that his brother Wilhelm Weber had prepared a theory for the wave celerity which was found to be the same as the till then forgotten result for Eq 1 of Thomas Young. Wilhelm finally published this (11) in 1866. Going further than Young, he combined the two first-order linear relations for the elasticity of the pipe walls and the acceleration of the fluid column to give a wave equation including the wave celerity in the form

$$c = \sqrt{\frac{R}{2k\rho}} \tag{8}$$

where his elastic modulus was defined as k = dR/dp, which in modern notation is $k = R^2/(Ee)$ for circular pipes with R = D/2, hence giving Eq 1.

Subsequent to the brothers Weber, there were a number of studies in this field, including a comprehensive series of experiments in flexible tubes by Marey (41, 42). Marey, though, lacked the necessary mathematics to develop a theory, so this was done for him by Resal, editor of the Journal de Mathématiques Pures et Appliquées. Resal (43, 44) rederived independently the result of Young and Wilhelm Weber and seems to have been the first to write it explicitly in its familiar modern form of Eq 1. Contemporaneously Moens (45) had modified the Weber Eq 8 with a factor whose mean value was close to 1 (4) and finally in 1878 Korteweg (9) derived the complete result including fluid elasticity (Eq 2).

CONCLUSION

On the basis of the following statements:

- about waterhammer (pulse) pressure rise (26, p. 23): the magnitude of the pulse ... is proportional to the velocity of the transmission ...
- about liquid flow suddenly stopped by valve closure (15, p. 338): ... and acts as if it were a single solid ...
- about impact of solids (15, pp. 143-145): the strain produced by the impact of elastic bodies equals the ratio of the convective velocity to the acoustic speed
- about the acoustic speed in solids (21, p. 31): he calculated the velocity of the compression wave that travels through a material following an impact
- about the analogy between solids and liquids (21, p. 31):

 Young showed that his modulus applied both to compression and to extension of rods and also extended its application to liquids

and in addition to his well known pressure wave speed Eq 1, Young arguably arrived at the concepts embodied in the Joukowsky Eq 3, which is the fundamental equation for waterhammer.

ACKNOWLEDGEMENT

Much of the material presented is based on the papers (18), (20) and (21) and on the biographies listed in the Appendix. Figures 1 and 2 are reproduced with kind permission of the Royal Society (London, UK).

REFERENCES

- (1) Young, T. (1808). "Hydraulic investigations, subservient to an intended Croonian lecture on the motion of the blood." *Philosophical Transactions of the Royal Society of London* 98, 164-186 [Errata in Ref (26), p. 31]. Also in (1809) *Nicholson's Journal* 22, 104-124.
- (2) Anderson, A. (2000). "Celebrations and challenges waterhammer at the start of the 20th and 21st centuries." *Proceedings of the 8th International Conference on Pressure*

- *Surges, The Hague, The Netherlands*, 317-322, A. Anderson, ed., BHR Group, Cranfield, UK; Professional Engineering Publishing, Bury St Edmunds, UK.
- (3) Boulanger, A. (1913). "Étude sur la propagation des ondes liquides dans les tuyaux élastiques." ("Study on the propagation of liquid waves in elastic tubes.") *Travaux et Mémoires de l'Université de Lille, Nouvelle Serie, II. Médecine-Sciences* 8, Tallandier, Lille, France; Gauthier-Villars, Paris, France (in French).
- (4) Lambossy, P. (1950; 1951). "Aperçu historique et critique sur le problème de la propagation des ondes dans un liquide compressible enfermé dans un tube élastique." ("Historical outline and review on the problem of wave propagation in a compressible liquid enclosed in an elastic tube.") *Helvetica Physiologica et Pharmacologica Acta* 8(2), 209-227; 9(2), 145-161 (in French).
- (5) Tijsseling, A.S., and Anderson, A. (2004). "A precursor in waterhammer analysis rediscovering Johannes von Kries." *Proceedings of the 9th International Conference on Pressure Surges, Chester, UK*, 739-751, S.J. Murray, ed., BHR Group, Cranfield, UK.
- (6) Galabin, A.L. (1876). "On the causation of the water-hammer pulse, and its transformation in different arteries as illustrated by the graphic method." *Medico-Chirurgical Transactions*, published by the Royal Medical and Chirurgical Society of London, 41(2), 361-388.
- (7) Lindsay, R.B. (1966). "The story of acoustics." *Journal of the Acoustical Society of America* 39(4), 629-644. [Reprinted in 1973: R.B. Lindsay, ed., *Acoustics: Historical and Philosophical Development*, Benchmark Papers in Acoustics, Dowden Hutchinson and Ross, Stroudsberg, Pa, USA, 5-20.]
- (8) Anderson, A (1976), "Menabrea's note on waterhammer: 1858." ASCE *Journal of the Hydraulics Division*, 102(1), 29-39.
- (9) Korteweg, D.J. (1878). "Ueber die Fortpflanzungsgeschwindigkeit des Schalles in elastischen Röhren." ("On the velocity of propagation of sound in elastic tubes.") *Annalen der Physik und Chemie, New Series* 5(12), 525-542 (in German).
- (10) Weber, E-H. (1850). "Ueber die Anwendung der Wellenlehre vom Kreislaufe des Blutes und insbesondere auf die Pulslehre." ("On the application of wave theory to the circulation of blood and in particular to the pulse.") Berichte über die Verhandlungen der Königlichen Sächsischen Gesellschaft der Wissenschaften zu Leipzig, Leipzig, Germany, Mathematical-Physical Section, 2, 164-204 (in German).
- (11) Weber, W. (1866). "Theorie der durch Wasser oder andere incompressible Flüssigkeiten in elastischen Röhren fortgepflanzten Wellen." ("Theory of waves propagating in water or in other incompressible liquids contained in elastic tubes.") Berichte über die Verhandlungen der Königlichen Sächsischen Gesellschaft der Wissenschaften zu Leipzig, Leipzig, Germany, Mathematical-Physical Section, 18, 353-357 (in German).
- (12) Giaquinta, A.R. (1971). "The historical development of the engineering analysis of blood flow." *La Houille Blanche* 26(4), 327-341.
- (13) Goupil, M. (1907). "Notice sur les principaux travaux concernant le coup de bélier et spécialement sur le mémoire et les expériences du Professeur N. Joukovsky (1898)." ["Report on the most important achievements in waterhammer and in particular on the work and experiments of Professor N. Joukovsky (1898)."] *Annales des Ponts et Chaussées* 77(1), 199-221 (in French).
- (14) Wood, F.M. (1970). History of water-hammer. C.E. Research Report No. 65, Department of Civil Engineering, Queen's University at Kingston, Ontario, Canada.

- (15) Young, T. (1807). "A course of lectures on natural philosophy and the mechanical arts." Vol. I: Text, Vol. II: Plates, Joseph Johnson, London, UK. [See also Young, T. (1802), "A syllabus for a course of lectures on natural and experimental philosophy." Four volumes, The Royal Institution, London, UK. Original notebooks available: http://www.ucl.ac.uk/Library/special-coll/tyoung.shtml commented by G.N. Cantor in Ref (20)].
- (16) Young, T. (1845). "A course of lectures on natural philosophy and the mechanical arts." A new edition, with references and notes by Philip Kelland, Taylor and Walton, London, UK. Available from: http://books.google.com/; http://www.archive.org/details/courseoflectures01younrich; http://www.archive.org/details/courseoflectures02younrich.
- (17) Young, T. (2002). "A course of lectures on natural philosophy and the mechanical arts." Four volumes, with a new introduction by Nicholas J. Wade (University of Dundee), Thoemmes, Bristol, UK.
- (18) Hilts, V.L. (1978). "Thomas Young's "Autobiographical sketch"." *Proceedings of the American Philosophical Society* 122(4), 248-260. French translation by A. Chappert in 1981: "L'Autobiographical sketch de Thomas Young Traduction française." *Revue d'Histoire des Sciences* 34(2), 137-147 (in French). Original papers available as Young, T. (1826 or 1827). "An article intended for a future edition of the Encyclopædia Britannica." Galton Papers 120/1, UCL Library, University College London, UK.
- (19) Helmholtz, H. (1873). Popular lectures on scientific subjects. Longmans, Green and Co, London, UK; Appleton, New York, USA. Second edition (1881), p. 220. Facsimile reprint in 1999, Thoemmes Continuum, London, UK. Available from: http://www.archive.org/details/lecturesonscient00helmiala.
- (20) Cantor, G.N. (1970). "Thomas Young's lectures at the Royal Institution." *Notes and Records of the Royal Society of London* 25(1), 87-112.
- (21) Beal, A.N. (2000). "Who invented Young's modulus?" *The Structural Engineer* 78(14), 27-32.
- (22) Timoshenko, S.P. (1953). "History of strength of materials." McGraw-Hill, New York. (Reprint in 1983, Dover Publications, New York, USA.)
- (23) Saint-Venant, A.J.C. Barré de (1883). In: "Théorie de l'élasticité des corps solides de Clebsch, A. Clebsch, translated and annotated by A.J.C. Barré de Saint-Venant", Dunod, Paris, France, Note finale du 60 (in French). (Facsimile reprint in 1966, Johnson Reprint, New York, USA.)
- (24) Young, T. (1800). "Outlines of experiments and enquiries respecting sound and light." *Philosophical Transactions of the Royal Society of London* 90, 106-150.
- (25) Young, T. (1802). "On the velocity of sound." *Journal of the Royal Institution of Great Britain* 1, 214-216.
- (26) Young, T. (1809). "The Croonian lecture. On the functions of the heart and arteries." *Philosophical Transactions of the Royal Society of London* 99, 1-31.
- (27) Todhunter, I., and Pearson, K. (Volume I, 1886; Volume II, Parts I and II, 1893). "A history of elasticity and strength of materials." Cambridge University Press, Cambridge, UK. (Reprinted in 1960 as: "A history of the theory of elasticity and of the strength of materials from Galilei to Lord Kelvin.", Dover Publications, New York, USA.)
- (28) Straub, H. (1952). "A history of civil engineering (an outline from ancient to modern times)." Translator E. Rockwell; Leonard Hill, London, UK; MIT Press, Cambridge, USA.

- (29) Navier, C.L.M.H. (1827). "Mémoire sur les lois de l'équilibre et du mouvement des corps solides élastiques." ("Memoir on the laws of equilibrium and motion of elastic solid bodies.") *Mémoires de l'Académie des Sciences* 7, 375-393 (in French).
- (30) Young, T. (1805). "An essay on the cohesion of fluids." *Philosophical Transactions of the Royal Society of London* 95, 65-87.
- (31) Whitehurst, J. (1775). "Account of a machine for raising water, executed at Oulton, in Cheshire, in 1772. In a letter from Mr. John Whitehurst to Dr. Franklin." *Philosophical Transactions* 65, 277-279.
- (32) Boulton, M. (1798). Repertory of Arts 9 (Patent).
- (33) Montgolfier, J.M. de (1803). "Note sur le bélier hydraulique, et sur la manière d'en calculer les effets." ("Note on the hydraulic ram, and on the method of calculating the power.") *Journal des Mines* 13(73), 42-51 (in French).
- (34) Young, T. (1802). "An account of an experiment on the velocity of water through a vertical pipe." *Journal of the Royal Institution of Great Britain* 1, 231-233. Also in (1803) *Nicholson's Journal* 6, 59-61.
- (35) Young, T. (1803). "Remarks on resistance of fluids." *Journal of the Royal Institution of Great Britain* 2, 14-16.
- (36) Young, T. (1803). "Further considerations on the resistance of fluids." *Journal of the Royal Institution of Great Britain* 2, 78-80.
- (37) Laird, J.D. (1980). "Thomas Young, M.D. (1773-1829)." *American Heart Journal* 100(1), 1-8.
- (38) Jackson, J.D., and Launder, B.E. (2007). "Osborne Reynolds and the publication of his papers on turbulent flow." *Annual Review of Fluid Mechanics* 39, 19-35.
- (39) Sutera, S.P., and Skalak, R. (1993). "The history of Poiseuille's law." *Annual Review of Fluid Mechanics* 25, 1-19.
- (40) Brown, G.O. (2002). "The history of the Darcy-Weisbach equation for pipe flow resistance." *Proceedings of the 150th Anniversary Conference of ASCE*, J. Rogers, and A. Fredrich, eds., ASCE, Reston, Va., USA, 34–43.
- (41) Marey, E.J. (1858). "Recherches sur la circulation du sang (études hydrauliques)". ("Investigations on the circulation of blood (hydraulic studies)".) *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* 46, 483-485 (in French).
- (42) Marey, E.J. (1875-1880). "Physiologie expérimentale." ("Experimental physiology.") Ecole Pratique des Hautes Etudes. Travaux du Laboratoire de M. Marey, 4 Volumes, G. Masson, Libraire de l'Académie de Médecine, Paris, France (in French).
- (43) Resal, H. (1876), "Note sur les petits mouvements d'un fluide incompressible dans un tuyau élastique". ("Note on the small movements of an incompressible fluid in an elastic tube.") *Journal de Mathématiques Pures et Appliquées*, 3rd Series, 2, 342-344 (in French).
- (44) Resal, H. (1876), "Sur les petits mouvements d'un fluide incompressible dans un tuyau élastique." ("On the small movements of an incompressible fluid in an elastic tube.") *Comptes Rendues Hebdomadaires de l'Académie des Sciences* 82, 698-699 (in French).
- (45) Moens, A.I. (1878). "Die Pulscurve." ("The pulse curve.") E.J. Brill, Leyden, The Netherlands (in German).

APPENDIX A: BIBLIOGRAPHY ON YOUNG'S LIFE AND ACHIEVEMENTS

- (46) Robinson, A. (2007). "Thomas Young and the Rosetta Stone." *Endeavour, New Series* 31(2), 59-64.
- (47) Matthews, R. (2006). "The curious life of a polymath." *New Scientist* 192(2573), 59. Review of Robinson (2006a).
- (48) Schwab, I.R. (2006). "AJO history of ophthalmology series. Thomas Young (1773-1829)." *American Journal of Ophthalmology* 142(3), 487.
- (49) Robinson, A. (2006a). "The last man who knew everything Thomas Young, the anonymous polymath who proved Newton wrong, explained how we see, cured the sick, and deciphered the Rosetta stone, among other feats of genius." Pi Press, New York, USA; Oneworld Publications, Oxford, UK.
- (50) Robinson, A. (2006b). "Thomas Young: physicist, physician and polymath." *Physics World* 19(3), 30-33.
- (51) Robinson, A. (2005). "A polymath's dilemma." Nature 438(7066), 291.
- (52) Hondros, E.D. (2005). "Dr. Thomas Young Natural philosopher." *Journal of Materials Science* 40(9-10), 2119-2123.
- (53) Landsman, K. (2003). "Wie was Thomas Young?" ("Who was Thomas Young?") *Nederlands Tijdschrift voor Natuurkunde* 69(2), 24-28 (in Dutch).
- (54) Martindale, C. (2001). "Oscillations and analogies: Thomas Young, MD, FRS, Genius." *American Psychologist* 56(4), 342-345.
- (55) Gauger, G.E. (1997). "The great mind of Thomas Young (1773-1829)." *Documenta Ophthalmologica* 94(1-2), 113-121.
- (56) Bruce Fye, W. (1997). "Thomas Young." *Clinical Cardiology* 20(1), 87-88.
- (57) Griffin, J.P. (1995). "Dr Thomas Young polymath." *Adverse Drug Reactions and Toxicological Reviews* 14(2), 77-81.
- (58) Kline, D.L. (1993). "Thomas Young, forgotten genius: an annotated narrative biography." Vidan Press, Cincinnati, USA.
- (59) Hopkins, R.W. (1991). "Presidential address: Energy, poise, and resilience Daniel Bernoulli, Thomas Young, J. L. M. Poiseuille, and F. A. Simeone." *Journal of Vascular Surgery* 13(6), 777-784.
- (60) Behrman, S. (1975). "Thomas Young, the physician." Clio Medica 10, 277-284.
- (61) Oldham, F. (1974). "Thomas Young." British Medical Journal 4(5937), 150-152.
- (62) Koelbing, H.M., (1974). "Thomas Young (1773-1829), die physiologische Optik und die Ägyptologie." ("Thomas Young (1773-1829), physiological optics and Egyptology.") *Gesnerus, Swiss Journal of the History of Medicine and Sciences* 31(1), 56-75 (in German).
- (63) Hodgkin, A. (1974). "Address of the president Sir Alan Hodgkin, O.M. at the Anniversary Meeting, 30 November 1973." *Proceedings of the Royal Society of London, Series B, Biological Sciences* 185(1078), v-xx.
- (64) Fonda, G. (1973). "Bicentenary of the birth of Thomas Young, M.D., F.R.S." *British Journal of Ophthalmology* 57(11), 803-808.
- (65) Hill, B. (1973). "Phenomenon Young'. Thomas Young, M.D., F.R.S." *Practitioner* 210(260), 831-835.
- (66) Cox, G.A. (1973). "Thomas Young 1773-1829." Physics Education 8(6), 396-399.
- (67) Douglas, A.V. (1973). "Thomas Young, 1773-1829." Journal of the Royal Astronomical Society of Canada 67, 150.

- (68) Herivel, J. (1973). "Thomas Young (1773-1829)." Endeavour 32(115), 15-18.
- (69) Rubinowicz, A. (1957). "Thomas Young and the theory of diffraction." *Nature* 180(4578), 160-162.
- (70) Wood, A. (1954). "Thomas Young Natural philosopher, 1773-1829," completed by Frank Oldham, with a memoir of Alexander Wood by Charles E. Raven, Cambridge University Press, Cambridge, UK.
- (71) Lamor, J. (1934). "Thomas Young." Nature 133, 276-279.
- (72) Oldham, F. (1933). "Thomas Young, F.R.S. Philosopher and physician." Edward Arnold & Co, London, UK.
- (73) Robinson, H.S. (1929). "Thomas Young A chronology and a bibliography with estimates of his work and character." *Medical Life* 36, 527-540.
- (74) Rowell, H.S. (1912). "Thomas Young and Göttingen." Nature 88, 519.
- (75) Peacock, G. (1855). "Life of Thomas Young, M.D., F.R.S., &c." John Murray, London.
- (76) Arago, D.-F.J. (1854). "Thomas Young Gedächtnißrede, gelesen in der öffentlichen Sitzung der Akademie der Wissenschaften am 26. November 1832." In: "Franz Arago's sämtliche Werke." ("François Arago's collected works."), Verlag von Otto Wigand, Leipzig, Germany, Vol. 1, 191-233. Republished in 1929, with epilogues of A. Einstein and M. von Rohr, in: *Die Naturwissenschaften* 17(20), 347-364 (in German).
- (77) Arago, D.-F.J. (1836). "Biographical memoir of Dr Thomas Young." *Edinburgh New Philosophical Journal* 20(40), 213-240.
- (78) Arago, D-F.J. (1835). "Éloge historique du docteur Thomas Young." *Mémoires de l'Académie des Sciences* 13, lvii-cv (in French).
- (79) Gurney, H. (1831). "Memoir of the life of Thomas Young." John & Arthur Arch, London.
- (80) Morley, J. (1994). "The importance of being historical: civil engineers and their history." ASCE *Journal of Professional Issues in Engineering Education and Practice* 120(4), 419-428.

APPENDIX B: YOUNG'S LIFE

Thomas Young has inspired many people to write biographical papers in all sorts and sizes (20-21, 37, 46-79), mostly of hagiographic nature (80). The major biography is Peacock's "Life of Thomas Young" (75), which is based on a large collection of letters and on Hudson Gurney's "Memoir of Thomas Young" (79). Gurney is Young's long-time friend and his "Memoir" is an extension of Young's own biographical sketch, which was published by Hilts in 1978 (18) shortly after its rediscovery around 1976. Other biographies are those of Oldham (72), Wood (70), Kline (58) and Robinson (49).

Young was born as the eldest son in a Quaker family on 13 June 1773 in Milverton (Somerset, UK). At the age of two he could read fluently and before the age of four he had read the bible twice. At the age of fourteen he was fluent in the classic languages and requested to be the "director general" of the Latin and Greek "of the whole party" (18, p. 251). Although he had several teachers and tutors in his early education, he may be regarded largely as self taught. From 1792 to 1803 he studied medicine in London, Edinburgh, Göttingen (Germany) and Cambridge. By coincidence, in Göttingen he lived in the building where in 1833 Wilhelm Weber (with Gauss) invented an electromagnetic telegraph. He was a physician – and not a physicist – his whole life, running a private practice from 1799 to 1814. In 1794, at the age of 21 and officially still a student, he was elected Fellow of the Royal Society, rewarding his paper on vision (read before the society in 1793). He was Professor of Natural Philosophy in the Royal Institution from 1801 to 1803. However, his friends were of opinion that his longer continuance, in the situation of a public teacher, would be unfavourable to his success in medicine, and after having lectured for two winters, he gave up the professorship. In the intervening summer of 1802 he had the pleasure of hearing Napoleon speak at the National Institute in Paris. In the same year, he became Foreign Secretary of the Royal Society, holding this position for the rest of his life. In 1804 he married Eliza Maxwell, which whom he had a happy marriage without children. In 1807 he was an unsuccessful candidate for the post of Physician to Middlesex Hospital, but in 1810, after a very arduous contest, he succeeded to become Physician at St. George's in London and remained this until his death. He was Adviser to the Admiralty on shipbuilding, Secretary of the Board of Longitude, and Superintendent of the vital "Nautical Almanac" from 1818 on, besides physician to and inspector of calculations for the Palladian Insurance Company from 1824 on. He was elected Foreign Member of the French Academy of Science in 1827, succeeding Volta. The French scientists Arago and Gay-Lussac were amongst his friends as well as Humboldt in Germany. Thomas Young died in London on 10 May 1829.