

Ruin excursions, the $G/G/\infty$ queue and tax payments in renewal risk models

Citation for published version (APA):

Albrecher, H., Borst, S. C., Boxma, O. J., & Resing, J. A. C. (2010). *Ruin excursions, the $G/G/\infty$ queue and tax payments in renewal risk models*. (Report Eurandom; Vol. 2010039), (Calculus : symposium on the integration of symbolic computation and mechanized reasoning : proceedings; Vol. 14). Eurandom.

Document status and date:

Published: 01/01/2010

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EURANDOM PREPRINT SERIES
2010-039

**Ruin Excursions, the $G/G/\infty$ queue and
tax payments in renewal risk models**

H. Albrecher, S.C. Borst, O.J. Boxma, J. Resing
ISSN 1389-2355

RUIN EXCURSIONS, THE $G/G/\infty$ QUEUE AND TAX PAYMENTS IN RENEWAL RISK MODELS

H. ALBRECHER,* *University of Lausanne*
S.C. BORST,** *Eindhoven University of Technology*
O.J. BOXMA,*** *Eindhoven University of Technology*
J. RESING,**** *Eindhoven University of Technology*

Abstract

In this paper we investigate the number and maximum severity of the ruin excursion of the insurance portfolio reserve process in the Cramér-Lundberg model with and without tax payments. We also provide a relation of the Cramér-Lundberg risk model with the $G/G/\infty$ queue and use it to derive some explicit ruin probability formulas. Finally, the renewal risk model with tax is considered, and an asymptotic identity is derived that in some sense extends the tax identity of the Cramér-Lundberg risk model.

Keywords: classical risk model; ruin probability; $G/G/\infty$ queue; tax; renewal model

2000 Mathematics Subject Classification: Primary 91B30

Secondary 60K30

1. Introduction

Consider the classical Cramér-Lundberg model in risk theory to describe the surplus process $\{R_t\}$ at time t of an insurance portfolio. Starting with an initial capital x , premium is collected according to a constant premium intensity (normalized to) 1.

* Postal address: Department of Actuarial Science, Faculty of Business and Economics, University of Lausanne, Quartier UNIL-Dorigny, 1015 Lausanne, Switzerland

** Postal address: Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands and Alcatel-Lucent, Bell Labs, 600 Mountain Avenue, P.O. Box 636, Murray Hill, NJ 07974-0636, USA

*** Postal address: Eindhoven University of Technology and EURANDOM, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

**** Postal address: Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

Claims occur according to a homogeneous Poisson process with intensity λ and are paid at the times of their occurrence. The claim sizes are independent and identically distributed random variables with distribution function $H(\cdot)$. Define $\phi_0(x) = \mathbb{P}(R_t \geq 0 \text{ for all } t | R_0 = x)$ as the probability of survival and correspondingly the ruin probability as $\psi_0(x) = 1 - \phi_0(x)$. Let further V_{\max} be the maximum workload in an $M/G/1$ queue with arrival rate λ and service time distribution $H(\cdot)$. Then the following relation between the Cramér-Lundberg risk model and the $M/G/1$ queueing model is classical:

$$\phi_0(x) = e^{-\lambda \int_x^\infty \mathbb{P}(V_{\max} > y) dy}. \quad (1)$$

Let $G(\cdot)$ denote the distribution function of V_{\max} . One way to show (1) is to use the well-known relation

$$G(u) = \mathbb{P}(V_{\max} < u) = 1 - \frac{1}{\lambda} \frac{d}{du} \ln \mathbb{P}(V < u), \quad (2)$$

where V is the stationary workload in the same $M/G/1$ queue as described above, and use the sample path duality result $\phi_0(x) = \mathbb{P}(V < x)$ (see e.g. Asmussen & Albrecher [5] for a recent survey). In [2] another more direct proof of (1) was given and subsequently used to establish a simple proof of the tax identity

$$\phi_\gamma(x) = \left(\phi_0(x) \right)^{\frac{1}{1-\gamma}} = e^{-\frac{\lambda}{1-\gamma} \int_x^\infty \mathbb{P}(V_{\max} > y) dy}, \quad (3)$$

where $\phi_\gamma(x) = 1 - \psi_\gamma(x)$ is the survival probability in a Cramér-Lundberg model with tax rate $0 \leq \gamma \leq 1$, i.e. whenever the risk process is in its running maximum (and hence in a profitable position), a constant proportion γ of the incoming premium is paid as tax ($\gamma = 0$ corresponds to the Cramér-Lundberg model without tax). For extensions of this identity in various directions see [1, 3, 4, 7, 10].

In this paper we will provide a relation of the Cramér-Lundberg risk model with the $G/G/\infty$ queue, which will give rise to another view towards identity (1) and some explicit ruin probability formulas. Subsequently, we will consider the renewal risk model with tax, and establish an asymptotic identity that may be interpreted as an extension of the tax identity (3). We start with some refined results on the number and maximum severity of the ruin excursion in the Cramér-Lundberg model with and without tax.

2. Maximum severity of the ruin excursion

Consider the Cramér-Lundberg model with tax rate γ . Ruin can only occur during an ‘interruption’, i.e., a period in between running maxima. Denote the k th interruption period by P_k . Interruptions occur according to a Poisson process with intensity λ . The probability that no ruin occurs during an interruption that starts at surplus level z is given by $G(z) = 1 - \bar{G}(z)$ (cf. (2)). Let R_{\min} be the lowest surplus value during the ruin excursion. Let further $A_k(x, d)$ be the probability that ruin occurs during the k th interruption P_k and $R_{\min} < -d$, where $d \geq 0$. Then, for $k \in \mathbb{N}$

$$A_k(x, d) = \int_{t=0}^{\infty} \lambda^k \frac{t^{k-1}}{(k-1)!} e^{-\lambda t} \left[\int_{v=0}^t G(x + (1-\gamma)v) \frac{dv}{t} \right]^{k-1} \bar{G}(x + (1-\gamma)t + d) dt. \quad (4)$$

Here we have used that the sum of k independent exponential arrival intervals is Erlang(k, λ) distributed, and given that their sum is t , the interruption epochs are uniformly distributed on $[0, t]$.

Proposition 2.1. *Let $A(x, d)$ be the probability that ruin occurs and the lowest surplus value of the ruin excursion is smaller than $-d \leq 0$. Then*

$$A(x, d) = \int_x^{\infty} \frac{\phi'_\gamma(w+d)}{\phi_\gamma(w+d)} \frac{\phi_\gamma(x)}{\phi_\gamma(w)} dw. \quad (5)$$

Proof. We have

$$\begin{aligned} A(x, d) &= \sum_{k=1}^{\infty} A_k(x, d) = \int_{t=0}^{\infty} \lambda e^{-\lambda t} \bar{G}(x + (1-\gamma)t + d) e^{\lambda \int_{v=0}^t G(x+(1-\gamma)v) dv} dt \\ &= \int_{t=0}^{\infty} \lambda \bar{G}(x + (1-\gamma)t + d) e^{-\lambda \int_{v=0}^t \bar{G}(x+(1-\gamma)v) dv} dt. \end{aligned} \quad (6)$$

Now the result follows from (2) and (3). \square

Remark 1. Clearly $d = 0$ gives $A(x, 0) = 1 - \phi_\gamma(x) = \psi_\gamma(x)$, so that in this case we indeed recover the usual ruin probability.

Remark 2. An alternative way to establish (6) is to use the joint distribution of the maximum surplus before ruin $R_{\max} = \sup_{t \geq 0} R_t I_{\{R_u \geq 0 \text{ for all } u \in [0, t]\}}$ and the

maximum deficit of the ruin excursion R_{\min} . Concretely,

$$\begin{aligned} & \mathbb{P}(R_{\max} \in [y, y + dy]; R_{\min} \leq -d) \\ &= \frac{d}{dy} \left[1 - e^{-\frac{\lambda}{1-\gamma} \int_{v=x}^y \mathbb{P}(V_{\max} > v) dv} \right] \cdot \mathbb{P}(V_{\max} > y + d | V_{\max} > y) \\ &= \frac{\lambda}{1-\gamma} \mathbb{P}(V_{\max} > y + d) \cdot e^{-\frac{\lambda}{1-\gamma} \int_{v=x}^y \mathbb{P}(V_{\max} > v) dv}, \end{aligned}$$

which also yields (6) upon integration over $y \geq x$. Note in addition that the time spent in the running maximum until ruin is given by $(R_{\max} - x)/(1 - \gamma)$.

Proposition 2.2. *The generating function $\Phi(z, x, d) := \sum_{k=1}^{\infty} z^k A_k(x, d)$ is given by*

$$\Phi(z, x, d) = z \int_x^{\infty} \frac{\phi'_\gamma(w + d)}{\phi_\gamma(w + d)} \left(\frac{\phi_\gamma(x)}{\phi_\gamma(w)} \right)^z e^{-\lambda(1-z)(w-x)/(1-\gamma)} dw. \quad (7)$$

Proof. From (4) it follows that

$$\begin{aligned} \Phi(z, x, d) &= z \int_{t=0}^{\infty} \lambda e^{-\lambda t} e^{z\lambda \int_{v=0}^t G(x+(1-\gamma)v) dv} \overline{G}(x + (1-\gamma)t + d) dt \\ &= z \int_{t=0}^{\infty} \lambda \overline{G}(x + (1-\gamma)t + d) e^{-\lambda(1-z)t} e^{-\lambda z \int_{v=0}^t \overline{G}(x+(1-\gamma)v) dv} dt, \end{aligned}$$

so that the assertion again follows from (2) and (3). \square

Denote by K the number of the interruption that leads to ruin (K is a defective random variable on the positive integers). Then starting at (7) with $d = 0$, some elementary calculations lead to the following result:

Corollary 2.1.

$$\begin{aligned} \mathbb{E}[K | \text{Ruin occurs with } R_0 = x] &= \frac{\frac{\partial}{\partial z} \Phi(z, x, 0) \Big|_{z=1}}{\psi_\gamma(x)} \\ &= \ln \phi_\gamma(x) \left(1 - \frac{1}{\psi_\gamma(x)} \right) - \frac{\lambda}{1-\gamma} \left(x - \frac{\phi_\gamma(x)}{\psi_\gamma(x)} \int_x^{\infty} \frac{w \phi'_\gamma(w)}{\phi_\gamma^2(w)} dw \right). \end{aligned}$$

On the other hand, one may rewrite (4) as follows:

$$\begin{aligned} A_k(x, d) &= \int_{t=0}^{\infty} \frac{\lambda}{(k-1)!} e^{-\lambda t} \left[\lambda \int_{v=0}^t G(x + (1-\gamma)v) dv \right]^{k-1} \overline{G}(x + (1-\gamma)t + d) dt \\ &= \int_{t=0}^{\infty} \frac{e^{-\lambda t}}{(k-1)!} \left[\lambda t - \int_{v=0}^t \frac{\phi'_0(x + (1-\gamma)v)}{\phi_0(x + (1-\gamma)v)} \right]^{k-1} \frac{\phi'_0(x + (1-\gamma)t + d)}{\phi_0(x + (1-\gamma)t + d)} dt \\ &= \int_{t=0}^{\infty} \frac{e^{-\lambda t}}{(k-1)!} \left[\lambda t - \ln \frac{\phi_\gamma(x + (1-\gamma)t)}{\phi_\gamma(x)} \right]^{k-1} \frac{\phi'_0(x + (1-\gamma)t + d)}{\phi_0(x + (1-\gamma)t + d)} dt. \quad (8) \end{aligned}$$

Integrating over d and some elementary algebra then gives the following expressions:

Corollary 2.2. *The expected maximum severity of the ruin excursion, with ruin occurring at the k th interruption, is given by*

$$\begin{aligned} & \mathbb{E}[|R_{\min}| \cdot I_{\{\text{ruin at } P_k\}} | R_0 = x] \\ &= -\frac{1}{(1-\gamma)^k} \int_x^\infty \frac{e^{-\lambda(w-x)/(1-\gamma)}}{(k-1)!} \left[\lambda(w-x) - \ln \frac{\phi_0(w)}{\phi_0(x)} \right]^{k-1} \ln \phi_0(w) \, dw. \end{aligned}$$

Furthermore, the expected maximum severity of the ruin excursion given that ruin occurs, is given by

$$\mathbb{E}[|R_{\min}| \mid \text{ruin occurs with } R_0 = x] = -\frac{\phi_\gamma(x)}{\psi_\gamma(x)} \int_x^\infty \frac{\ln \phi_\gamma(w)}{\phi_\gamma(w)} \, dw.$$

Remark 3. From the above formulas, it is straightforward to write down the probability that the ruin excursion stays above surplus level $-d < 0$, given that ruin occurs, as

$$\frac{A(x, 0) - A(x, d)}{\psi_\gamma(x)} = \frac{1}{\psi_\gamma(x)} \int_x^\infty \left[\frac{\phi'_\gamma(w)}{\phi_\gamma(w)} - \frac{\phi'_\gamma(w+d)}{\phi_\gamma(w+d)} \right] \frac{\phi_\gamma(x)}{\phi_\gamma(w)} \, dw.$$

For the case without tax ($\gamma = 0$), this formula can be compared with the following related classical formula for the maximum severity M of ruin, which is defined as the smallest value of the risk process after ruin before level 0 (instead of the running maximum) is reached again:

$$P(M \leq d | R_0 = x \text{ and ruin occurs}) = \frac{\phi_0(x+d) - \phi_0(x)}{\phi_0(d)(1 - \phi_0(x))}$$

(see Picard [8]).

3. Relation with the $G/G/\infty$ queue

Consider the following situation. We have a sequence of pairs of random variables $(X_1, Y_1), (X_2, Y_2), (X_3, Y_3), \dots$, for which we want to calculate

$$\phi(x) = P \left(Y_i \leq x + \sum_{j=1}^i X_j \text{ for all } i = 1, 2, \dots \right). \quad (9)$$

As a first interpretation, the function $\phi(x)$ is the survival probability in the risk model, if the X_i 's represent the increase of the surplus during periods in which the surplus process is in its running maximum (in the absence of tax payments, the X_i 's equivalently represent the lengths of the periods during which the surplus process is

in its running maximum) and the Y_i 's represent the maximal decreases of the surplus process in periods during which the surplus process is not in a profitable situation (i.e., the Y_i 's correspond to identically distributed copies of the random variable V_{\max}).

A second interpretation of the function $\phi(x)$ is as the steady-state probability that at an arrival instant in a $G/G/\infty$ queue the residual service times of all the customers present in the system are less than x . Here, the X_i 's represent the interarrival times of the customers and the Y_i 's represent the service times of the customers.

For the moment we assume that for different i and j the pairs of random variables (X_i, Y_i) and (X_j, Y_j) are independent and identically distributed. Furthermore, we assume that, within a pair, the random variables X_i and Y_i are independent.

Remark 4. These assumptions are satisfied in the Cramér-Lundberg risk model, where the claim arrival process is a Poisson process. However, when the claim arrival process is a general renewal process the random variables Y_i and X_{i+1} are dependent. In the related $G/G/\infty$ queueing model this will mean that the service time of a customer depends on the previous interarrival time.

Let us denote by $F(\cdot)$ the common distribution function of the random variables X_i (with corresponding probability density function $f(\cdot)$). Furthermore, we denote by $G(\cdot)$ the common distribution function of the random variables Y_i .

Conditioning on the value of X_1 we obtain

$$\phi(x) = \int_{x_1=0}^{\infty} \phi(x+x_1)G(x+x_1)f(x_1)dx_1. \quad (10)$$

Iteration of this equation yields

$$\begin{aligned} \phi(x) &= \int_{x_1=0}^{\infty} \int_{x_2=0}^{\infty} \phi(x+x_1+x_2)G(x+x_1+x_2)G(x+x_1)f(x_2)f(x_1)dx_2dx_1 \\ &\vdots \\ &= \lim_{M \rightarrow \infty} \int_{x_1=0}^{\infty} \dots \int_{x_M=0}^{\infty} \phi(x + \sum_{j=1}^M x_j) \prod_{i=1}^M \left\{ G(x + \sum_{j=1}^i x_j) f(x_i) \right\} dx_M \dots dx_1. \end{aligned}$$

Example 3.1. (X_i 's are deterministic.) If the X_i 's are deterministic, say $X_i = w$, we have

$$\phi(x) = \phi(x+w)G(x+w) = \prod_{i=1}^{\infty} G(x+w \cdot i).$$

Example 3.2. (Y_i 's are deterministic.) If the Y_i 's are deterministic, say $Y_i = v$, we have

$$\phi(x) = \begin{cases} 1 & \text{for } x \geq v, \\ 1 - F(v - x) & \text{for } x < v. \end{cases}$$

Example 3.3. (X_i 's are exponential with parameter λ .) This is the case of the Cramér-Lundberg risk model. For an $M/G/\infty$ queue it is well-known (see e.g. [9]) that the steady-state distribution of the number of customers is Poisson distributed and that the residual service times of the customers are all i.i.d. according to the excess lifetime distribution

$$G_e(x) := \frac{1}{\mathbb{E}[Y]} \int_0^x \bar{G}(y) dy.$$

Hence we find

$$\phi(x) = \sum_{n=0}^{\infty} \frac{(\lambda \mathbb{E}[Y])^n}{n!} e^{-\lambda \mathbb{E}[Y]} [G_e(x)]^n = e^{-\lambda \mathbb{E}[Y](1-G_e(x))} = e^{-\lambda \int_x^{\infty} \bar{G}(y) dy}, \quad (11)$$

which can be interpreted as yet another approach to establish formula (1). Of course, formula (11) can also be obtained from equation (10) which in this case takes the form

$$\phi(x) = \lambda \int_0^{\infty} \phi(x + x_1) G(x + x_1) e^{-\lambda x_1} dx_1.$$

Introducing $T(x) := e^{-\lambda x} \phi(x)$ yields

$$T(x) = \lambda \int_x^{\infty} T(u) G(u) du,$$

which gives $T'(x) = -\lambda G(x) T(x)$. It follows that $T(x) = C e^{-\lambda \int_0^x G(y) dy}$, so that $\phi(x) = C e^{\lambda \int_0^x \bar{G}(y) dy}$ with C some constant yet to be determined. Letting $x \rightarrow \infty$, we find $C = e^{-\lambda \int_0^{\infty} \bar{G}(y) dy}$, and hence $\phi(x) = e^{-\lambda \int_x^{\infty} \bar{G}(y) dy}$.

Example 3.4. (Y_i 's are exponential with parameter ν .) For a $G/M/\infty$ queue it is well-known (see e.g. [9]) that the steady-state probability that an arriving customer finds n customers in the system is given by

$$p_n = \sum_{r=n}^{\infty} (-1)^{r-n} \binom{r}{n} B_r,$$

where B_r is given by

$$B_r = \prod_{i=1}^r \left(\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right)$$

and $\tilde{F}(s)$ is the LST of the interarrival time distribution. Exploiting the lack-of-memory property of the exponential distribution, we hence have

$$\begin{aligned}
\phi(x) &= \sum_{n=0}^{\infty} p_n (1 - e^{-\nu x})^n \\
&= \sum_{n=0}^{\infty} \sum_{r=n}^{\infty} (-1)^{r-n} \binom{r}{n} \prod_{i=1}^r \left(\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right) (1 - e^{-\nu x})^n \\
&= \sum_{r=0}^{\infty} \prod_{i=1}^r \left(\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right) \sum_{n=0}^r \binom{r}{n} (-1)^{r-n} (1 - e^{-\nu x})^n \\
&= \sum_{r=0}^{\infty} \left(\prod_{i=1}^r \left(\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right) \right) (-e^{-\nu x})^r.
\end{aligned}$$

In the special case that the interarrival times are exponential as well (with parameter λ), we have

$$\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} = \frac{\lambda}{i\nu}$$

and correspondingly

$$\phi(x) = \sum_{r=0}^{\infty} \prod_{i=1}^r \left(\frac{\lambda}{i\nu} \right) (-e^{-\nu x})^r = \sum_{r=0}^{\infty} \left(-\frac{\lambda}{\nu} e^{-\nu x} \right)^r / r! = e^{-\frac{\lambda}{\nu} e^{-\nu x}} = e^{-\lambda \int_x^{\infty} e^{-\nu y} dy} \quad (12)$$

as before.

If on the other hand the interarrival times are Erlang(2, λ) distributed, we have

$$\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} = \frac{\lambda^2}{(i\nu)^2 + 2\lambda i\nu}$$

and consequently

$$\phi(x) = \sum_{r=0}^{\infty} \prod_{i=1}^r \left(\frac{\lambda^2}{(i\nu)^2 + 2\lambda i\nu} \right) (-e^{-\nu x})^r = \sum_{r=0}^{\infty} \left(\frac{\lambda}{\nu} \right)^{2r} \frac{1}{r!} \prod_{i=1}^r \left(\frac{1}{i + 2\frac{\lambda}{\nu}} \right) (-e^{-\nu x})^r.$$

Introducing $\alpha = 2\lambda/\nu$ and using

$$\prod_{i=1}^r \left(\frac{1}{i + \alpha} \right) = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha + r + 1)}$$

gives

$$\phi(x) = \Gamma(\alpha + 1) \sum_{r=0}^{\infty} \frac{\left[-\left(\frac{\lambda}{\nu}\right)^2 e^{-\nu x} \right]^r}{r! \Gamma(\alpha + r + 1)} = \frac{\Gamma(\alpha + 1)}{\left(\frac{\lambda}{\nu} e^{-\nu x/2}\right)^\alpha} \cdot J_\alpha \left(\alpha e^{-\nu x/2} \right) \quad (13)$$

where $J_\alpha(\cdot)$ is the Bessel function of the first kind, defined by

$$J_\alpha(x) = \sum_{r=0}^{\infty} \frac{(-1)^r}{r! \Gamma(r + \alpha + 1)} \left(\frac{x}{2}\right)^{2r + \alpha}.$$

Formula (13) can also be obtained via equation (10): Plugging $f(x) = \lambda^2 x e^{-\lambda x}$ and $G(x) = 1 - e^{-\nu x}$ into (10), differentiating twice yields

$$(e^{-\lambda x} \phi(x))'' = \lambda^2 \phi(x) (1 - e^{-\nu x})$$

or equivalently

$$\phi''(x) - 2\lambda\phi'(x) + \lambda^2 e^{-\nu x} \phi(x) = 0.$$

This ordinary differential equation has the solution

$$\phi(x) = \left(\nu e^{\nu x/2} / \lambda\right)^\alpha \left[C_1 \Gamma(1 + \alpha) J_\alpha(\alpha e^{-x\nu/2}) + C_2 \Gamma(1 - \alpha) J_{-\alpha}(\alpha e^{-x\nu/2}) \right],$$

where C_1, C_2 are constants and again $\alpha = 2\frac{\lambda}{\nu}$. The boundary condition $\lim_{x \rightarrow \infty} \phi(x) = 1$ then gives $C_2 = 0$ and $C_1 = 1$, hence (13).

It is interesting to examine the asymptotic behavior of $\phi(x_\lambda)$, with $x_\lambda := \kappa + \frac{1}{\nu} \log \lambda$, as $\lambda \rightarrow \infty$. It is easily verified that

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \phi(x_\lambda) &= \lim_{\lambda \rightarrow \infty} \sum_{r=0}^{\infty} \left(\frac{\lambda}{\nu}\right)^{2r} \frac{1}{r!} \prod_{i=1}^r \left(\frac{1}{i + 2\frac{\lambda}{\nu}}\right) \left(\frac{-e^{-\kappa\nu}}{\lambda}\right)^r \\ &= \sum_{r=0}^{\infty} \frac{1}{r!} \left(\frac{-e^{-\kappa\nu}}{2\nu}\right)^r = e^{-\frac{1}{2}e^{-\kappa\nu}/\nu}. \end{aligned}$$

Note that this limit is the same as the value of $\phi(x_\lambda)$ in the case of exponential interarrival times with parameter $\lambda/2$ (cf. (12)).

4. An asymptotic result for renewal risk models with tax

Assume that potential ‘catastrophes’ occur according to a delayed renewal process with initial delay T_0 and interrenewal periods T_1, T_2, \dots . At time $S_n := T_0 + \dots + T_n$, an actual catastrophe occurs if V_n exceeds $f(S_n)$, with $f(\cdot)$ some increasing function, and V_0, V_1, V_2, \dots a sequence of independent and identically distributed random variables. The random variables T_{n+1} and V_n may be dependent. Let the 0–1 variable $I_n :=$

$I_{\{V_n > f(S_n)\}}$ indicate whether or not an actual catastrophe occurs at time S_n , and denote

$$p(t) := \mathbb{P}\{V_n > f(t)\}.$$

We are interested in the probability of the event E_τ that no actual catastrophe occurs during the time interval $[0, \tau]$, i.e.,

$$E_\tau = \cup_{n=-1}^{\infty} \{S_n \leq \tau < S_{n+1}; I_0 = \dots = I_n = 0\},$$

with the notational convention that $S_{-1} := 0$.

Now consider the surplus process in the Sparre Andersen risk model where claims of generic size Y occur according to a renewal process with generic interrenewal time X , and a marginal tax rate γ applies whenever the free surplus is at a running maximum. Let Q be a single-server queue with generic interarrival time X and generic service time Y . Let V_{\max} and T be a pair of random variables with as joint distribution that of the maximum workload during a busy period of Q and the subsequent idle period. Further suppose that we take the joint distribution of T_{n+1} and V_n to be that of T and V_{\max} , and $f(t) = x + (1 - \gamma)t$. Then the probability of the event E_τ with $\tau = (v - x)/(1 - \gamma)$ equals the probability $\phi_\gamma(x, v)$ that the surplus process reaches level v , starting from level x , before ruin occurs. In particular, the survival probability in the renewal model with tax is $\phi_\gamma(x) = \mathbb{P}\{E_\infty\}$, with $E_\infty = \{V_n \leq x + (1 - \gamma)S_n \text{ for all } n = 0, 1, 2, \dots\}$.

Remark 5. Following Section 3, the probability of the event E_∞ may also be interpreted as the probability that no customer with a remaining service time exceeding x is present in a $G/G/\infty$ system where the joint distribution of the interarrival time and subsequent service time is that of $(1 - \gamma)T_{n+1}$ and V_n , given that the past interarrival time is T_0 .

In order to characterize the probability of interest, i.e., $\mathbb{P}\{E_\tau\}$, we will consider a scenario where the interrenewal periods are relatively short (compared to the time interval $[0, \tau]$), i.e., the number of potential catastrophes is relatively large, while the probability that an actual catastrophe occurs is relatively small, such that the value of the ratio $p(t)/\mathbb{E}\{T\}$ is moderate. More specifically, we assume an asymptotic regime

where time is accelerated by a factor s , i.e., with interrenewal periods $T^{(s)} := T/s$, while the function $f^{(s)}(\cdot)$ is simultaneously boosted in such a manner that the ratio $p^{(s)}(t)/\mathbb{E}\{T^{(s)}\} = p(t)/\mathbb{E}\{T\}$, i.e., $p^{(s)}(t) = p(t)/s$. For each fixed value of s , denote the resulting event E_τ by $E_\tau^{(s)}$.

The next theorem states the main result of this section.

Theorem 4.1. *Under the above-mentioned assumptions,*

$$\mathbb{P}\left\{E_\tau^{(s)}\right\} \rightarrow \exp\left(-\lambda \int_{t=0}^{\tau} p(t)dt\right) \quad (14)$$

as $s \rightarrow \infty$, with $\lambda := 1/\mathbb{E}\{T\}$.

Remark 4.1. Theorem 4.1 suggests that the expression on the right-hand side should provide a reasonable approximation for $\mathbb{P}\left\{E_\tau^{(s)}\right\}$ in the above-described asymptotic regime where the interrenewal periods are relatively short compared to the time interval $[0, \tau]$. Note that (14) has a similar form as the earlier result (1) for the Cramér-Lundberg risk process.

In order to prove Theorem 4.1, we will establish lower and upper bounds for the unscaled process. Lemmas 4.2 and 4.4 then show that these two bounds, while crude, coincide in the asymptotic regime under consideration.

For compactness, we henceforth drop the subscript τ from the notation $E_\tau^{(s)}$, and simply write $E^{(s)}$ or just E . Note that

$$\lim_{K \rightarrow \infty} \frac{\tau}{K} \sum_{k=1}^K p\left(k \frac{\tau}{K}\right) = \lim_{K \rightarrow \infty} \frac{\tau}{K} \sum_{k=1}^K p\left(\left(k-1\right) \frac{\tau}{K}\right) = \int_{t=0}^{\tau} p(t)dt. \quad (15)$$

Let us now focus on the lower bound. Let $K \geq 1$ and $N \geq 1$ be integers and $t_0 = 0 \leq t_1 \leq \dots \leq t_K = \tau$. For any $k = 1, \dots, K$, define the events

$$D_k := \{S_{kN} > t_k\},$$

$$F_k := \{V_{(k-1)N} \leq f(t_{k-1}), \dots, V_{kN-1} \leq f(t_{k-1})\},$$

and

$$E^{\text{lower}} := \bigcap_{k=1}^K D_k \cap \bigcap_{k=1}^K F_k.$$

Lemma 4.1. *The event E^{lower} implies the event E .*

Proof. Suppose that the event E^{lower} occurs, i.e., all the events D_k and F_k occur. Let i be such that $(k-1)N \leq i \leq kN-1$ for some $k=1, \dots, K$. The event D_k gives $S_i \geq S_{(k-1)N} > t_{k-1}$, while the event F_k implies $V_i \leq f(t_{k-1})$. Since the function $f(\cdot)$ is increasing, it follows that $V_i \leq f(S_i)$. Hence $I_i = 0$ for all $i=0, \dots, KN-1$. The event D_K implies that there exists an $n \leq KN-1$ with $S_n \leq \tau < S_{n+1}$. Thus the event E occurs. \square

Lemma 4.2.

$$\lim_{s \rightarrow \infty} \mathbb{P} \left\{ E^{(s)} \right\} \geq e^{-\lambda \int_{t=0}^{\tau} p(t) dt}. \quad (16)$$

Proof. Lemma 4.1 yields that

$$\begin{aligned} \mathbb{P} \{E\} &\geq \mathbb{P} \{E^{\text{lower}}\} = \mathbb{P} \left\{ \bigcap_{k=1}^K D_k \cap \bigcap_{k=1}^K F_k \right\} \geq \mathbb{P} \left\{ \bigcap_{k=1}^K F_k \right\} - \mathbb{P} \left\{ \overline{\bigcap_{k=1}^K D_k} \right\} \\ &\geq \prod_{k=1}^K \mathbb{P} \{F_k\} - \sum_{k=1}^K \mathbb{P} \{\overline{D}_k\} \\ &= \prod_{k=1}^K \mathbb{P} \{V_{(k-1)N} \leq f(t_{k-1}), \dots, V_{kN-1} \leq f(t_{k-1})\} - \sum_{k=1}^K \mathbb{P} \{S_{kN} \leq t_k\} \\ &= \prod_{k=1}^K (\mathbb{P} \{V \leq f(t_{k-1})\})^N - \sum_{k=1}^K \mathbb{P} \{S_{kN} \leq t_k\}. \end{aligned}$$

Choose now $N = \lceil N(s) \rceil$, with $N(s) = (1 + \epsilon) \frac{\tau s}{K \mathbb{E}\{T\}}$, and $t_k = \frac{k\tau}{K}$, $k=1, \dots, K$.

Then

$$\begin{aligned} \mathbb{P} \{S_{kN} \leq t_k\} &= \mathbb{P} \left\{ T_0/s + T_1/s + \dots + T_{\lceil N(s) \rceil} / s \leq \frac{k\tau}{K} \right\} \\ &= \mathbb{P} \left\{ T_0 + T_1 + \dots + T_{\lceil N(s) \rceil} \leq \frac{kN(s)\mathbb{E}\{T\}}{1 + \epsilon} \right\}, \end{aligned}$$

which by the law of large numbers tends to zero as $s \rightarrow \infty$. Also,

$$\begin{aligned} \lim_{s \rightarrow \infty} \prod_{k=1}^K (\mathbb{P} \{V < f(t_{k-1})\})^{N(s)} &= \lim_{s \rightarrow \infty} \prod_{k=1}^K e^{-N(s)p^{(s)}(t_{k-1})} = e^{-\sum_{k=1}^K \lim_{s \rightarrow \infty} N(s)p^{(s)}(t_{k-1})} \\ &= e^{-\sum_{k=1}^K \frac{\tau p(t_{k-1})}{K \mathbb{E}\{T\}}} = e^{-\frac{\tau}{K \mathbb{E}\{T\}} \sum_{k=1}^K p(t_{k-1})}. \end{aligned}$$

We deduce that

$$\lim_{s \rightarrow \infty} \mathbb{P} \left\{ E^{(s)} \right\} \geq e^{-\frac{\tau}{K \mathbb{E}\{T\}} \sum_{k=1}^K p(t_{k-1})}$$

for any $K \geq 1$. Letting $K \rightarrow \infty$ and applying (15), we obtain the lower bound (16). \square

Next, we establish an upper bound that asymptotically matches the lower bound. Let $K \geq 1$ and $N \geq 1$ be integers and $t_0 = 0 \leq t_1 \leq \dots \leq t_K = \tau$. For any $k = 1, \dots, K$, define the events

$$G_k := \{V_{(k-1)N} \leq f(t_k), \dots, V_{kN-1} \leq f(t_k)\},$$

and

$$E^{\text{upper}} := \bigcup_{k=1}^K D_k \cup \bigcap_{k=1}^K G_k.$$

Lemma 4.3. *The event E implies the event E^{upper} .*

Proof. Suppose that the event E occurs, i.e., there exist an $n(\tau)$ with $S_{n(\tau)} \leq \tau < S_{n(\tau)+1}$ and $I_0 = \dots = I_{n(\tau)} = 0$. Also assume that all the events \bar{D}_k occur, i.e., $S_{kN} \leq t_k$ for all $k = 1, \dots, K$, because otherwise there is nothing to prove. This in particular implies that $n(\tau) \geq KN - 1$, and hence $I_0 = \dots = I_{KN-1} = 0$, i.e., $V_i \leq f(S_i)$ for all $i = 0, \dots, KN - 1$. Let i be such that $(k-1)N \leq i \leq kN - 1$ for some $k = 1, \dots, K$, so that $S_i \leq S_{kN}$. Since the function $f(\cdot)$ is increasing, it follows that $V_i \leq f(t_k)$, and thus all the events G_k occur, and hence the event E^{upper} occurs. \square

Lemma 4.4.

$$\lim_{s \rightarrow \infty} \mathbb{P} \left\{ E^{(s)} \right\} \leq e^{-\lambda \int_{t=0}^{\tau} p(t) dt}. \quad (17)$$

Proof. Lemma 4.3 yields that

$$\begin{aligned} \mathbb{P} \{E\} &\leq \mathbb{P} \{E^{\text{upper}}\} \\ &= \mathbb{P} \left\{ \bigcup_{k=1}^K D_k \cup \bigcap_{k=1}^K G_k \right\} \\ &\leq \mathbb{P} \left\{ \bigcap_{k=1}^K G_k \right\} + \mathbb{P} \left\{ \bigcup_{k=1}^K D_k \right\} \\ &\leq \prod_{k=1}^K \mathbb{P} \{G_k\} + \sum_{k=1}^K \mathbb{P} \{D_k\} \\ &= \prod_{k=1}^K (\mathbb{P} \{V \leq f(t_k)\})^N + \sum_{k=1}^K \mathbb{P} \{S_{kN} > t_k\}. \end{aligned}$$

We now take $N = \lceil N(s) \rceil$, with $N(s) = (1 - \epsilon) \frac{\tau s}{K \mathbb{E}\{T\}}$, and $t_k = \frac{k\tau}{K}$, $k = 1, \dots, K$, and proceed to evaluate the above upper bound in the asymptotic regime of interest. Note

that

$$\begin{aligned} \mathbb{P}\{S_{kN} > t_k\} &= \mathbb{P}\left\{T_0/s + T_1/s + \cdots + T_{k\lceil N(s)\rceil}/s > \frac{k\tau}{K}\right\} \\ &= \mathbb{P}\left\{T_0 + T_1 + \cdots + T_{k\lceil N(s)\rceil} > \frac{kN(s)\mathbb{E}\{T\}}{1-\epsilon}\right\}, \end{aligned}$$

which tends to zero as $s \rightarrow \infty$ because of the law of large numbers. Also,

$$\begin{aligned} \lim_{s \rightarrow \infty} \prod_{k=1}^K (\mathbb{P}\{V \leq f(t_k)\})^{N(s)} &= \lim_{s \rightarrow \infty} \prod_{k=1}^K e^{-N(s)p^{(s)}(t_k)} = e^{-\sum_{k=1}^K \lim_{s \rightarrow \infty} N(s)p^{(s)}(t_k)} \\ &= e^{-\sum_{k=1}^K \frac{\tau p(t_k)}{K\mathbb{E}\{T\}}} = e^{-\frac{\tau}{K\mathbb{E}\{T\}} \sum_{k=1}^K p(t_k)}. \end{aligned}$$

We conclude that

$$\lim_{s \rightarrow \infty} \mathbb{P}\{E^{(s)}\} \leq e^{-\frac{\tau}{K\mathbb{E}\{T\}} \sum_{k=1}^K p(t_k)}$$

for any $K \geq 1$. Letting $K \rightarrow \infty$ and invoking (15), we obtain the upper bound (17).

□

Acknowledgement

H. Albrecher would like to acknowledge support from the Swiss National Science Foundation Project 200021-124635/1.

References

- [1] H. Albrecher, A. Badescu and D. Landriault (2008) On the dual risk model with taxation. *Insurance Mathematics and Economics* **42**, 1086–1094.
- [2] H. Albrecher, S.C. Borst, O.J. Boxma and J.A.C. Resing (2009) The tax identity in risk theory – a simple proof and an extension. *Insurance Mathematics and Economics* **44**, 304–306.
- [3] H. Albrecher and C. Hipp (2007) Lundberg’s risk process with tax. *Bl. DGVFM*, **28**(1):13–28, 2007.
- [4] H. Albrecher, J. Renaud and X. Zhou (2008) A Levy insurance risk process with tax. *Journal of Applied Probability* **45**, 363–375.
- [5] S. Asmussen and H. Albrecher. *Ruin Probabilities*. 2nd edn. World Scientific, New Jersey, 2010.
- [6] J.M. Harrison and S.I. Resnick (1976) The stationary distribution and first exit probabilities of a storage process with general release rule. *Math. Oper. Res.* **1**, 347–358.

- [7] A. Kyprianou and X. Zhou (2009) General tax structures and the Lévy insurance risk model. *Journal of Applied Probability* **46**, 1146–1156.
- [8] P. Picard (1994) On some measures of the severity of ruin in the classical Poisson model. *Insurance Mathematics and Economics* **14**, 107–115.
- [9] L. Takács (1962) *Introduction to the Theory of Queues*. Oxford University Press, New York.
- [10] J. Wei, H. Yang and R. Wang (2010) On the Markov-modulated insurance risk model with tax. *Bl. DGVFM* **31**(1), 65–78.