

Mimicking the satellite cell niche

Citation for published version (APA):

Boonen, K. J. M., Baaijens, F. P. T., & Post, M. J. (2007). *Mimicking the satellite cell niche*. Poster session presented at Mate Poster Award 2007 : 12th Annual Poster Contest.

Document status and date:

Published: 01/01/2007

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Mimicking the satellite cell niche

K.J.M. Boonen, F.P.T. Baaijens, M.J. Post

Eindhoven University of Technology, Department of Biomedical Engineering

Introduction

Because of their effectiveness in regenerating large muscle defects *in vivo*, we investigate the use of satellite cells derived from muscle biopsies as a cell source for muscle tissue engineering. Despite their *in vivo* potency, however, their proliferative and myogenic capacity *in vitro* has been disappointing.¹

We therefore concentrate on optimizing culture conditions by mimicking the natural niche environment of these cells. The elasticity of the environment is an important aspect of this niche and has been shown to influence differentiation pathways of stem cells². We hope to improve the *in vitro* capacities of satellite cells by varying substrate elasticity. The potential for myogenesis is evaluated from morphological and functional parameters.

The model system used in these experiments provides for high-throughput and well-controlled measurements and can ultimately be expanded to 3D tissue culture conditions using biomaterials and/or different cell types.

Material and methods

Tibialis Anterior (TA) muscles from Swiss White mice are digested in collagenase and triturated to release single fibres from which satellite cells are liberated with a 19G needle. The satellite cells are plated on Matrigel™-coated polyacrylamide (PA) gels (Figure 1) with elastic moduli (E) ranging from 1.4 to 72 kPa on top of 13 mm ø coverslips and cultured in plating medium (DMEM advanced, 10% HS, 20% FBS, 0.5% CEE, 1% pen/strep, 1% L-glut.).




Figure 1: Creation of coated PA gels on top of coverslips

Results

- Satellite cells proliferate on all substrates
- Fusion first takes place on 17 kPa, which mimics physiological elasticity of striated muscle.
- Subsequently, satellite cells fuse on 8.1 and 35 kPa
- Eventually fusion takes place on all E-moduli




Figure 2: Proliferation and differentiation of satellite cells on substrates with varying elasticity

Conclusions

Timing of differentiation of satellite cells seems to be dependent on the elasticity of their environment. Fusion first takes place on physiological elasticity of striated muscle, subsequently at the elastic moduli closest to this physiological value and lastly at the outer values.

Overall, we have developed a robust tissue culture system, that gives us the possibility to pursue long term proliferation of satellite cell derived progenitor cells and evaluate their proliferative and differentiative capacities.

In the future, different biochemical and biophysical cues will be used to influence and optimize these processes.

References:

- [1] Collins, C. A., I. Olsen, et al. (2005). "Stem Cell Function, Self-Renewal, and Behavioral Heterogeneity of Cells from the Adult Muscle Satellite Cell Niche." *Cell* 122(2): 289-301.
- [2] Engler, A. J., S. Sen, et al. (2006). "Matrix elasticity directs stem cell lineage specification." *Cell* 126(4): 677-89.