

Pressure-related deep tissue injury coincides with areas of high maximum shear strains

Citation for published version (APA):

Ceelen, K. K., Stekelenburg, A., Baaijens, F. P. T., Bader, D. L., Nicolaij, K., & Oomens, C. W. J. (2007). *Pressure-related deep tissue injury coincides with areas of high maximum shear strains*. Poster session presented at Mate Poster Award 2007 : 12th Annual Poster Contest.

Document status and date:

Published: 01/01/2007

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Pressure-related deep tissue injury coincides with areas of high maximum shear strains

K.K. Ceelen, A. Stekelenburg, F.P.T. Baaijens, K. Nicolay, D.L. Bader, C.W.J. Oomens

Biomechanics and Tissue Engineering, Soft Tissue Biomechanics & Engineering

Introduction

Deep tissue injury has recently been recognized as a specific type of pressure ulcer, and was defined as pressure-related damage to subcutaneous tissues under an intact skin. These wounds often require long hospitalization because of poor healing and they seriously decrease quality of life.

Prevention and early detection are hampered by incomplete understanding of the underlying damage pathways. Previous animal experiments demonstrated that the ischemic area was much larger than the damaged area and suggested that strains were directly responsible for tissue injury (Stekelenburg et al. 2007):




Figure 1: MR-image of cross-section of rat hindlimb during 2h compression (left), perfusion during compression (blue = low, red = high perfusion)(center), T2-weighted MR-image 4h after unloading (high intensity = damage)(right).


Is there a relation between tissue deformation and deep tissue injury?

MR-tagging can be used to measure internal deformations during compression, but these measurements provoke damage. Therefore they cannot be combined with measuring damage due to compression. A dedicated 2D plane stress finite element model was developed to calculate experiment-specific internal strains and compare them with measured damage. MR-tagging experiments were used to validate this model.

Validation

Numerical and tagging displacements were interpolated on a grid in the region of interest (red square in figure 1) and then processed to obtain strains. Correlation between numerical and experimental maximum shear strains and strain energy densities was good:

Figure 3: Correlation of model and tagging strain quantities ($n = 4$).



Damage correlation

The model was then used to simulate damage experiments and the overlap between damage locations and maximum compressive and shear strains was analyzed:




Figure 4: Maximum shear strains in region of interest (red square in figure 1) of 2 different experiments with damage (white cross/black circle) and deformed contours (black curves) superimposed.

The area of high maximum shear strains coincides with the damage, and the amount of damage increases as the size of the area with high maximum shear strains increases:




Figure 5: Relative damaged area in each category ($n = 9$).

Yes, internal shear strains cause deep tissue injury.

- High shear strains cause muscle damage within 2 hours, while ischemia does not cause damage until after 4 hours.
- Defining a strain damage threshold would be a major leap forward in prevention and early detection of deep tissue injury.