

Momentum transport in TCV across sawteeth events

Citation for published version (APA):

Duval, B. P., Bortolon, A., Federspiel, L., Felici, F., Furno, I., Goodman, T. P., Karpushov, A., Paley, J. I., & Piras, F. (2010). *Momentum transport in TCV across sawteeth events*. 191-191.

Document status and date:

Published: 01/01/2010

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EXS/P4 — Poster Session

EXS/P4-01

Momentum Transport In TCV Across Sawteeth Events

Duval, B.P.¹, Bortolon, A.¹, Federspiel, L.¹, Furno, I.¹, Karpushov, A.¹, Paley, J.¹, Piras, F.¹, and TCV Team¹

¹*Ecole Polytechnique Fédérale de Lausanne-CRPP-EPEL, Association Euratom-Confédération Suisse-CH-1015 Lausanne, Switzerland*

Corresponding Author: basil.duval@epfl.ch

The scaling of toroidal rotation has been studied over a wide range of in plasma and shape parameters on the TCV tokamak using a diagnostic neutral beam that injects a negligible momentum into the plasma. As on other Tokamaks, the toroidal rotation magnitude on TCV, for a given shape and plasma conditions, is observed to scale inversely with the plasma current. More specifically, this describes the maximum toroidal rotation in plasmas with strong sawteeth activity. The toroidal rotation gradient from the plasma edge was similar over a wide range of plasma currents and only diverged inside the sawtooth inversion radius. By extrapolating the maximum core rotation from the rotation profiles outside the sawtooth inversion radii and adding a correction for the observed ion temperature changes, the scaling of the intrinsic toroidal rotation with plasma current is no longer observed.

The TCV CXRS diagnostic was configured to measure the toroidal rotation profile with a low spatial resolution (4 points across the radial co-ordinate) and a high temporal resolution (2 ms). Using ECH X2 power deposited close to the sawteeth inversion radius, the sawteeth repetition time was extended to over 12 ms. The measurement used a newly available Real-Time node to generate a sequence of trigger pulses from analysis of the soft X-ray such that each spectroscopic frame was taken at 2, 4, 6, 8, ... ms after each sawtooth. The radial toroidal rotation profile evolution over a sawtooth was determined using coherent resampling over many sawtooth events.

Initial results indicate that, at the sawtooth crash, the plasma core accelerates in the co current direction and then the core rotation profile relaxes back to the counter current direction with a time constant ~ 10 ms. This implies that the scaling of intrinsic rotation with plasma current may be better understood as a peaked rotation profile (or an inwards momentum pinch) in the plasma core region that is flattened by sawtooth activity. This change of viewpoint should be integrated into the theory of momentum generation and transport in tokamaks such as ITER in that these mechanisms are not as universally affected by the plasma current as was at first thought.