

Magnetic resonance techniques to study damage evolution in muscle

Citation for published version (APA):

Stekelenburg, A., Oomens, C. W. J., Bouten, C. V. C., Bader, D. L., & Nicolaij, K. (2002). *Magnetic resonance techniques to study damage evolution in muscle*. Poster session presented at Mate Poster Award 2002 : 7th Annual Poster Contest.

Document status and date:

Published: 01/01/2002

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Magnetic Resonance Techniques to study damage evolution in muscle

A.Stekelenburg, C.W.J.Oomens, C.V.C.Bouten, D.L.Bader, K.Nicolay

Eindhoven University of Technology, Department of Biomedical Engineering

Introduction

Pressure sores are localised areas of tissue breakdown in soft tissues. They are primarily caused by prolonged mechanical loading. It is not known how these external mechanical loads result in tissue damage.

A new rat model was developed by Bosboom [1] in which the tibialis anterior muscle was compressed between an indenter and the tibia. It was demonstrated that T2-weighted MRI is a promising alternative for histological examination to evaluate muscle tissue. However, the model contained too many inaccuracies to study the influence of duration and magnitude of pressure and the influence of (re)perfusion on the development of muscle damage.

Objective

To study damage thresholds, perfusion and deformation of soft tissue during and after mechanical loading using Magnetic Resonance techniques

Methods

To be able to simultaneously collect MR data and apply pressure to the muscle, a MR-compatible loading apparatus is being developed, in which an anaesthetized rat can be placed.




Figure 1 Loading device, front and side view

The hind leg of the rat is placed in a radio-frequency coil. A water-filled indenter (which makes it visible on the MR images, see figure 2) can be moved through the legs of the coil.




Figure 2 Indentor visible on MR image

The force will be applied by a linear actuator and can be applied in less than 100 ms, which allows for tracking muscle deformation using MRI tagging methods. The deformation will be compared to finite element modelling of tissue deformation.




Figure 3 Different positions of RF coil which makes fixation of the hind limb easier

MRI measurements will be complemented with 1H- and 31P-MRS studies of the metabolic status of the tissue. 1H-MRS will yield data on lactate formation (as an index of anaerobic glycolysis), while 31P-MRS will provide information on tissue energy charge and intracellular pH, both of which are expected to decline during prolonged loading.

Discussion

The local loading conditions will be carefully controlled and measured for each individual animal. Finite element modeling will be used to determine the mechanical conditions within the muscles during loading. Critical material input parameters for the finite element modeling will be deduced from the MRI data.




Figure 4 Three dimensional finite element mesh of muscle, skin and indenter

Conclusions

With this new set-up we will be able to learn more about the role of deformation, perfusion and damage mechanics in the aetiology of pressure sores.

References:

- [1] BOSBOOM, 2001: *Deformation as a trigger for pressure sore related muscle damage*