

Exponential-uniform identities related to records

Citation for published version (APA):

Gnedin, A. V., & Marynych, A. (2012). *Exponential-uniform identities related to records*. (Report Eurandom; Vol. 2012021). Eurandom.

Document status and date:

Published: 01/01/2012

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EURANDOM PREPRINT SERIES

2012-021

November, 2012

Exponential Uniform Identities Related to Records

Alexander Gnedin, Alexander Marynych

ISSN 1389-2355

Exponential–Uniform Identities Related to Records

Alexander Gnedin* and Alexander Marynych†

Abstract

We consider a rectangular grid induced by the south-west records from the planar Poisson point process in \mathbb{R}_+^2 . A random symmetry property of the matrix whose entries are the areas of tiles of the grid implies cute multivariate distributional identities for certain rational functions of independent exponential and uniform random variables.

Keywords: records, planar Poisson process, distributional identities

AMS 2000 Subject Classification: Primary 60G70, 60E99

1. Introduction Let E_1, E_2, \dots and U_1, U_2, \dots be two independent sequences of independent rate-one exponential and $[0, 1]$ -uniform random variables, respectively. A prototype of the distributional identities appearing in this note is the identity

$$\left(\frac{E_1}{U_1} + \frac{E_2}{U_1 U_2} + \dots + \frac{E_n}{U_1 \dots U_n} \right) (1 - U_1 \dots U_{n+1}) \stackrel{d}{=} \left(E_1 + \frac{E_2}{U_1} + \dots + \frac{E_{n+1}}{U_1 \dots U_n} \right) (1 - U_1 \dots U_n), \quad (1)$$

which was used in [4] to explain coincidence of the values in two quite different problems of optimal stopping. Some probabilities related to the simplest instance of (1),

$$\frac{E_1}{U_1} (1 - U_1 U_2) \stackrel{d}{=} \left(E_1 + \frac{E_2}{U_1} \right) (1 - U_1), \quad (2)$$

had been evaluated in [7].

We will show that (1) along with more general identities for matrix functions in the exponential and uniform variables follow from symmetry properties of the set of *records* (also known as Pareto-extremal points [2]) from the planar Poisson process in the positive quadrant. This continues the line of [5], where it was argued that the planar Poisson process is a natural framework for two gems of combinatorial probability: Ignatov’s theorem [6] and the Arratia-Barbour-Tavaré lemma on the scale-invariant Poisson processes on \mathbb{R}_+ [1].

We shall evaluate the areas of tiles for a rectangular grid induced by the set of records. The identities obtained in this way are genuinely multivariate, albeit they stem from the arrays with identical marginal distributions. In particular, (2) appears by a row summation in

$$\begin{pmatrix} (1 - U_1)E_1 & (1 - U_1)\frac{E_2}{U_1} \\ U_1(1 - U_2)E_1 & (1 - U_2)E_2 \end{pmatrix} \stackrel{d}{=} \begin{pmatrix} (1 - U_2)E_2 & (1 - U_1)\frac{E_2}{U_1} \\ U_1(1 - U_2)E_1 & (1 - U_1)E_1 \end{pmatrix}. \quad (3)$$

2. A random tiling induced by records We use the self-explaining notations $\nearrow, \searrow, \swarrow, \nwarrow$ for four coordinate-wise partial orders on the positive quadrant. For instance, relations $a \nearrow b$ and $b \swarrow a$ for $a, b \in \mathbb{R}_+^2$ both mean that b is located strictly north-east of a .

Let \mathcal{P} be the planar Poisson point process with unit intensity in \mathbb{R}_+^2 . It will be convenient to understand \mathcal{P} as a random set, rather than counting measure. The event $(t, x) \in \mathcal{P}$ is interpreted as the

*Queen Mary University of London, email: a.gnedin@qmul.ac.uk

†Eindhoven University of Technology, email: O.Marynych@tue.nl. The research of AM was financially supported by a Free Competition Grant of the Netherlands Organisation for Scientific Research (NWO).




Figure 1: The rectangular tiling and areas with the same distribution.

value x observed at time t . Note that with probability one no two atoms of \mathcal{P} lie on the same vertical or horizontal line. An atom $r \in \mathcal{P}$ is said to be a (lower) *record* if there is no earlier observation with a smaller value, that is $a \nearrow r$ for no $a \in \mathcal{P}$. The set of records, denoted henceforth \mathcal{R} , is a point process with the intensity function e^{-tx} . The collection of records ordered by increase of the time component is a two-sided infinite \searrow -chain

$$\dots \searrow r_{-2} \searrow r_{-1} \searrow r_0 \searrow r_1 \searrow r_2 \searrow \dots,$$

which we label by nonzero integers in such a way that the records r_0 and r_1 are separated by the bisectrix $t = x$.

Drawing vertical lines at t -locations of records (record times), and drawing horizontal lines at their x -locations (record values) divides the positive quadrant into rectangular tiles. Let C_{ij} be the area of the tile whose north-west corner is the intersection point of the horizontal line through r_i and the vertical line through r_j (see Figure 1). In particular, C_{ii} for $i \in \mathbb{Z}$ is the area of a tile spanned on records r_i, r_{i+1} .

Given a record at location (t, x) , the next record is just the next point of \mathcal{P} south-west of (t, x) , hence distributed like $(t + E/x, xU)$, as is easily seen from the independence and homogeneity properties of \mathcal{P} . The sequence r_1, r_2, \dots is Markovian with just described transitions, whence

$$(C_{ij}; i, j = 1, 2, \dots) \stackrel{d}{=} \left(U_1 \cdots U_{i-1} (1 - U_i) \frac{E_j}{U_1 \cdots U_{j-1}}; i, j = 1, 2, \dots \right). \quad (4)$$

Note that the left-hand side of (4) is independent of r_1 . Since the law of \mathcal{R} is not changed by reflection about the bisectrix, we also have

$$(C_{i,j}; i, j = 1, 2, \dots) \stackrel{d}{=} (C_{-j-1, -i-1}; i, j = 1, 2, \dots). \quad (5)$$

As an illustration, the areas of two rectangles (Figure 1) spanned on r_1, r_4 and r_{-3}, r_0 , respectively, have the same distributions.

For $n \in \mathbb{N}$ and $k \in \mathbb{Z}/\{0\}$ let $M_{k,n} = (C_{k+i-1, k+j-1}, i, j = 1, \dots, n)$ be the $n \times n$ matrix associated with records $r_k, r_{k+1}, \dots, r_{k+n}$. Obviously from the above, $M_{k,n}$ is independent of r_k and satisfies $M_{k,n} \stackrel{d}{=} M_{1,n}$ for $k = 1, 2, \dots$. For $k < 0$, $M_{k,n}$ is not independent of r_k , since the transition probability from $r_k = (t, x)$ to r_{k+1} accounts for the condition that $-k$ records must lie south-east of (t, x) above the bisectrix. Also, $M_{k,n} \stackrel{d}{=} M_{1,n}$ fails for $-n < k \leq 0$: for instance, $C_{0,0}$ and $C_{1,1}$ have different distributions. Nevertheless, we will show that $M_{k,n} \stackrel{d}{=} M_{1,n}$ holds for $k \leq -n$, which by virtue of (5) is equivalent to the following random symmetry property of $M_{1,n}$.

Let $M_{1,n}^*$ be the matrix obtained by reflecting $M_{1,n}$ about the antidiagonal, that is by exchanging each entry (i, j) with entry $(n - j + 1, n - i + 1)$.

Proposition 1 For $n = 1, 2, \dots$

$$M_{1,n}^* \stackrel{d}{=} M_{1,n}. \quad (6)$$

Identity (3) is the $n = 2$ instance of (6). Identity (1) appears by calculating the sum of all entries of $M_{1,n+1}$ except the entries in the $(n + 1)$ st row. Further identities can be derived by applying functions, e.g., taking the product of matrix elements in the first row of $M_{1,n}$:

$$\frac{E_1 \cdots E_n (1 - U_1)^n}{U_1^{n-1} U_2^{n-2} \cdots U_{n-1}} \stackrel{d}{=} \frac{E_n^n (1 - U_1) \cdots (1 - U_n)}{U_1 U_2^2 \cdots U_{n-1}^{n-1}}.$$

3. Records in a finite box To prove (6) we consider records in finite rectangles. Let $A \subset \mathbb{R}_+^2$ be a finite open rectangle with sides parallel to the coordinate axes. Atom $a \in \mathcal{P} \cap A$ will be called A -record if no other atom $b \in \mathcal{P} \cap A$ lies south-west of a . The set of A -records induces a random partition of A in rectangular tiles. Denote by $N = (N_{i,j})$ the random matrix of areas of the tiles. The number of rows (or columns) of N is a random variable that is one plus the number of A -records. Let $N^* = (N_{i,j}^*)$ be the array obtained by reflecting N about the antidiagonal, which is defined conditionally on the number of A -records.

Lemma 2 For every rectangle A we have

- (i) $N \stackrel{d}{=} N^*$,
- (ii) also $N \stackrel{d}{=} N^*$ conditionally given the number of A -records is n , for each $n \geq 0$,
- (iii) the distribution of N depends on A only through the area of A .

Proof. Applying a hyperbolic shift $(t, x) \mapsto (\lambda t, x/\lambda)$ with some $\lambda > 0$, rectangle A can be mapped onto a square. The mapping preserves the area and the coordinate-wise orders, hence preserves the distribution of N . For A a square (i) and (ii) follow by symmetry of A about the north-east diagonal (see Figure 2).

Lemma 3 For $n \geq 1$ the following conditional distributions coincide:

- (a) the distribution of $M_{k,n}$ given the area v of the rectangle spanned on records r_k and r_{n+k} , where $k \geq 1$ or $k \leq -n - 1$,
- (b) the distribution of N for a rectangle A of area v , given that the number of A -records is $n - 1$.

Proof. For any fixed rectangle A , the set of A -records is independent of the Poisson point process outside A . On the other hand, given that the north-west and the south-east corners of A are records, \mathcal{P} has no atoms south-east of these corners, hence the set of A -records coincides with $\mathcal{R} \cap A$. That is to say, given that two records are located at the corners of A , the records inside A are distributed like A -records. In the same way, taking, for instance, $k = 1$, we have: given that r_1 and r_{n+1} are at the corners of rectangle




Figure 2: Records in a square

A (below the line $x = t$), the set $\{r_2, \dots, r_n\}$ has the same distribution as the set of A -atoms, conditioned on the event that the number of A -records is $n - 1$. Now the assertion follows from Lemma 2 (iii).

Proving Proposition 1 is now easy. Combining Lemma 3 with Lemma 2 (ii) we see that the distributional identity (6) holds conditionally on the area of the rectangle spanned on r_1 and r_{n+1} , hence (6) also holds unconditionally. Note that the area is equal to the sum of all entries of the $n \times n$ matrix, that is distributed like

$$(1 - U_1 \cdots U_n) \left(E_1 + \frac{E_2}{U_1} + \frac{E_3}{U_1 U_2} + \cdots + \frac{E_n}{U_1 \cdots U_{n-1}} \right).$$

We could not find a proof of (1) by computing densities or transforms, or by connecting to other known identities like ‘beta-gamma algebra’ [3]. Spanning a grid on the point process of k -corners [5] we were able to show that similar identities hold with uniform distribution replaced by beta($1, \theta$).

References

- [1] R. Arratia, A. Barbour and S. Tavaré(2006). A tale of three couplings: Poisson- Dirichlet and GEM approximations for random permutations, *Combin. Probab. Comput.* **15**, 31–62.
- [2] Yu. M. Baryshnikov (2000). Supporting-points processes and some of their applications. *Probab. Theory Related Fields*, **117**, 163-182.
- [3] D. Dufresne (1998). Algebraic properties of beta and gamma distributions, and applications, *Adv. Appl. Math.* **20**, 285-299.
- [4] A. Gnedin (2004) Best choice from the planar Poisson process, *Stoch. Proc. Applications* **111**, 317–354.
- [5] A. Gnedin (2008). Corners and records of the Poisson process in the quadrant, *Elect. Comm. in Probab.* **13**, 187–193.
- [6] C. M. Goldie and L. C. G. Rogers (1984). The k -records are i.i.d., *Probability Theory Rel Fields* **67**, 197–211.
- [7] Samuels, S.M. (2004). Why do these quite different best-choice problems have the same solutions? *Adv. Appl. Prob.* **36**, 398–416.