

A missing link in blending?

Citation for published version (APA):

Stegeman, Y. W., Chesters, A. K., Vosse, van de, F. N., & Meijer, H. E. H. (1999). *A missing link in blending?*. Poster session presented at Mate Poster Award 1999 : 4th Annual Poster Contest, .

Document status and date:

Published: 01/01/1999

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Introduction

Blend properties largely depend on flow history. CFD packages can predict this flow history and theories relating rheology with morphology exist. The link between flow history and morphology is still missing.

Objective

To investigate droplet behaviour in time-dependent flows by means of analytical, experimental and numerical methods.

Slender Body Model

Assumptions

- ◇ single, neutrally buoyant, Newtonian droplet
- ◇ axisymmetric elongational viscous Stokes flow
- ◇ droplet shape is a slender ellipsoid of revolution

Droplet elongation rate

The droplet elongation rate $\dot{\epsilon}_d = \dot{L}/L$ depends only on length L and capillary number $Ca = \eta_c R_0 \dot{\epsilon}_c / \sigma$ [1]:

$$\frac{\dot{\epsilon}_d}{\dot{\epsilon}_c} = \frac{L^3 + \frac{2\eta_c}{3\eta_d} + \frac{5}{12} - \frac{\sigma g}{f\eta_c R_0 \dot{\epsilon}_c} - \frac{\sigma g}{4\eta_d R_0 \dot{\epsilon}_c}}{L^3 + \frac{3}{4} + \frac{\eta_c}{2\eta_d} + \frac{4\eta_d}{3f\eta_c} - \frac{2}{3f}}$$

- L droplet length
- $\dot{\epsilon}_c$ cont. phase elongation rate
- $\dot{\epsilon}_d$ droplet elongation rate
- σ interfacial tension
- η_d droplet viscosity
- η_c continuous phase viscosity
- R_0 equivalent droplet radius
- f, g shape functions $f(L), g(L)$




fig. 1 Droplet elongation rate $\dot{\epsilon}_d(L, \dot{\epsilon}_c, \sigma, \eta_d, \eta_c, R_0)$

By integrating $\dot{\epsilon}_d$ (figure 1), the droplet deformation L in any time-dependent flow can be predicted, relating morphology with flow history.

References:

- [1] Y.W. STEGEMAN, A.K. CHESTERS, F.N. Vd VOSSE AND H.E.H. MEIJER, Proceedings PPS-15, 's-Hertogenbosch, 1999.
- [2] J.M.H. JANSSEN, *Dynamics of Liquid-Liquid Mixing*, PhD-thesis, Eindhoven University of Technology, 1993.

Results

Step in elongation rate $\dot{\epsilon}_c$

In figure 2, model predictions are compared with experimental results by Janssen [2]. The numbers denote the ratio between the imposed and the critical capillary number Ca_c (Ca_c : breakup in steady flow).




fig. 2 Droplet response to a step in the capillary number.

Triangular $\dot{\epsilon}_c$ - time profile

Predicted droplet response and experimental results obtained in a cross slot flow are shown in figure 3.




fig. 3 Droplet response to a triangular Ca -time profile.

Conclusions

The analytical model provides:

- ◇ adequate prediction of droplet deformation in time dependent flows
- ◇ link between morphology and flow history
- ◇ crucial input for CFD packages

Acknowledgement

This research is part of the FAIR project BioMix, funded by the European Community.