

## Faithful tropicalisation and torus actions

***Citation for published version (APA):***

Draisma, J., & Postinghel, E. (2014). *Faithful tropicalisation and torus actions*. (arXiv; Vol. 1404.4715 [math.AG]). s.n.

***Document status and date:***

Published: 01/01/2014

***Document Version:***

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

***Please check the document version of this publication:***

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

***General rights***

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

***Take down policy***

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

# FAITHFUL TROPICALISATION AND TORUS ACTIONS

JAN DRAISMA AND ELISA POSTINGHEL

ABSTRACT. For any affine variety equipped with coordinates, there is a surjective, continuous map from its Berkovich space to its tropicalisation. Exploiting torus actions, we develop techniques for finding an explicit, continuous section of this map. In particular, we prove that such a section exists for linear spaces, Grassmannians of planes (reproving a result due to Cueto, Hübich, and Werner), matrix varieties defined by the vanishing of maximal minors or of  $3 \times 3$ -minors, and for the hypersurface defined by Cayley's hyperdeterminant.

## 1. INTRODUCTION

Let  $K$  be a field with a non-Archimedean valuation  $v : K \rightarrow \mathbb{R}_\infty := \mathbb{R} \cup \{\infty\}$ , let  $\mathbb{A}^n \supseteq \mathbb{G}_m^n$  be the  $n$ -dimensional affine space over  $K$  and the  $n$ -dimensional torus with coordinates  $x_1, \dots, x_n$ , respectively, and let  $\mathbb{P}^{n-1}$  be the  $(n-1)$ -dimensional projective space over  $K$  with homogeneous coordinates  $x_1, \dots, x_n$ . For a closed subvariety  $X$  of  $\mathbb{G}_m^n$  or  $\mathbb{A}^n$  or  $\mathbb{P}^{n-1}$ , defined over  $K$ , we write  $\text{Trop}(X)$  for the tropicalisation of  $X$  sitting inside  $\mathbb{R}^n$  or  $\mathbb{R}_\infty^n$  or  $(\mathbb{R}_\infty^n \setminus \{(\infty, \dots, \infty)\})/\mathbb{R}(1, \dots, 1)$ , respectively.

Write  $X^{\text{an}}$  for the analytification of  $X$  in Berkovich's sense [Ber90, Chapter 1]. We work with the negative logarithms of multiplicative seminorms, so in the affine case  $X^{\text{an}}$  is the set of all ring valuations  $K[X] \rightarrow \mathbb{R}_\infty$  extending  $v$ , equipped with the topology of pointwise convergence. Write  $\infty$  for the valuation of  $K[\mathbb{A}^n] = K[x_1, \dots, x_n]$  that maps a polynomial to the valuation of its constant term. In the projective case, let  $\widehat{X} \subseteq \mathbb{A}^n$  be the affine cone over  $X$ . Then, as a topological space,  $X^{\text{an}}$  equals  $\widehat{X}^{\text{an}} \setminus \{\infty\}$  modulo the equivalence relation under which  $w_1$  and  $w_2$  are equivalent if and only if there exists a constant  $C \in \mathbb{R}$  such that for each degree- $d$  homogeneous polynomial  $f$  in the graded coordinate ring  $K[\widehat{X}]$  we have  $w_1(f) = dC + w_2(f)$ .

There is a continuous surjection

$$\pi : X^{\text{an}} \rightarrow \text{Trop}(X), \quad w \mapsto (w(x_1), \dots, w(x_n)).$$

This can be taken either as a definition of  $\text{Trop}(X)$  or as a theorem when other definitions are chosen [EKL06, Pay09, Dra08]. Indeed, in [Pay09] it is proved that  $X^{\text{an}}$  is the projective limit of the tropicalisations  $\text{Trop}(X)$  for all choices of coordinates. The tropicalisation is the support of a finite polyhedral complex by [BG84].

In this paper we discuss a number of high-dimensional examples where  $\pi$  has a continuous section. The results are motivated by exciting recent work for Grassmannians of planes [CHW] and for curves [BPR]. In particular, we will give another,

---

JD is supported by a Vidi grant from the Netherlands Organisation for Scientific Research (NWO).

EP is supported by the Research Foundation - Flanders (FWO).

more geometric proof of the main result of [CHW] that Grassmannians of planes admit such a section. In the very recent paper [GRW] (written concurrently with our paper) it is proved that, if  $X$  is a subvariety of  $\mathbb{G}_m^n$ , then a section exists on the locus in  $\text{Trop}(X)$  where the tropical multiplicity equals one [GRW]. This beautiful general theorem implies parts of our results, e.g. for linear spaces. The emphasis in our paper, however, is on explicit sections in concrete examples.

Throughout, we will assume that the valuation  $K \rightarrow \mathbb{R}_\infty$  is surjective. This is no restriction for our purposes. Indeed,  $(K, v)$  always embeds into a valued field  $(L, v_L)$  with  $v_L$  surjective. This does not change  $\text{Trop}(X)$ , and a suitable section  $\text{Trop}(X) \rightarrow X_L^{\text{an}}$  can be composed with the restriction map  $X_L^{\text{an}} \rightarrow X^{\text{an}}$  to obtain a section  $\text{Trop}(X) \rightarrow X^{\text{an}}$ .

We will use the following notation and facts. Given a point  $\xi \in \mathbb{R}_\infty^n$  we write

$$K[x]_\xi := \left\{ \sum_{\alpha} c_{\alpha} x^{\alpha} \mid v(c_{\alpha}) + \alpha \cdot \xi \geq 0 \right\}$$

for the *tilted group ring* [Pay09]. This ring contains the valuation ring of  $K$  and it has an ideal with the same definition but with  $\geq$  replaced by  $>$ . The quotient by this ideal is an algebra over the residue field  $k$  of  $K$ . By surjectivity of the valuation, this algebra is in fact a polynomial ring over  $k$  in at most  $n$  variables—generators can be obtained as the images  $y_i$  of  $c_i x_i$  where  $i$  ranges through the set where  $\xi_i \neq \infty$  and where the coefficients  $c_i \in K$  are chosen such that  $v(c_i) + \xi_i = 0$ . Let  $I(X)$  be the ideal of  $X$  in  $K[x]$ . The image of  $I(X) \cap K[x]_\xi$  in the polynomial ring  $k[y_i \mid i : \xi_i \neq \infty]$  is called the *initial ideal*  $\text{in}_{\xi} I(X)$  of  $I(X)$  relative to  $\xi$ , and the scheme  $\text{in}_{\xi} X$  defined by it is called the *initial degeneration* of  $X$ . The point  $\xi$  lies in the tropical variety if and only if  $\text{in}_{\xi} I(X)$  does not contain monomials [MS, Chapter 3].

The remainder of this paper is organised as follows. In Section 2 we prove that if  $Y \subseteq \mathbb{A}^n$  is a linear space, then the surjection  $Y^{\text{an}} \rightarrow \text{Trop}(Y)$  has a continuous section. In Section 3, given an action of an  $m$ -dimensional subtorus of  $\mathbb{G}_m^n$  on a subvariety  $X \subseteq \mathbb{A}^n$ , we construct an action of  $\mathbb{R}^m$  on a retract  $Z \subseteq X^{\text{an}}$ , which maps surjectively and  $\mathbb{R}^m$ -equivariantly onto  $\text{Trop}(X)$ . In Section 4 we introduce techniques for finding sections  $\text{Trop}(X) \rightarrow Z$  when  $X$  is obtained by smearing around a linear space  $Y$  with a torus action. As an example, we treat the variety in  $\mathbb{G}_m^{m \times p}$  of matrices of less than full rank. In Sections 5 and 6 we apply our techniques to Grassmannians of two-spaces and to matrices of rank two, respectively. We conclude with a brief discussion of  $A$ -discriminants in Section 7.

**Acknowledgments.** We thank Joe Rabinoff for interesting discussions and, more specifically, for pointing out that the retract  $Z$  of  $X^{\text{an}}$  that we define in Section 3 is, in fact, a strong deformation retract.

## 2. LINEAR SPACES

In this section we assume that  $Y$  is a linear subspace through the origin  $0 \in \mathbb{A}^n$ . Tropical linear spaces are well-understood through their circuits and cocircuits [YY07, AK06], and the proof of the following theorem is very natural from that perspective.

**Theorem 2.1.** *For any linear subspace  $Y \subseteq \mathbb{A}^n$  the projection  $\pi_Y : Y^{\text{an}} \rightarrow \text{Trop}(Y)$  has a continuous section.*

Without loss of generality, we may restrict to the case where  $Y$  is not contained in any coordinate hyperplane, so that  $\text{Trop}(Y)$  is the closure of  $\text{Trop}(Y) \cap \mathbb{R}^n$ . Nevertheless, we will need to check carefully that the section we construct is also continuous on  $\text{Trop}(Y) \setminus \mathbb{R}^n$ . We will use that for  $\eta \in \text{Trop}(Y) \cap \mathbb{R}^n$  the initial degeneration  $\text{in}_\eta Y$  is a linear subspace of  $\mathbb{A}_k^n$  of the same dimension as  $Y$ . For general  $\eta \in \text{Trop}(Y)$  we have  $\text{in}_\eta Y = \text{in}_\eta Y'$ , where  $Y'$  is the subspace of  $Y$  consisting of all  $y$  with  $x_i(y) = 0$  for all  $i$  with  $\eta_i = \infty$ .

*Proof of Theorem 2.1.* We define the section  $\sigma : \text{Trop}(Y) \rightarrow Y^{\text{an}}$  as follows. Pick  $\eta \in \text{Trop}(Y)$ , set  $S := \{i \in [n] \mid \eta_i = \infty\}$ , and let  $Y' \subseteq Y$  be the subspace of all  $y \in Y$  with  $x_i(y) = 0$  for all  $i \in S$ . Choose a maximal set  $J \subseteq [n]$  such that the  $y_j, j \in J$  (from the definition of the tilted polynomial ring) are linearly independent on the subspace  $\text{in}_\eta Y = \text{in}_\eta Y'$ . In particular,  $J$  is disjoint from  $S$ . Then the corresponding  $x_j, j \in J$  are linearly independent on  $Y'$ , and the inclusion  $K[x_j \mid j \in J] \rightarrow K[Y']$  is an isomorphism. In general, a basis  $J$  of the matroid defined by  $Y'$  will be called *compatible* with  $\eta$  (and  $\eta$  with  $J$ ) if it arises this way.

For  $f \in K[Y]$  we can uniquely write  $f|_{Y'} = \sum_\alpha c_\alpha x^\alpha$  where the  $\alpha$  run through  $\mathbb{N}^J$ , and we set

$$\sigma(\eta)(f) := \min_\alpha (v(c_\alpha) + \alpha \cdot \eta).$$

This is clearly a valuation that maps  $x_j, j \in J$  to  $\eta_j$  and that maps the  $x_i$  with  $i \in S$  to  $\infty$ . What about  $x_i$  with  $i \notin J \cup S$ ? Up to a scalar factor, there exists a unique non-zero linear relation

$$\sum_{j \in J \cup \{i\}} d_j x_j \in I(Y').$$

After scaling by a common factor we may assume that  $v(d_j) + \eta_j \geq 0$  for all  $j \in J \cup \{i\}$  and that equality holds for at least one  $i$ . Then this element lies in  $I(Y') \cap K[x_i \mid i \notin S]_\eta$ . If  $v(d_i) + \eta_i$  were strictly positive, then projecting down into  $k[y_i \mid i \notin S]$  would yield a relation among the  $y_j$  with  $j \in J$ , a contradiction to the choice of  $J$ . Hence  $v(d_i) + \eta_i = 0$ . If  $v(d_j) + \eta_j$  were strictly positive for all  $j \in J$ , then projecting down would yield  $y_i \in \text{in}_\eta I(Y')$ , which contradicts  $\eta \in \text{Trop}(Y')$ . Hence  $v(d_j) + \eta_j = 0$  for some  $j \in J$ . This shows that

$$\sigma(\eta)(x_i) = \min_{j \in J} (v(-d_j/d_i) + \eta_j) = \eta_i,$$

as required. So  $\sigma(\eta) \in Y^{\text{an}}$  is a point in the fibre of  $\pi_Y$  above  $\eta$ .

To define  $\sigma(\eta)$ , we have made the choice of a basis  $J$  in the matroid defined by  $\text{in}_\eta I(Y')$ . But in fact, this choice does not influence the outcome. Indeed, *any* valuation  $w \in Y^{\text{an}}$  with  $\pi_Y(w) = \eta$  must satisfy  $w(f) \geq \sigma(\eta)(f)$  for all  $f \in K[Y]^1$ . In particular, this must hold for all valuations constructed from other bases of the matroid. This shows that  $\sigma$  is well-defined on all of  $\text{Trop}(Y)$ .

It remains to show that  $\sigma$  is continuous. This is immediate from the formula for  $\sigma(\eta)$  on a subset of  $\text{Trop}(Y)$  where  $S$  and  $J$  compatible with  $\eta$  are fixed. Let  $Y'$  be as above. Suppose that a sequence  $\eta^{(l)}, l = 1, 2, \dots$  in this set converges to a point  $\eta \in \text{Trop}(Y)$ . Note that the set of  $i$  with  $\eta_i = \infty$  contains  $S$  but may be strictly larger, and may even contain elements of  $J$ . Even so, for every non-zero relation

<sup>1</sup>So  $\sigma(\eta)$  is the *Shilov boundary point* [Ber90, Chapter 2] in the fibre in  $Y^{\text{an}}$  above  $\eta$ .

$\sum_{j \in J \cup \{i\}} d_j x_j \in I(Y')$  for  $i \notin J \cup S$  we have  $\min_{j \in J} (v(d_j) + \eta_j^{(l)}) = v(d_i) + \eta_i^{(l)}$ . This closed condition then also holds in the limit:

$$(1) \quad \min_{j \in J} (v(d_j) + \eta_j) = v(d_i) + \eta_i.$$

Let  $w$  be the valuation of  $K[Y]$  defined by mapping  $f \in K[Y]$  with  $f|_{Y'} = \sum_{\alpha \in \mathbb{N}^J} c_\alpha x^\alpha$  to  $\min_{\alpha \in \mathbb{N}^J} (v(c_\alpha) + \alpha \cdot \eta)$ . Then  $w$  is, indeed, a valuation of  $K[Y]$ , which maps  $x_j$  to  $\eta_j$  for  $j \in J$  (because  $x_j|_{Y'}$  is a single term) and for  $j \in S$  (because  $x_j|_{Y'}$  has no terms) and for  $j \notin J \cup S$  (by (1)). Moreover,  $w(f)$  is minimal among all such valuations, so  $w(f) = \sigma(\eta)(f)$ . This shows that  $\sigma$  is continuous on the closure of the set of all  $\eta$  compatible with a given  $S$  and  $J$ . These closures form a finite closed cover of  $\text{Trop}(Y)$ , hence  $\sigma$  is continuous everywhere.  $\square$

**Remark 2.2.** In the *constant coefficient case*, where  $Y$  is a linear space defined over a subfield of  $K$  on which the valuation is trivial, the choice of  $J$  above can be made more constructive, as follows. Given  $\eta \in \text{Trop}(Y) \cap \mathbb{R}^n$ , take a permutation  $\pi \in S_n$  such that  $\eta_{\pi(1)} \geq \dots \geq \eta_{\pi(n)}$ . Then construct  $J$  by setting  $J_0 := \emptyset$  and

$$J_i := \begin{cases} J_{i-1} \cup \{i\} & \text{if } x_i|_Y \text{ linearly independent of } \langle x_j|_Y \mid j \in J_{i-1} \rangle, \text{ and} \\ J_{i-1} & \text{otherwise.} \end{cases}$$

Then  $J := J_n$  is a basis of the matroid defined by  $Y$  compatible with  $\eta$ . This is the greedy algorithm for finding a maximal-weight basis in a matroid [Sch03, Chapter 40].

Conversely, given a basis  $J$  of that matroid, we can construct all  $\eta \in \text{Trop}(Y) \cap \mathbb{R}^n$  compatible with it by choosing the  $\eta_j$  with  $j \in J$  arbitrarily and setting  $\eta_i$  for  $i \notin J$  equal to the minimal value  $\eta_j$  for  $j$  in the unique circuit contained in  $J \cup \{i\}$ . We will use this explicit construction in Sections 5 and 6. These remarks apply, *mutatis mutandis*, also to  $\eta \in \text{Trop}(Y) \setminus \mathbb{R}^n$ .

### 3. TORUS ACTIONS

Let  $\varphi : \mathbb{G}_m^m \rightarrow \mathbb{G}_m^n$  be a homomorphism of tori. This is of the form  $\varphi(t_1, \dots, t_m) = (t^{a_1}, \dots, t^{a_n})$  where  $a_1, \dots, a_n \in \mathbb{Z}^m$ . Let  $A \in \mathbb{Z}^{n \times m}$  be the matrix with rows  $a_1, \dots, a_n$ . Let  $X \subseteq \mathbb{A}^n$  or  $X \subseteq \mathbb{G}_m^n$  be a closed affine subvariety stable under the  $\mathbb{G}_m^m$ -action on  $\mathbb{A}^n$  (or  $\mathbb{G}_m^n$ ) given by  $\varphi$ . Then  $\mathbb{R}^m$  has a continuous action on  $\text{Trop}(X)$  given by

$$(\tau, \xi) \mapsto A\tau + \xi.$$

The column space of  $A$  is contained in the *lineality space* of  $\text{Trop}(X)$ . In this section we investigate to what extent this action can be lifted to  $X^{\text{an}}$ . For this, we denote by  $\lambda : \mathbb{G}_m^m \times X \rightarrow X, (t, x) \mapsto \varphi(t)x$  the action of  $\mathbb{G}_m^m$  on  $X$  and by  $\lambda^* : K[X] \rightarrow K[\mathbb{G}_m^m \times X]$  its comorphism.

**Lemma 3.1.** *There exists a commutative diagram of continuous maps:*

$$\begin{array}{ccccc} \mathbb{R}^m \times X^{\text{an}} & \longrightarrow & (\mathbb{G}_m^m \times X)^{\text{an}} & \xrightarrow{w \mapsto w \circ \lambda^*} & X^{\text{an}} \\ \downarrow \text{id} \times \pi & & & & \downarrow \pi \\ \mathbb{R}^m \times \text{Trop}(X) & \xrightarrow{(\tau, \xi) \mapsto A\tau + \xi} & & & \text{Trop}(X). \end{array}$$

*Proof.* The right-most map in the top row of the diagram is the analytification of the torus action. The only map that needs a definition is the left-most map in that row. It sends  $(\tau, w)$  to the valuation of  $K[\mathbb{G}_m^m \times X]$  defined by

$$\sum_{\beta \in \mathbb{Z}^m} f_\beta t^\beta \mapsto \min_{\beta} (w(f_\beta) + \beta \cdot \tau).$$

To see that the diagram commutes, pick  $(\tau, w) \in \mathbb{R}^m \times X^{\text{an}}$  and let  $w' \in X^{\text{an}}$  be the image of that pair along the top row. We have  $\lambda^* x_i = t^{a_i} x_i$ , and hence

$$w'(x_i) = w(x_i) + a_i \cdot \tau.$$

This implies that  $\pi(w') = \pi(w) + A\tau$ , as claimed.  $\square$

Let  $\mu$  denote the composition of the two maps in the first row. Unwinding the definitions, we find that  $\mu$  sends  $(\tau, w)$  to a valuation on  $K[X]$  defined as follows. Pick  $f \in K[X]$  and decompose  $f = \sum_{\beta \in \mathbb{Z}^m} f_\beta$  into  $\mathbb{G}_m^m$ -weight vectors, i.e., with  $\lambda^* f_\beta = t^\beta f_\beta$ . Then

$$\mu(\tau, w)(f) = \min_{\beta} (w(f_\beta) + \beta \cdot \tau).$$

We remark that if  $A\tau = 0$ , then  $\mu(\tau, w) = \mu(0, w)$ . Indeed, then  $\tau$  is perpendicular to the rows of  $A$ , hence to any  $\mathbb{Z}$ -linear combination of these, and the  $\beta$  for which there exist non-zero  $f_\beta \in K[X]$  of weight  $\beta$  are such linear combinations.

In general,  $\mu$  is not an action of  $\mathbb{R}^m$  on  $X^{\text{an}}$ . Indeed,  $\mu(0, w)$  may not equal  $w$ . For instance, set  $X = \mathbb{A}^2$  with coordinate ring  $K[x_1, x_2]$ , and let  $m = 2$  and let  $\varphi$  be the identity. Let  $w$  be any element of  $X^{\text{an}}$  and set  $\xi := \pi(w)$ . Then the image of  $(0, w)$  along the first row equals the valuation  $w'$  of  $K[x_1, x_2]$  defined by

$$\sum_{i,j} c_{ij} x_1^i x_2^j \mapsto \min_{i,j} (v(c_{ij}) + i\xi_1 + j\xi_2).$$

Of course, it may very well be that  $w \neq w'$ . However, the following lemma shows that  $\mu(0, w) \neq w$  is the *only* obstacle to  $\mu$  being an action.

**Lemma 3.2.** *Define  $Z$  as the image of  $\mu$ . Then  $Z$  is a closed subset of  $X^{\text{an}}$  and the restriction of  $\mu$  to  $\mathbb{R}^m \times Z$  defines a continuous action of  $\mathbb{R}^m$  on  $Z$ . Moreover, the map  $w \mapsto \mu(0, w)$  defines a continuous retraction from  $X^{\text{an}}$  to  $Z$ .*

*Proof.* First, for  $\tau_1, \tau_2 \in \mathbb{R}^m$  and  $f \in K[X]$  with  $\mathbb{G}_m^m$ -weight decomposition  $f = \sum_{\beta \in \mathbb{Z}^m} f_\beta \in K[X]$  and  $w \in X^{\text{an}}$  we compute

$$\begin{aligned} \mu(\tau_1, \mu(\tau_2, w))(f) &= \min_{\beta \in \mathbb{Z}^m} (\beta \cdot \tau_1 + \mu(\tau_2, w)(f_\beta)) \\ &= \min_{\beta \in \mathbb{Z}^m} (\beta \cdot \tau_1 + \beta \cdot \tau_2 + w(f_\beta)) = \mu(\tau_1 + \tau_2, w)(f). \end{aligned}$$

This implies that  $\mu(0, \mu(\tau, w)) = \mu(\tau, w)$ , so that 0 acts as the identity on  $Z$ . Hence  $\mu$  is an action on  $Z$ . Furthermore,  $Z$  can be characterised as the set of all  $w \in X^{\text{an}}$  with  $\mu(0, w) = w$ . As  $\mu$  is continuous, this implies that  $Z$  is closed. The last statement is immediate.  $\square$

The following refinement of the statement that  $Z$  is a retract of  $X^{\text{an}}$  was pointed out to us by Joe Rabinoff.

**Proposition 3.3.** *In the setting above,  $Z$  is a strong deformation retract of  $X^{\text{an}}$ .*

*Proof.* This can be derived using the general techniques of [Ber90, Chapter 6]; here is a shortcut in our language. For  $r \in [0, \infty]$  and  $w \in X^{\text{an}}$  let  $w_r$  be the function  $K[X] \rightarrow \mathbb{R}_\infty$  defined as follows. Take  $f \in K[X]$ , expand  $f(\varphi(t)x) := \sum_{\beta \in \mathbb{Z}^m} f_\beta t^\beta$ , and rewrite this Laurent series with  $K[X]$ -coefficients as a formal power series

$$\sum_{\beta \in \mathbb{Z}^m} f_\beta t^\beta = \sum_{\gamma \in (\mathbb{Z}_{\geq 0})^m} g_\gamma (t-1)^\gamma$$

around the identity element  $1 = (1, \dots, 1)$  of  $\mathbb{G}_m^m$ . Set

$$w_r(f) := \min_{\gamma} (w(g_\gamma) + |\gamma|r), \text{ where } |\gamma| := \gamma_1 + \dots + \gamma_m.$$

We argue that this minimum is attained, and that it can be replaced by a minimum over a finite set of  $\gamma$ s that does not depend on  $w$  or  $r$ . In the rewriting process, we replace each Laurent monomial  $t^\beta$  by the formal power series of  $((t-1)+1)^\beta$  around 1. This shows that each  $g_\gamma$  is a  $\mathbb{Z}$ -linear combination of the  $f_\beta$ . In particular, for all  $\gamma$  we have  $w(g_\gamma) \geq \min_{\beta} w(f_\beta)$ , and for  $r > 0$  this suffices to conclude that the minimum is attained.

Conversely, we claim that each  $f_\beta$  is a  $\mathbb{Z}$ -linear combination of the  $g_\gamma$ . This is immediate if all  $\beta$  with  $f_\beta \neq 0$  are already in  $(\mathbb{Z}_{>0})^m$  (since then we are just rewriting polynomials, and the rewriting can be reversed). The general case can be reduced to this, since multiplication of power series with a fixed power series of the form  $((t-1)+1)^\beta$  is a  $\mathbb{Z}$ -linear isomorphism with inverse equal to multiplication with  $((t-1)+1)^{-\beta}$ . Consequently, we find that the minimum is attained for  $r = 0$ , as well, and that  $\min_{\gamma} w(g_\gamma) = \min_{\beta} w(f_\beta) = \mu(0, w)(f)$ .

Combining the two  $\mathbb{Z}$ -linear transitions, all countably many  $g_\gamma$  are  $\mathbb{Z}$ -linear combinations of finitely many among them. If  $d$  is the maximum value of  $|\gamma|$  among these finitely many, then we can replace the minimum defining  $w_r(f)$  by the minimum over all  $\gamma$  with  $|\gamma| \leq d$ . Then it is evident that  $w_r(f)$  depends continuously on the pair  $(w, r) \in X^{\text{an}} \times [0, \infty]$ .

Now  $w_r$  is a point in  $X^{\text{an}}$  that depends continuously on  $(w, r)$ . For  $r = \infty$  we have

$$w_\infty(f) = w(g_0) = w\left(\sum_{\beta} f_\beta\right) = w(f),$$

so  $w_\infty = w$ . As mentioned above, we have  $w_0 = \mu(0, w)$ . Finally, we must argue that if  $w$  already lies in  $Z$ , that is, if  $w = \mu(0, w)$ , then  $w_r = w$  for all  $r \in (0, \infty]$ . But in this case the  $\gamma = 0$  term in the definition of  $w_r$  equals  $\min_{\beta} w(f_\beta)$  and all other terms are (strictly) larger than this, so that  $w_r = w$  as desired.  $\square$

We conclude this section with two remarks on quotients. The first concerns the categorical quotient  $X//\mathbb{G}_m^m$  of  $X$  by the action of  $\mathbb{G}_m^m$ , i.e., the affine variety with coordinate ring equal to the ring of  $\mathbb{G}_m^m$ -invariants in  $K[X]$ . The morphism  $X \rightarrow X//\mathbb{G}_m^m$  gives rise to a morphism of analytic spaces, which sends a valuation  $w \in X^{\text{an}}$  to its restriction to the  $\mathbb{G}_m^m$ -invariants.

**Lemma 3.4.** *The map  $X^{\text{an}} \rightarrow (X//\mathbb{G}_m^m)^{\text{an}}$  factorises as*

$$X^{\text{an}} \rightarrow Z \rightarrow Z/\mathbb{R}^m \rightarrow (X//\mathbb{G}_m^m)^{\text{an}}.$$

*Proof.* We need to show that, for  $\tau \in \mathbb{R}^m$  and  $w \in X^{\text{an}}$ , the restriction of  $w' := \mu(\tau, w)$  to the  $\mathbb{G}_m^m$ -invariants  $f \in K[X]$  does not depend on  $\tau$  and equals the

restriction of  $w$  to  $\mathbb{G}_m^m$ -invariants. But this is immediate:  $f$  has weight zero, and hence

$$w'(f) = \mu(\tau, w)(f) = w(f) + 0 \cdot \tau = w(f),$$

as desired.  $\square$

The second remark concerns the passage from affine cones to projective varieties. Suppose that  $X \subseteq \mathbb{A}^n$  is an affine cone, and denote by  $\mathbb{P}X \subseteq \mathbb{P}^{n-1}$  the corresponding projective variety. The points of  $(\mathbb{P}X)^{\text{an}}$  are equivalence classes of points of  $X^{\text{an}} \setminus \{\infty\}$ .

**Lemma 3.5.** *The map  $Z \rightarrow (\mathbb{P}X)^{\text{an}}$  factorises as*

$$Z \rightarrow Z/U \rightarrow (\mathbb{P}X)^{\text{an}},$$

where  $U := A^{-1}\mathbb{R}(1, \dots, 1)$ .

*Proof.* We need to show that if  $A\tau = (C, \dots, C)$  for some  $C \in \mathbb{R}$  and if  $w \in Z$ , then  $w' := \mu(\tau, w)$  is equivalent to  $w$ . Thus let  $f$  be a homogeneous polynomial of degree  $d$  in the graded ring  $K[X]$ , and decompose  $f = \sum_{\beta \in \mathbb{Z}^m} f_\beta$ . Then  $\beta \cdot \tau = dC$  for all  $\beta$  with  $f_\beta$  non-zero, and hence

$$w'(f) = \min_{\beta} (w(f_\beta) + \beta \cdot \tau) = dC + \min_{\beta} w(f_\beta) = \mu(0, w) = w.$$

$\square$

#### 4. SMEARING A SUBSPACE AROUND BY A TORUS

Let  $Y \subseteq \mathbb{A}^n$  be a linear subspace not contained in any coordinate hyperplane and let  $\varphi : \mathbb{G}_m^m \rightarrow \mathbb{G}_m^n$  be a torus homomorphism given by an  $n \times m$  integer matrix  $A$ . Define

$$X := \overline{\{\varphi(t)y \mid y \in Y, t \in \mathbb{G}_m^m\}},$$

so that  $X$  is stable under the action of  $\mathbb{G}_m^m$ . Let  $X^0, Y^0$  be the open subsets of  $X, Y$ , respectively, where none of the coordinates vanish. Then we have  $\text{Trop}(X^0) = \text{Trop}(X) \cap \mathbb{R}^n$  and  $\text{Trop}(Y^0) = \text{Trop}(Y) \cap \mathbb{R}^n$  and

$$\text{Trop}(X^0) = A\mathbb{R}^m + \text{Trop}(Y^0);$$

this follows, for instance, from [Pay09, Proposition 2.5]. Let  $\mu : \mathbb{R}^m \times X^{\text{an}} \rightarrow Z$  be the map constructed in Section 3. We then obtain a continuous map

$$\mathbb{R}^m \times \text{Trop}(Y) \rightarrow Z, (\tau, \eta) \mapsto \mu(\tau, \sigma_Y(\eta)),$$

where  $\sigma_Y$  is the section of  $Y^{\text{an}} \rightarrow \text{Trop}(Y)$  constructed in Section 2. We would like to use this map to construct a section  $\text{Trop}(X) \rightarrow Z$  of the surjection  $Z \rightarrow \text{Trop}(X)$ , or at least a section  $\text{Trop}(X^0) \rightarrow Z^0$ , where  $Z^0$  is the preimage of  $X^0$  in  $Z^0$ . There are two basic strategies for doing so. The first strategy is given in the following proposition.

**Proposition 4.1.** *If the map  $\mathbb{R}^m \times \text{Trop}(Y^0) \rightarrow \text{Trop}(X^0)$ ,  $(\tau, \eta) \mapsto A\tau + \eta$  has a continuous section, then so does the map  $\pi : Z^0 \rightarrow \text{Trop}(X^0)$ . Moreover, if the former section can be chosen  $\mathbb{R}^m$ -equivariant, then so can the latter.*

Here the action of  $\mathbb{R}^m$  on  $\mathbb{R}^m \times \text{Trop}(Y^0)$  is given by addition in the first coordinate and the trivial action on  $\text{Trop}(Y^0)$ .



*Proof.* The composition

$$\sigma : (\text{Trop}(X^0) \rightarrow \mathbb{R}^m \times \text{Trop}(Y^0) \rightarrow Z^0)$$

of a continuous section  $\text{Trop}(X^0) \rightarrow \mathbb{R}^m \times \text{Trop}(Y^0)$  and the map  $(\tau, \eta) \mapsto \mu(\tau, \sigma_Y(\eta))$  is a section  $\text{Trop}(X^0) \rightarrow Z^0$ . The second statement is immediate.  $\square$

Here is an application of this construction.

**Theorem 4.2.** *Let  $m \leq p$  natural numbers. Let  $X \subseteq \mathbb{A}^{m \times p}$  be the matrix variety defined by the vanishing of all  $m \times m$ -minors. Then the map  $\pi : (X^0)^{\text{an}} \rightarrow \text{Trop}(X^0)$  has a continuous section.*

*Proof.* Let  $Y$  be the linear subspace contained in  $X$  consisting of all matrices  $y$  such that  $\mathbf{1}^T y = 0$ , and let  $\mathbb{G}_m^m$  act on matrices by scaling the rows. Let  $A$  be the corresponding  $(mp) \times m$ -matrix of integers. It has a 1 at position  $((i, j), k)$  if  $k = i$  and a 0 otherwise. Then  $X = \overline{\mathbb{G}_m^m Y}$  and hence  $\text{Trop}(X^0) = A\mathbb{R}^m + \text{Trop}(Y^0)$ . Note that  $\text{Trop}(Y^0)$  is the set of matrices  $\eta$  such that the  $p$  tropical hyperplanes in  $\mathbb{R}^m$  defined by the columns of  $\eta$  all contain  $\mathbb{R}(1, \dots, 1)$ .

We claim that the map

$$\mathbb{R}^m \times \text{Trop}(Y^0) \rightarrow \text{Trop}(X^0), (\tau, \eta) \mapsto A\tau + \eta$$

has a continuous section, defined as follows. Let  $\xi \in \text{Trop}(X^0)$ . Then for  $\tau \in \mathbb{R}^m$  the condition that  $\xi - A\tau$  lies in  $\text{Trop}(Y^0)$  is equivalent to the condition that for each  $j = 1, \dots, p$  the minimum  $\min_i(\xi_{ij} - \tau_i)$  is attained at least twice. This means that  $-\tau$  lies on the intersection of the  $p$  tropical hyperplanes in  $\mathbb{R}^m$  with coefficient vectors given by the columns of  $\xi$ . Take  $-\tau$  in the *stable intersection* (see [RGST05, Section 5] and [Mik06, Section 4]) of the first  $m - 1$  of these hyperplanes; this is a set of the form  $(c_1, \dots, c_m) + \mathbb{R}(1, \dots, 1)$ . For definiteness chose  $\tau$  such that  $\tau_1 = 0$ . By basic properties of stable intersections, this  $\tau$  depends continuously on  $\xi$ . We claim that it has the property that  $\xi - A\tau$  lies in  $\text{Trop}(Y^0)$ , i.e., that  $-\tau$  also lies in the tropical hyperplanes defined by the remaining columns of  $\xi$ . This is true on the subset of  $\text{Trop}(X^0)$  where the *set-theoretic* intersection of the hyperplanes given by the first  $m - 1$  columns is of the required form  $(c_1, \dots, c_m) + \mathbb{R}(1, \dots, 1)$ , because on this set there is no other choice for  $\tau$  up to tropical scalar multiplication (and, since  $\xi \in \text{Trop}(X^0)$ ,  $\xi - A\tau$  lies in  $\text{Trop}(Y^0)$  for *some*  $\tau$ ). But the set of such  $\xi$  is dense in  $\text{Trop}(X^0)$ , because it is the image of  $\mathbb{R}^m \times U$ , where  $U$  is the dense subset of  $\text{Trop}(Y^0)$  where the intersection of the hyperplanes given by the first  $m - 1$  columns of  $\eta$  equals exactly  $\mathbb{R}(1, \dots, 1)$ .

Thus we may apply Proposition 4.1 and find a continuous section  $\text{Trop}(X^0) \rightarrow Z^0 \subseteq (X^0)^{\text{an}}$ . As constructed, this section is not quite  $\mathbb{R}^m$ -equivariant, as we chose  $\tau_1 = 0$ . If, instead, we choose  $\tau_1$  equal to  $\xi_{11}$ , then we do obtain an  $\mathbb{R}^m$ -equivariant continuous section.  $\square$

**Remark 4.3.** We do not know whether the section constructed in this proof extends to all of  $\text{Trop}(X)$ , or if not, whether it can be adapted to extend.

The second strategy for constructing a section  $\text{Trop}(X^0) \rightarrow (X^0)^{\text{an}}$  is to show that the map  $\mathbb{R}^m \times \text{Trop}(Y^0) \rightarrow (X^0)^{\text{an}}$  factors through the map  $\mathbb{R}^m \times \text{Trop}(Y^0) \rightarrow \text{Trop}(X^0)$ . We will now formulate sufficient conditions for this to happen.

The first of these conditions is purely polyhedral, namely, we require that for each  $\eta \in \text{Trop}(Y^0)$  the set

$$T_\eta := \{\tau \in \mathbb{R}^m \mid A\tau + \eta \in \text{Trop}(Y^0)\},$$

which is the support of a polyhedral complex, is connected. Observe that these sets encode the ambiguity in the decomposition of  $\xi$ : if  $\xi$  equals both  $\eta_1 + A\tau_1$  and  $\eta_2 + A\tau_2$ , then  $\tau_1 - \tau_2 \in T_{\eta_1}$ . The second condition is more algebraic. Let  $\eta \in \text{Trop}(Y^0)$ . Extend the valuation  $w := \sigma_Y(\eta)$  from  $K[Y]$  to the field  $K(Y)$ ; this can be done since it sends no non-zero polynomials to infinity. Let  $y \in Y(K(Y))$  be the generic point of  $Y$ . By slight abuse of notation, we have  $w(y) = \eta$ . Let  $\tau \in \mathbb{R}^m$  be such that the line segment  $[0, \tau]$  is contained in  $T_\eta$ . Then we require that for all sufficiently small  $\epsilon > 0$  there exists a valued extension  $(L, w_L)$  of  $K(Y)$  and a  $t \in \mathbb{G}_m^m(L)$  such that  $w_L(t) = \epsilon\tau$  and  $\varphi(t)y \in Y^0(L)$ .

**Proposition 4.4.** *Suppose that the torus homomorphism  $\varphi : \mathbb{G}_m^m \rightarrow \mathbb{G}_m^n$  and the linear space  $Y$  satisfy the two aforementioned requirements. Then, for  $\xi = A\tau_1 + \eta_1 \in \text{Trop}(X^0)$  with  $\tau_1 \in \mathbb{R}^m$  and  $\eta_1 \in \text{Trop}(Y^0)$  the expression*

$$\sigma(\xi) := \mu(\tau_1, \sigma_Y(\eta_1)) \in Z \subseteq X^{\text{an}}$$

*does not depend on the chosen decomposition of  $\xi \in \text{Trop}(X^0)$ . The map  $\sigma : \text{Trop}(X^0) \rightarrow Z^0$  thus defined is a continuous,  $\mathbb{R}^m$ -equivariant section of the surjection  $Z^0 \rightarrow \text{Trop}(X^0)$ .*

Before we give the proof, we discuss a simple example in the plane.

**Example 4.5.** Let  $Y \subseteq \mathbb{A}^2$  be given by the linear equation  $x_1 - x_2 = 0$ , and let  $\varphi : \mathbb{G}_m \rightarrow \mathbb{G}_m^2$  be given by  $\varphi(t) = (t, t^{-1})$ , so that  $A = (1, -1)^T$ . Then we have

$$\text{Trop}(Y) = \{(\eta_1, \eta_2) \in \mathbb{R}_\infty^2 \mid \eta_1 = \eta_2\} \text{ and } A\mathbb{R}^1 = \{(\tau, -\tau) \mid \tau \in \mathbb{R}\}.$$

We have  $X = \overline{\varphi(\mathbb{G}_m)Y} = \mathbb{A}^2$  and  $\text{Trop}(X^0) = A\mathbb{R}^1 + \text{Trop}(Y^0)$  and  $T_\eta = \{0\}$  for all  $\eta \in \text{Trop}(Y^0)$ . In the second requirement we can just take  $t = 1$  for all  $\eta$ . Thus both requirements are met. For  $\xi = (\xi_1, \xi_2) = A\tau + \eta = (\tau + \eta_1, -\tau + \eta_1)$  and  $f = \sum_{i,j} c_{ij}x_1^i x_2^j$  we find

$$\sigma(\xi)(f) = \min_{k \in \mathbb{Z}} \min_{i-j=k} (k\tau + v(c_{ij}) + (i+j)\eta_1) = \min_{i,j} (v(c_{ij}) + i\xi_1 + j\xi_2),$$

which extends to all of  $\text{Trop}(X)$ , and in fact equals the section obtained in Section 2 when regarding  $X$  as a linear space.  $\diamond$

*Proof of Proposition 4.4.* For the first statement we need to prove that if  $\xi$  can also be decomposed as  $A\tau_2 + \eta_2$  then

$$\mu(\tau_2, \sigma_Y(\eta_2)) = \mu(\tau_1, \sigma_Y(\eta_1)).$$

This is equivalent to

$$\mu(\tau_2 - \tau_1, \sigma_Y(\eta_2)) = \mu(0, \sigma_Y(\eta_1)).$$

Now  $\tau_1 - \tau_2 \in T_{\eta_1}$ , and since  $T_{\eta_1}$  is connected it suffices to prove the following local version of this equality. Let  $\eta \in \text{Trop}(Y^0)$  and  $\tau \in \mathbb{R}^m$  be such that the segment  $[0, \tau]$  lies entirely in  $T_\eta$ . Then we want to show that

$$\mu(-\tau, \sigma_Y(A\tau + \eta)) = \mu(0, \sigma_Y(\eta)).$$

By definition of  $\mu$ , it suffices to prove this when applied to a non-zero  $f \in K[X]$  that is homogeneous with respect to the  $\mathbb{G}_m^m$ -action, say of weight  $\beta$ . We will prove, in fact, that the function

$$\ell : [0, 1] \rightarrow \mathbb{R}, \quad \epsilon \mapsto \mu(-\epsilon\tau, \sigma_Y(A\epsilon\tau + \eta))(f)$$

is constant on the interval  $[0, 1]$ . Since  $\ell$  is a continuous function and  $[0, 1]$  is connected, it suffices to prove that  $\ell$  has a local minimum at every point in  $[0, 1]$ . We give the argument at the point 0; it follows at other points in a similar manner.

Set  $w := \sigma_Y(\eta)$ , and let  $y \in Y(K(Y))$  be the generic point. Then for  $\epsilon > 0$  sufficiently small a valued field extension  $(L, w_L) \supseteq (K(Y), w)$  and  $t \in \mathbb{G}_m^m(L)$  exist as in the second requirement, that is, with  $w_L(t) = \epsilon\tau$  and  $\varphi(t)y \in Y^0(L)$ . After shrinking  $\epsilon$  if necessary we may assume that  $\eta$  and  $\epsilon A\tau + \eta$  are both compatible with the same basis  $J \subseteq [n]$  of the matroid defined by  $Y$ . Expand the restriction  $f|_Y$  as  $\sum_{\alpha \in \mathbb{N}^J} c_\alpha x^\alpha$ . Then on the one hand we have

$$w_L(f|_Y(\varphi(t)y)) = w_L(t^\beta f|_Y(y)) = \beta \cdot \epsilon\tau + \sigma_Y(\eta)(f),$$

where we have used that  $\varphi(t)y \in Y(L)$  and that  $f$  is homogeneous of  $\mathbb{G}_m^m$ -weight  $\beta$ . On the other hand, we have

$$w_L(f|_Y(ty)) = w_L\left(\sum_{\alpha \in \mathbb{N}^J} c_\alpha t^{\alpha A} y^\alpha\right) \leq \min_{\alpha} (v(c_\alpha) + \alpha \cdot A\epsilon\tau + \alpha \cdot \eta) = \sigma_Y(A\epsilon\tau + \eta)(f).$$

Thus we find that

$$\ell(\epsilon) = \sigma_Y(A\epsilon\tau + \eta)(f) - \beta \cdot \epsilon\tau \geq \sigma_Y(\eta)(f) = \ell(0),$$

as desired. This shows that the section  $\sigma : \text{Trop}(X^0) \rightarrow Z$  is well-defined. To see that  $\sigma$  is continuous, decompose  $\text{Trop}(Y^0)$  into finitely many closed polyhedra  $P_i$  and let  $P'_i$  denote the image of

$$\mathbb{R}^m \times P_i \rightarrow \text{Trop}(X^0), (\tau, \eta) \mapsto A\tau + \eta.$$

By basic linear algebra over  $\mathbb{R}$ , on each  $P'_i$  this map has a continuous (in fact, affine-linear) section  $P'_i \rightarrow \mathbb{R}^m \times P_i$ . This shows that the restriction of  $\sigma$  to each  $P'_i$  is continuous. Since the  $P'_i$  form a finite closed cover of  $\text{Trop}(X^0)$ , the map  $\sigma$  is continuous on  $\text{Trop}(X^0)$ .

Finally, we need to verify that  $\sigma$  is  $\mathbb{R}^m$ -equivariant. Let  $\xi = A\tau + \eta \in \text{Trop}(X^0)$  with  $\tau \in \mathbb{R}^m$  and  $\eta \in \text{Trop}(Y^0)$ . Let  $\tau' \in \mathbb{R}^m$ . Then we have

$$\sigma(A\tau' + \xi) = \sigma(A(\tau' + \tau) + \eta) = \mu(\tau' + \tau, \sigma_Y(\eta)) = \mu(\tau', \mu(\tau, \sigma_Y(\eta))) = \mu(\tau', \sigma(\xi)),$$

as desired.  $\square$

**Remark 4.6.** While Proposition 4.4 gives a section only over  $\text{Trop}(X^0)$ , we will see that, at least in the case of Grassmannians of planes, this section extends to all of  $\text{Trop}(X)$ .


## 5. GRASSMANNIANS OF PLANES

In this section we set  $n := \binom{m}{2}$  and consider  $\mathbb{A}^n$  with coordinates  $x_{ij}$  for  $1 \leq i < j \leq n$ . We also write  $x_{ji} = -x_{ij}$  for  $i > j$ , and  $\xi_{ji} = \xi_{ij}$  for tropical coordinates. Let  $X := \widehat{\text{Gr}}(2, m) \subseteq \mathbb{A}^n$  denote the affine cone over the Grassmannian of planes, given as the image of the polynomial map

$$\psi : (\mathbb{A}^m)^2 \rightarrow \mathbb{A}^n, (y, z) \mapsto (y_i z_j - y_j z_i)_{i < j}.$$

Let  $Y$  denote the subvariety of  $X$  given as the image

$$Y := \psi(\mathbb{A}^m \times \{\mathbf{1}\}),$$


 FIGURE 1. A spanning tree in  $K_9$  with minimal-weight edge  $i_0 j_0$ 

where  $\mathbf{1}$  is the all-one vector. Then  $Y$  is a linear space with generic point  $(y_i - y_j)_{i < j}$ . Furthermore, let  $\varphi : \mathbb{G}_m^m \rightarrow \mathbb{G}_m^n$  be given by

$$\varphi(t) := (t_i t_j)_{i < j},$$

and let  $A$  be the corresponding  $n \times m$ -matrix of nonnegative integers. Then  $X$  is  $\mathbb{G}_m^m$ -stable, and in fact  $X = \overline{\varphi(\mathbb{G}_m^m) \cdot Y}$ . Hence we are in the situation of Section 4. Let  $\mu : \mathbb{R}^m \times X^{\text{an}} \rightarrow Z \subseteq X^{\text{an}}$  be the map from Section 3, which restricts to an action of  $\mathbb{R}^m$  on  $Z$ . We will prove the following theorem.

**Theorem 5.1.** *The surjective projection from  $Z \subseteq \widehat{\text{Gr}}(2, m)^{\text{an}}$  to  $\text{Trop}(\widehat{\text{Gr}}(2, m))$  has a continuous,  $\mathbb{R}^m$ -equivariant section.*

Our proof consists of two parts. We first construct a continuous section in the spirit of Proposition 4.1, which relies on the choice of a hyperplane in  $\text{Trop}(\mathbb{P}^{m-1})$ . Then we use the technique of Proposition 4.4 to verify that the constructed section is, in fact, natural and independent of the choice of hyperplane. This then also implies  $\mathbb{R}^m$ -equivariance.

We will use that the matroid on the variables  $x_{ij}$  defined by  $Y$  is the graphical matroid of the complete graph  $K_m$ . This is immediate from the definition of  $Y$ , and was also exploited in [AK06, Section 4]. Thus a basis  $J$  as in Section 2 is a tree with vertex set  $[m]$ . We will write  $\Gamma$  instead of  $J$ . Given such a tree  $\Gamma$ , one finds all  $\eta \in \text{Trop}(Y)$  compatible with  $\Gamma$  as follows. First, give arbitrary values in  $\mathbb{R}_\infty$  to all  $\eta_{ij}$  with  $ij$  an edge in the tree  $\Gamma$ . Then, for each edge  $ij$  in  $K_m \setminus \Gamma$  set  $\eta_{ij}$  equal to the minimum of the  $\eta_{kl}$  over all edges  $kl$  in the simple path from  $i$  to  $j$  in  $\Gamma$ . See Figure 1.

*Proof of Theorem 5.1, construction of a continuous section.* Up to tropical scaling, the non-infinity points of  $\text{Trop}(X)$  are in one-to-one correspondence with tropical projective lines in the simplex  $\Delta := \text{Trop}(\mathbb{P}^{m-1})$  (see [SS04, Theorem 3.8] for  $\text{Trop}(X^0)$ ). The non-infinity points of  $\text{Trop}(Y)$  correspond bijectively to the tropical lines that pass through the all-zero point 0. The action of  $\mathbb{R}^m$  is by translation of the lines.

Thus to go from  $\xi \in \text{Trop}(X)$  to a pair  $(\tau, \eta) \in \mathbb{R}^m \times \text{Trop}(Y)$  one is tempted to proceed as follows. Let  $\ell$  be the line represented by  $\xi$ , let  $\tau \in \mathbb{R}^m$  be such that  $-\tau + \mathbb{R}(1, \dots, 1)$  is a point on  $\ell$ , and set  $\eta := -A\tau + \xi$ . Then  $\eta$  represents the

translate of  $\ell$  over  $-\tau$ , which therefore passes through 0. By construction, the pair  $(\tau, \eta)$  satisfies  $A\tau + \eta = \xi$ , so that the valuation  $\sigma(\xi) := \mu(\tau, \sigma_Y(\eta))$  maps to  $\xi$ .

There are various problems with this definition of  $\sigma$ , but we can sharpen it as follows. A first, minor problem is that if  $\xi = \infty$ , then  $\xi$  does not represent a line. In that case, we just set  $\sigma(\xi)$  equal to  $\infty \in X^{\text{an}}$ . The second, and more serious, problem is that  $\ell$  may not contain points  $\tau \in \mathbb{R}^m / \mathbb{R}(1, \dots, 1)$ . To remedy this, we will use a stratification of  $X = \widehat{\text{Gr}}(2, m)$  and  $\text{Trop}(X)$  defined as follows (and also used, in slightly different terminology, in [CHW]). Let  $J$  be any subset of  $[m]$  of cardinality  $\geq 2$ , set  $I := [m] \setminus J$ , and let  $X_J$  denote the subset of  $X$  where

$$\{i \in [m] \mid x_{ij} = 0 \text{ for all } j \in [m]\} = I.$$

So  $X_{[m]}$  is the largest one among these strata, and it parameterises lines that intersect  $\mathbb{G}_m^m$ . Each  $X_J$  is the dense set in the (cone over the) smaller Grassmannian of 2-spaces contained in  $\mathbb{A}^J \times \{0\}^I$  consisting of all spaces that intersect  $\mathbb{G}_m^J \times \{0\}^I$ . Similarly, non-infinity points of  $\text{Trop}(X_J)$  parameterise lines in the face  $\Delta_J$  of  $\Delta$  (where all  $I$ -coordinates are  $\infty$ ) that intersect the relative interior of  $\Delta_J$ . Let  $Y_J$  denote the  $J$ -analogue of  $Y$ , that is, the subspace of  $K^{\binom{J}{2}}$  parameterised by  $(y_j - y_{j'})_{j < j'}$ , identified with a subspace of  $K^{\binom{m}{2}}$  by extending with zero coordinates. Then  $Y_J \setminus \{0\}$  is a subset of  $X_J$ , and in fact we have  $\varphi(\mathbb{G}_m^J) \cdot (Y_J \setminus \{0\}) = X_J$ . Note also that  $Y_J$  is not a *subspace* of  $Y = Y_{[m]}$  but rather its *image* under projecting some coordinates to 0. We also define  $Y_\emptyset$  as the point  $\{0\}$ .

We choose  $\tau \in \mathbb{R}_\infty^m$  as a function of  $\xi$  as follows. If  $\xi = \infty$ , then set  $\tau := \infty$ . Otherwise, let  $H$  be the tropical hyperplane in  $\Delta$  with the tropical equation  $\zeta_1 \oplus \dots \oplus \zeta_m$ , and let  $\tau$  represent the stable intersection of  $H$  and the line  $\ell$  represented by  $\xi$ . By continuity of stable intersection, the projective point  $\tau + \mathbb{R}(1, \dots, 1)$  depends continuously on non-infinite  $\xi$ .

Next, we choose  $\eta$  as a function of  $\xi$ . If  $\xi = \infty$ , then  $\eta := \infty$ . Otherwise, let  $J$  be such that  $\xi \in \text{Trop}(X_J)$ . Then  $\ell$  lies in  $\Delta_J$  and intersects the relative interior of  $\Delta_J$ . As a consequence,  $\tau_j \neq \infty$  if and only if  $j \in J$ . Set  $\eta_{ij} := \xi_{ij} - \tau_i - \tau_j$  for  $i, j \in J$  and  $\eta_{ij} := \infty$  if one of  $i, j$  lies in  $I$ . Then  $\eta$  lies in  $\text{Trop}(Y_J)$ .

The pair  $(\tau, \eta)$  thus constructed does not depend continuously on  $\xi$ , but we claim that the valuation

$$\sigma(\xi) := \mu(\tau, \sigma_{Y_J}(\eta)) \in X^{\text{an}}$$

does. Note that we abuse notation slightly, since  $\tau$  will in general have some coordinates equal to  $\infty$ —but one readily verifies that, since  $A$  contains only non-negative entries,  $\mu$  extends to  $\mathbb{R}_\infty^m \times X^{\text{an}}$ . First observe that tropically scaling all coordinates of  $\tau$  with  $c \in \mathbb{R}$  and all coordinates of  $\eta$  with  $-2c$  leads to the same valuation. Now let  $\xi^{(p)}, p = 1, 2, 3, \dots$  be a sequence of points in  $\text{Trop}(X)$  that converges to a non-infinity limit  $\xi \in \text{Trop}(X_J)$  with  $|J| \geq 2$ . After deleting an initial segment of the sequence, we may assume that each  $\xi^{(p)}$  lies in some  $\text{Trop}(X_{J^{(p)}})$  with  $J^{(p)} \supseteq J$ . Let  $\eta^{(p)} \in \text{Trop}(Y_{J^{(p)}})$  and  $\tau^{(p)} \in \mathbb{R}^m$  be the corresponding points, so that  $\xi^{(p)} = A\tau^{(p)} + \eta^{(p)}$  for all  $p$ . The projective points  $\tau^{(p)} + \mathbb{R}(1, \dots, 1)$  converge to  $\tau + \mathbb{R}(1, \dots, 1)$  (by continuity of stable intersection). Hence, after suitable tropical scalings of the  $\tau^{(p)}$  and the  $\eta^{(p)}$ , we achieve that  $\tau^{(p)} \rightarrow \tau$  for  $p \rightarrow \infty$ . Then for  $i, j \in J$  we find that

$$\eta_{ij}^{(p)} = \xi_{ij}^{(p)} - \tau_i^{(p)} - \tau_j^{(p)} \rightarrow \xi_{ij} - \tau_i - \tau_j = \eta_{ij} \text{ for } p \rightarrow \infty.$$




FIGURE 2. Basis exchange in the case  $J^{(p)} \setminus J = \{i\}$ , with  $\eta_{ij}^{(p)} \leq \eta_{ij'}^{(p)} \leq \eta_{ij''}^{(p)}$ .

We now argue that for each  $\mathbb{G}_m^m$ -homogeneous element  $f \in K[X]$  the value  $\sigma(\xi^{(p)})(f)$  converges to  $\sigma(\xi)(f)$ . Let  $\beta \in \mathbb{N}^m$  be the weight of  $f$ . If  $\beta_i > 0$  for some  $i \notin J$ , then  $f$  lies in the ideal generated by the coordinates  $x_{ij}$  for which one of  $i, j$  does not lie in  $J$ . In this case,  $\sigma(\xi)(f) = \infty$ . To see that  $\sigma(\xi^{(p)})(f)$  tends to infinity, expand  $f = \sum_{ij} x_{ij} f_{ij}$  where the sum is over pairs  $(i, j)$  that are not both in  $J$ . Then we have

$$\sigma(\xi^{(p)})(f) \geq \min_{ij} (\xi_{ij}^{(p)} + \sigma(\xi^{(p)})(f_{ij})).$$

Since each  $\xi_{ij}^{(p)}$  tends to infinity and each  $\sigma(\xi^{(p)})(f_{ij})$  is bounded from below, we find the desired convergence (a similar convergence argument applies when the limit  $\xi$  equals  $\infty$ ). If  $\beta_i = 0$  for all  $i \notin J$ , then  $f$  depends only on the coordinates  $x_{ij}$  with  $i, j \in J$ , and it suffices to show that

$$\sigma_{Y_{J^{(p)}}}(\eta^{(p)})(f) \rightarrow \sigma_{Y_J}(\eta)(f), \quad p \rightarrow \infty.$$

Using the definition of  $\sigma$  and the fact that  $\eta_{ij}^{(p)} \rightarrow \eta_{ij}$  for  $p \rightarrow \infty$  and  $i, j \in J$ , this convergence follows if there exists a tree on  $J^{(p)}$  compatible with  $\eta^{(p)}$  which contains a spanning tree on  $J \subseteq J^{(p)}$ . But this is a consequence of the basis exchange axiom: start with any tree  $\Gamma$  on  $J^{(p)}$  compatible with  $\eta^{(p)}$ . If the induced forest  $\Gamma|_J$  on  $J$  is not connected, pick arbitrary endpoints  $j, j' \in J$  that belong to different connected components of  $\Gamma|_J$ . Then replace, in  $\Gamma$ , the edge in the simple path from  $j$  to  $j'$  of smallest  $\eta^{(p)}$ -weight by  $jj'$  (which has the same weight). This creates a new  $\Gamma$  compatible with  $\eta^{(p)}$  such that  $\Gamma|_J$  has fewer connected components than before. Proceed in this fashion until  $\Gamma|_J$  is connected. See Figure 2 for an illustration of this procedure. This concludes the proof that  $\sigma$  is a continuous section  $\text{Trop}(X) \rightarrow Z$  of the surjection  $Z \rightarrow \text{Trop}(X)$ .  $\square$

*Proof of Theorem 5.1, naturality and equivariance.* In the previous proof, we decomposed  $\xi$  as  $A\tau + \eta$  by choosing for  $-\tau$  a point on the tropical line  $\ell$  represented by  $\xi$ . This point was obtained by stably intersecting  $\ell$  with a hypersurface. By verifying the conditions of Proposition 4.4, we now show that the chosen decomposition is, in fact, irrelevant for the section  $\sigma$ .

First, if  $\eta \in \text{Trop}(Y^0)$  corresponds to a tropical projective line  $\ell$ , then the  $\tau \in \mathbb{R}^m$  for which  $A\tau + \eta$  lies in  $\text{Trop}(Y)$  are those for which  $-\tau$  lies in  $\ell$ . Thus  $T_\eta = -\ell$  is connected. This settles the first requirement for Proposition 4.4.

Now let  $\eta \in \text{Trop}(Y^0)$  and  $\tau \in \mathbb{R}^m$  such that  $[0, \tau]$  lies in  $T_\eta$ . For sufficiently small  $\epsilon > 0$  both  $\eta$  and  $\eta' := A\epsilon\tau + \eta$  are compatible with a fixed tree  $\Gamma$ . By shrinking  $\epsilon$  we may, moreover, assume that there exists an edge  $i_0j_0$  in  $\Gamma$  such that both

$$\eta_{i_0j_0} \leq \eta_{ij} \text{ and } \eta'_{i_0j_0} \leq \eta'_{ij} \text{ for all } ij \in \Gamma$$

(and hence also for all  $ij \in K_m \setminus \Gamma$ ). The edge  $i_0j_0$  cuts the tree  $\Gamma$  into two connected components (see Figure 1). Let  $[m] = I \cup J$  be the vertex sets of these connected components, with  $i_0 \in I$  and  $j_0 \in J$ . We claim that  $\tau_i = \tau_{i_0}$  for all  $i \in I$  and  $\tau_j = \tau_{j_0}$  for all  $j \in J$ . Indeed, pick  $j \in J$  and consider the cycle formed by  $i_0, j$ , and then back along  $\Gamma$  to  $i_0$ . We have  $\eta'_{i_0j} = \eta'_{i_0j_0}$ , since this is the edge of  $\Gamma$  in said cycle with smallest  $\eta'$ -weight. On the other hand, we have

$$\eta'_{i_0j} = \eta_{i_0j} + \epsilon(\tau_{i_0} + \tau_j) = \eta_{i_0j_0} + \epsilon(\tau_{i_0} + \tau_j)$$

and

$$\eta'_{i_0j_0} = \eta_{i_0j_0} + \epsilon(\tau_{i_0} + \tau_{j_0}).$$

This shows that  $\tau_j = \tau_{j_0}$ . Similarly, we find that for all  $i \in I$  we have  $\tau_i = \tau_{i_0}$ .

To construct  $t$ , we may assume that one of  $\tau_{i_0}, \tau_{j_0}$  is zero and the other is non-negative—indeed, this can be achieved by adding a multiple of the all-one vector to  $\tau$ , which can be mimicked by multiplying  $t$  with a scalar of the right valuation. Without loss of generality, suppose that  $\epsilon\tau_{i_0} =: a$  is non-negative and  $\tau_{j_0}$  is zero. Then adding  $\epsilon A\tau$  to  $\eta$  has the effect of increasing all  $\eta_{ij}$  with  $i, j \in I$  by  $2a$ , keeping all  $\eta_{ij}$  with  $i, j \in J$  constant, and increasing all  $\eta_{ij}$  with  $i \in I$  and  $j \in J$  by  $a$ . As, by assumption, the minimal-weight edge in  $\Gamma$  remains the edge  $i_0j_0$ , the minimal  $\eta$ -weight of an edge of  $\Gamma$  with both vertices in  $J$  must be at least  $\eta_{i_0j_0} + a$ .

Now let  $w = \sigma_Y(\eta)$  and let  $y \in Y^0$  be the generic point. Its coordinates are  $x_{ij} = (y_i - y_j)_{ij}$  where the  $y_i$  are variables. It represents the subspace spanned by the rows of the matrix

$$\begin{bmatrix} y_1 & y_2 & \cdots & y_m \\ 1 & 1 & \cdots & 1 \end{bmatrix}$$

A point  $t$  sends  $y$  into  $Y^0$  if and only if it sends this subspace to another subspace containing the all-one vector, hence if and only if  $t^{-1}$  lies in the subspace. Hence we make the *Ansatz*

$$t_i = \frac{1}{c(y_i - y_{j_0}) + d},$$

where  $c, d$  are still to be determined. Now for each  $i$  the expression  $y_i - y_{j_0}$  expands as a sum of the  $x$ -variables corresponding to the edges on the path in  $\Gamma$  from  $i$  to  $j_0$ , and  $w(y_i - y_{j_0})$  equals the minimal  $\eta$ -weight among these edges. For  $i \in I$  this equals  $\eta_{i_0j_0}$ , as this is the minimal-weight edge overall. Thus we choose  $c \in K$  such that  $v(c) + \eta_{i_0j_0} = -a < 0$ . For  $i \in J$  the minimal weight along the path is at least  $a + \eta_{i_0j_0}$ , so if we choose a  $d \in K$  with valuation 0, then the denominator in the *Ansatz* gets the right valuation for both  $i \in I$  and  $i \in J$ .

We have thus constructed a  $t \in \mathbb{G}_m(K(Y))$  with  $w(t) = \epsilon\tau$  and  $\varphi(t)y \in Y(K(Y))$ . As all requirements of Proposition 4.4 are met, we have constructed an  $\mathbb{R}^m$ -equivariant section  $\text{Trop}(X^0) \rightarrow Z^0$ . This section agrees with the restriction to  $\text{Trop}(X^0)$  of the

section constructed in the previous proof. Hence the continuous section constructed there is  $\mathbb{R}^m$ -equivariant.  $\square$

**Remark 5.2.** In [CHW] the setting is projective rather than affine. Theorem 1.1 and Corollary 7.3 from that paper follow from our theorem by applying Lemmas 3.5 and 3.4, respectively.

## 6. RANK-TWO MATRICES

In this section we take  $n = m \cdot p$  and consider  $\mathbb{A}^n$  with coordinates  $x_{ij}$  with  $1 \leq i \leq m$  and  $1 \leq j \leq p$ . Let  $X \subset \mathbb{A}^n$  be the image of the polynomial map

$$\psi : (\mathbb{A}^m)^2 \times (\mathbb{A}^p)^2 \rightarrow \mathbb{A}^n, \quad (y, y'), (z, z') \mapsto (y_i z'_j - y'_i z_j)_{1 \leq i \leq m, 1 \leq j \leq p}.$$

It is the variety of matrices of rank at most two, and also the affine cone over the variety of secant lines of the Segre embedding of  $\mathbb{P}^{m-1} \times \mathbb{P}^{p-1}$  in  $\mathbb{P}^{n-1}$ .

Let  $Y$  be the subvariety of  $X$  defined as the image of

$$(\mathbb{A}^m \times \{\mathbf{1}\}) \times (\mathbb{A}^p \times \{\mathbf{1}\})$$

via  $\psi$ . Points of  $Y$  have coordinates  $x_{ij} = (y_i - z_j)$  in  $\mathbb{A}^n$ , so that  $Y$  is the zero locus of the linear forms

$$(2) \quad x_{ij} + x_{lk} = x_{ik} + x_{lj}, \quad 1 \leq i < l \leq m, 1 \leq j < k \leq p.$$

Consider also the homomorphism of tori given by

$$\varphi : \mathbb{G}_m^m \times \mathbb{G}_m^p \rightarrow \mathbb{G}_m^n, \quad (t, s) \mapsto (t_i s_j)_{1 \leq i \leq m, 1 \leq j \leq p}.$$

The corresponding  $n \times (m+p)$ -matrix  $A$  has a one-dimensional kernel spanned by  $(1, \dots, 1, -1, \dots, -1)$ . We have  $X = \overline{\varphi(\mathbb{G}_m^m \times \mathbb{G}_m^p) \cdot Y}$  and

$$\text{Trop}(X^0) = A(\mathbb{R}^m \times \mathbb{R}^p) + \text{Trop}(Y^0)$$

where  $X^0 \subseteq X$  and  $Y^0 \subseteq Y$  are the loci where no coordinate is zero. Let  $\mu : \mathbb{R}^{m+p} \times X^{\text{an}} \rightarrow Z \subseteq X^{\text{an}}$  be as constructed in Section 3. We will prove the following theorem.

**Theorem 6.1.** *The surjection  $X^{\text{an}} \rightarrow \text{Trop}(X)$ , where  $X$  is the variety of  $m \times p$ -matrices of rank at most two, has a continuous,  $\mathbb{R}^m$ -equivariant section  $\text{Trop}(X) \rightarrow Z$ .*

Note that we do not claim that the section is also  $\mathbb{R}^p$ -equivariant. While this might be the case, our construction below does not yield this.

For the proof of this theorem, we need to understand points in  $\text{Trop}(X)$  and its tropical subvariety  $\text{Trop}(Y)$ . By [DSS05, Corollary 3.8], a matrix  $\xi \in \mathbb{R}^{m \times p}$  lies in  $\text{Trop}(X^0)$  if and only if it has tropical rank at most 2, i.e., if and only if all its  $3 \times 3$ -submatrices are tropically singular. This extends directly to all of  $\text{Trop}(X)$ . To understand  $\text{Trop}(Y)$  note that the matroid defined by  $Y$  is the graphical matroid of the complete bipartite graph  $K_{m,p}$ ; this is immediate from the parameterisation  $x_{ij} = y_i - z_j$ . In other words,  $\eta \in \mathbb{R}_\infty^{m \times p}$  lies in  $\text{Trop}(Y)$  if and only if along each cycle in  $K_{m,p}$  the minimal  $\eta$ -weight of an edge is attained at least twice. We claim that this is equivalent to the condition that in every  $2 \times 2$ -submatrix of  $\eta$  the minimal entry appears at least twice. Indeed, necessity of the latter condition is obvious, as any  $2 \times 2$ -submatrix records the weights of a 4-cycle in  $K_{m,p}$ . For sufficiency, assume that the minimal  $\eta$ -weight in every 4-cycle is attained at least twice, and let  $C$  be a general (simple, even) cycle in  $K_{m,p}$ . Label  $C$  as  $i_1 - j_1 - i_2 - j_2 - \dots - i_a - j_a - i_1$ ,






FIGURE 3. Only four-cycles need to be tested for membership of  $\text{Trop}(Y^0)$ .

where the  $i$ s are in  $[m]$  and the  $j$ s are in  $[n]$  and where  $\alpha := \eta_{i_1, j_1}$  is the minimal weight of an edge in  $C$ . Assume, for a contradiction, that all other edges in  $C$  have  $\eta$ -weight strictly larger than  $\alpha$ . Then in the 4-cycle  $i_1 - j_1 - i_2 - j_2 - i_1$  the weight  $\eta_{i_1, j_2}$  must be  $\alpha$ . Next, in the 4-cycle  $i_1 - j_2 - i_3 - j_3 - i_1$  the weight  $\eta_{i_1, j_3}$  must also equal  $\alpha$ , etc. In this manner we find that  $\eta_{i_1, j_a}$  must also equal  $\alpha$ , a contradiction. See Figure 3 for an illustration. Armed with this characterisation of  $\text{Trop}(Y)$  we will now prove the theorem.

*Proof of Theorem 6.1.* As in the proof of Theorem 5.1, we use a stratification of  $X$ . For  $I \subseteq [m]$  and  $J \subseteq [p]$  let  $X_{IJ}$  denote the locus in  $X$  consisting of  $x$  such that the rows of  $x$  labelled by  $[m] \setminus I$  and the columns of  $x$  labelled by  $[p] \setminus J$  are identically zero and the submatrix  $x[I, J]$  does not have identically zero rows or columns. Let  $Y_{IJ}$  denote the  $(I, J)$ -analogue of  $Y$ . It is the image of  $Y$  under the map sending all coordinates outside the  $[I, J]$ -submatrix to zero.

For  $\xi \in \text{Trop}(X_{IJ})$  we let  $\tau \in \mathbb{R}_\infty^m$  be the tropical product  $\xi \odot (0, \dots, 0)^T$ , a point in the tropical convex hull of the columns of  $\xi$ . Then we have  $\tau_i \neq \infty$  if and only if  $i \in I$ . Let  $\xi' \in \text{Trop}(X_{IJ})$  be the matrix obtained from  $\xi$  by subtracting  $\tau_i$  from each  $\xi_{ij}$  with  $i \in I, j \in J$ . Then let  $\rho \in \mathbb{R}_\infty^p$  be the tropical product  $(0, \dots, 0) \odot \xi'$ , which records the minimal entry in each column of  $\xi'$ . Let  $\eta$  be the matrix obtained from  $\xi'$  by subtracting  $\rho_j$  from each  $\xi'_{ij}$  with  $i \in I, j \in J$ . By [DSS05, Lemma 6.2], the matrix  $\eta[I, J]$  has the property that in each of its  $2 \times 2$ -submatrices the minimal entry appears at least twice. By the discussion preceding the proof,  $\eta$  lies in  $\text{Trop}(Y_{IJ})$ .

We set

$$\sigma(\xi) := \mu((\tau, \rho), \sigma_{Y_{IJ}}(\eta)),$$

and claim that this depends continuously on  $\xi$ . To see this, let  $\xi^{(q)}$ ,  $q = 1, 2, \dots$  be a sequence in  $\text{Trop}(X)$  converging to  $\xi \in \text{Trop}(X_{IJ})$ , and construct  $\tau^{(q)}$  and  $\rho^{(q)}$  and  $\eta^{(q)}$  as above. After dropping finitely many initial terms, we have  $\xi^{(q)} \in \text{Trop}(X_{I^{(q)}J^{(q)}})$  with  $I^{(q)} \supseteq I$  and  $J^{(q)} \supseteq J$ . For  $i \in I$  and  $j \in J$  we find that  $\tau_i^{(q)} \rightarrow \tau_i$  for  $q \rightarrow \infty$  and also  $\lim_{q \rightarrow \infty} \rho_j^{(q)} = \rho_j$  and  $\lim_{q \rightarrow \infty} \eta_{ij}^{(q)} = \eta_{ij}$ .

Let  $f$  be a  $\mathbb{G}_m^{m+p}$ -weight element of  $K[X]$ . We have the same dichotomy as in the proof for the Grassmannian case: either  $f$  lies in the ideal generated by all variables  $x_{ij}$  with  $i \notin I$  or  $j \notin J$ , and in this case

$$\sigma(\xi^{(q)})(f) \rightarrow \infty = \sigma(\xi)(f) \text{ for } q \rightarrow \infty;$$

or  $f$  lies in the ring generated by the  $x_{ij}$  with  $i, j \in J$ . In the latter case, it suffices to show that

$$\sigma_{Y_{I^{(q)}J^{(q)}}}(\eta^{(q)})(f) \rightarrow \sigma_{Y_{I,J}}(\eta)(f).$$

Proceeding as for the Grassmannian of planes, we find that there exists, for each  $q$ , a tree  $\Gamma_q$  compatible with  $\eta^{(q)}$  that induces a tree (rather than a forest) on  $I \cup J$ . Using this tree, the right-hand side can be expressed in terms of coefficients  $\eta_{ij}^{(q)}$  with  $i \in I$  and  $j \in J$ . These converge to  $\eta_{ij}$ , and this gives the required convergence.

The map  $\text{Trop}(X) \rightarrow \mathbb{R}_{\infty}^m, \xi \rightarrow \xi \odot (0, 0, \dots, 0)^T = \tau$  is  $\mathbb{R}^m$ -equivariant, and this implies that  $\sigma$  is  $\mathbb{R}^m$ -equivariant. But the construction  $\xi \mapsto \rho$  is not  $\mathbb{R}^p$ -equivariant.  $\square$

**Remark 6.2.** The proof above uses only the technique of Proposition 4.1. Therefore, it is not as satisfactory as the proof for Grassmannians of two-spaces in Section 5, which used the technique of Proposition 4.4 to prove that the defined section is independent of the decomposition  $\xi = A\tau + \eta$  and hence equivariant. We have tried to mimick the proof for the Grassmannian, but failed because for suitable  $\eta \in \text{Trop}(Y^0)$  the set  $T_\eta$  can have dimension much larger than the expected four dimensions. This implies that the second requirement in Proposition 4.4 cannot be satisfied. Of course, this does not rule out the existence of alternative techniques for proving  $\mathbb{R}^{m+p}$ -equivariance.

## 7. A-DISCRIMINANTS

Linear spaces smeared around by tori, as discussed in Section 4, arise in the study of  $A$ -discriminants from [GKZ94]. Let  $\varphi : \mathbb{G}_m^m \rightarrow \mathbb{G}_m^n$  be a torus homomorphism with corresponding integer  $n \times m$ -matrix  $A$ , and let  $V$  be the closure in  $\mathbb{A}^n$  of the image of  $\varphi$ , a toric variety. The linear action of  $\mathbb{G}_m^m$  on  $\mathbb{A}^n$  gives rise to an action on the dual space  $(\mathbb{A}^n)^\vee$ , given by a torus homomorphism  $\varphi^\vee : \mathbb{G}_m^m \rightarrow \mathbb{G}_m^n$  corresponding to the matrix  $-A$ .

Let  $Y \subseteq (\mathbb{A}^n)^\vee$  be the annihilator of the tangent space  $T_{\varphi(1)}V$ . Since  $A$ , when regarded as a matrix over  $K$ , is the derivative of  $\varphi$  at 1,  $Y$  is the orthogonal complement of the column space of  $A$ . For  $t \in \mathbb{G}_m^m$ ,  $\varphi(t)$  maps  $T_{\varphi(1)}V$  into  $T_{\varphi(t)}V$ , hence we find that  $\varphi^\vee(t)$  maps  $V$  into the annihilator of  $T_{\varphi(t)}V$ . Thus the variety  $X$  defined as the Zariski closure of the union of these annihilators equals  $\overline{\varphi^\vee(\mathbb{G}_m^m) \cdot Y}$ . This is known as the Horn uniformisation of the *dual variety* of  $V$ . It was used in [DFS07] to characterise  $\text{Trop}(X^0)$  as

$$\text{Trop}(X^0) = -A\mathbb{R}^m + \text{Trop}(Y^0),$$

where, of course, the minus sign is only a reminder of the contragredience of the action of  $\mathbb{G}_m^m$  on  $(\mathbb{A}^n)^\vee$  and can also be left out. This leads to the following fundamental problem.

**Problem 7.1.** *For which torus homomorphisms  $\varphi : \mathbb{G}_m^m \rightarrow \mathbb{G}_m^n$  does the map from the analytification of the dual variety  $X$  of  $V = \overline{\text{im}\varphi} \subseteq \mathbb{A}^n$  to  $\text{Trop}(X)$  admit a continuous,  $\mathbb{R}^m$ -equivariant section into the subset  $Z \subseteq \text{Trop}(X)$  defined in Section 3?*

We do not have any general results at this point. Instead, we now consider the very special case of *Cayley's hyperdeterminant*, and we stay away from zero coordinates.




FIGURE 4. The tropical variety of Cayley's hyperdeterminant has 8 triangles and 6 quadrangles.

**Example 7.2.** Let  $n = 2^3$  and use coordinates  $x_{ijk}$ ,  $i, j, k \in \{0, 1\}$  on  $\mathbb{A}^8$ . Let  $m = 3 \cdot 2$  and use coordinates  $t_i, u_j, v_k$ ,  $i, j, k \in \{0, 1\}$  on  $\mathbb{G}_m^6$ . Let  $\varphi$  be the map  $(t, u, v) \mapsto (t_i u_j v_k)_{i,j,k}$ . Then  $V^0$  is the variety of rank-one tensors of format  $2 \times 2 \times 2$ . The dual variety  $X$  is a hypersurface whose defining equation is Cayley's hyperdeterminant

$$\begin{aligned} \Delta = & x_{000}^2 x_{111}^2 + x_{001}^2 x_{110}^2 + x_{010}^2 x_{101}^2 + x_{100}^2 x_{011}^2 \\ & - 2x_{000}x_{001}x_{110}x_{111} - 2x_{000}x_{010}x_{101}x_{111} - 2x_{000}x_{011}x_{100}x_{111} \\ & - 2x_{001}x_{010}x_{101}x_{110} - 2x_{001}x_{011}x_{110}x_{100} - 2x_{010}x_{011}x_{101}x_{100} \\ & + 4x_{000}x_{011}x_{101}x_{110} + 4x_{001}x_{010}x_{100}x_{111}. \end{aligned}$$

The tropical variety of  $X$  is known explicitly (though we will not use this knowledge): modulo its four-dimensional lineality space it is a 3-dimensional fan in 4-space. Intersecting with a 3-dimensional sphere yields a 2-dimensional spherical polyhedral complex, which consists of two nested tetrahedra glued by quadrangles along corresponding edges; see Figure 4.

The matrix  $A$  sends  $\tau = (\rho, \delta, \nu) \in \mathbb{R}^6$  to the  $2 \times 2 \times 2$ -array with entries  $(\rho_i + \delta_j + \nu_k)_{ijk}$ . The kernel of this map consists of vectors of the form  $(a\mathbf{1}, b\mathbf{1}, c\mathbf{1})$  with  $a + b + c = 0$ , so the column space  $\text{im}A$  has dimension 4. It defines the matroid on the vertices of the three-dimensional cube in which independence is affine independence. Since the complement of any four affinely independent vertices of the cube is again affinely independent, this matroid is self-dual. So the dual matroid, which is the matroid of the linear space  $Y$ , is the same matroid on 8 elements.

Up to symmetries of the cube, the seven-dimensional polyhedral fan  $\text{Trop}(Y^0)$  has six maximal cones, and they are depicted in Figure 5. Among these, the cones of type IIa, IIb, and IIIa lie in  $A\mathbb{R}^6$  plus the union of the cones of type I, IIIb, IIIc. For instance, take the array in type IIIa and add  $(c-b)/2, 0, (b-c)/2$  to the positions with entries  $b, a, (c \text{ and } d)$ , respectively. The array thus added lies in the column space of  $A$ , and the result is an array of type IIIb (with  $b$  and  $c$  replaced by  $(b+c)/2$  and  $d$  replaced by  $d + (b-c)/2$ ).

Now let  $C$  be a cone of type I, IIIb, or IIIc. Then the linear span of  $C$  intersects  $A\mathbb{R}^m$  only in scalar multiples of the all-one array. This follows from the fact that the span in  $\mathbb{R}^3$  of the differences of vertices of the cube with the same label ( $a$  or  $b$ ) is all of  $\mathbb{R}^3$  (this is not true for the other types!). Thus on  $A\mathbb{R}^m + C \subseteq \text{Trop}(X^0)$




FIGURE 5. The six orbits of maximal cones in  $\text{Trop}(Y^0)$ , with  $a \leq b \leq c \leq d$ .

we can define a section  $\sigma_C$  into  $\text{Trop}(X^0)$  as follows: write  $\xi$  as  $A\tau + \eta$  with  $\eta \in C$  and set  $\sigma(\xi) := \mu(\tau, \sigma_Y(\eta))$ . Note that, for any  $c \in \mathbb{R}$ , subtracting  $(c\mathbf{1}, c\mathbf{1}, c\mathbf{1})$  from  $\tau$  and adding  $3c$  times the all-one array to  $\eta$  yields the same value for  $\sigma(\xi)$ , so that  $\sigma$  is well-defined on  $\mathbb{A}\mathbb{R}^m + C$ .

Next we verify that if  $C'$  is a second cone of type I, IIIb, or IIIc, then  $\sigma_C$  and  $\sigma_{C'}$  agree on the intersection  $(\mathbb{A}\mathbb{R}^m + \sigma_C) \cap (\mathbb{A}\mathbb{R}^m + C')$ . This is immediate if

$$(\mathbb{A}\mathbb{R}^m + C) \cap (\mathbb{A}\mathbb{R}^m + C') = \mathbb{A}\mathbb{R}^m + (C \cap C')$$

as the recipes defining  $\sigma_C$  and  $\sigma_{C'}$  agree on the right-hand side. For each choice of  $C$  and  $C'$ , a vector witnessing that the left-hand side is *strictly larger* than the right-hand side can be found by solving a number of linear programs. If none of these linear programs turns out to be feasible, then equality holds. We have performed this test for all choices of  $C$  in the cones I, IIIb, IIIc, and  $C'$  in one of the orbits of these cones. Together with Proposition 4.1 this proves the following theorem.

**Theorem 7.3.** *Let  $X \subseteq K^{2 \times 2 \times 2}$  be the hypersurface defined by Cayley's hyperdeterminant, equipped with the natural action of  $\mathbb{G}_m^2 \times \mathbb{G}_m^2 \times \mathbb{G}_m^2$ . Let  $X^{\text{an}} \rightarrow Z$  be the retraction defined relative to this torus action. Then the surjection  $Z^0 \rightarrow \text{Trop}(X^0)$  has a continuous,  $\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$ -equivariant section.  $\diamond$*

#### REFERENCES

- [AK06] Federico Ardila and Caroline J. Klivans. The Bergman complex of a matroid and phylogenetic trees. *J. Comb. Theory, Ser. B*, 96(1):38–49, 2006.
- [Ber90] Vladimir G. Berkovich. *Spectral theory and analytic geometry over non-archimedean fields*, volume 33 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1990.
- [BG84] Robert Bieri and John R.J. Groves. The geometry of the set of characters induced by valuations. *J. reine angew. Math.*, 347:168–195, 1984.

- [BPR] Matthew Baker, Sam Payne, and Joseph Rabinoff. Nonarchimedean geometry, tropicalization, and metrics on curves. Preprint, available from <http://arxiv.org/abs/1104.0320>.
- [CHW] Maria Angelica Cueto, Mathias Hübich, and Annette Werner. Faithful tropicalization of the Grassmannian of planes. *Math. Ann.* To appear; preprint available from <http://arxiv.org/abs/1309.0450>.
- [DFS07] Alicia Dickenstein, Eva Maria Feichtner, and Bernd Sturmfels. Tropical discriminants. *J. Am. Math. Soc.*, 20(4):1111–1133, 2007.
- [Dra08] Jan Draisma. A tropical approach to secant dimensions. *J. Pure Appl. Algebra*, 212(2):349–363, 2008.
- [DSS05] Mike Develin, Francisco Santos, and Bernd Sturmfels. On the rank of a tropical matrix. In *Combinatorial and computational geometry*, pages 213–242. Cambridge: Cambridge University Press, 2005.
- [EKL06] Manfred Einsiedler, Mikhail Kapranov, and Douglas Lind. Non-archimedean amoebas and tropical varieties. *J. reine angew. Math.*, 601:139–157, 2006.
- [GKZ94] Israel M. Gelfand, Mikhail M. Kapranov, and Andrei V. Zelevinsky. *Discriminants, resultants, and multidimensional determinants*. Mathematics: Theory & Applications. Birkhäuser, Boston, MA, 1994.
- [GRW] Walter Gubler, Joe Rabinoff, and Annette Werner. Skeletons and tropicalizations. Preprint, soon available from [arxiv](http://arxiv.org).
- [Mik06] Grigory Mikhalkin. Tropical geometry and its applications. In Marta Sanz-Solé *et al.*, editor, *Proceedings of the international congress of mathematicians, Madrid, Spain, August 22–30, 2006. Volume II: Invited lectures.*, Zürich, 2006. European Mathematical Society.
- [MS] Diane MacLagan and Bernd Sturmfels. *Tropical Geometry*. Forthcoming, available from [homepages.warwick.ac.uk/staff/D.MacLagan/](http://homepages.warwick.ac.uk/staff/D.MacLagan/).
- [Pay09] Sam Payne. Fibers of tropicalization. *Math. Z.*, 262(2):301–311, 2009.
- [RGST05] Jürgen Richter-Gebert, Bernd Sturmfels, and Thorsten Theobald. First steps in tropical geometry. In G. L. *et al.* Litvinov, editor, *Idempotent Mathematics and Mathematical Physics. Proceedings of the International Workshop, Vienna, Austria, February 3–10, 2003*, volume 377 of *Contemporary Mathematics*, pages 289–317, Providence, RI, 2005. AMS.
- [Sch03] Alexander Schrijver. *Combinatorial Optimization. Polyhedra and Efficiency*. Number 24 in Algorithms and Combinatorics. Springer, Berlin, etc., 2003.
- [SS04] David E. Speyer and Bernd Sturmfels. The tropical Grassmannian. *Adv. Geom.*, 4(3):389–411, 2004.
- [YY07] Josephine Yu and Debbie S. Yuster. Representing tropical linear spaces by circuits. In *Proceedings of the 19th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2007), July 2007, Tianjin, China*, 2007. Preprint available from <http://arxiv.org/abs/math/0611579>.

(Jan Draisma) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, TECHNISCHE UNIVERSITEIT EINDHOVEN, P.O. BOX 513, 5600 MB EINDHOVEN, THE NETHERLANDS; AND CENTRUM VOOR WISKUNDE EN INFORMATICA, AMSTERDAM, THE NETHERLANDS  
*E-mail address:* [j.draisma@tue.nl](mailto:j.draisma@tue.nl)

(Elisa Postinghel) DEPARTMENT OF MATHEMATICS, KATHOLIEKE UNIVERSITEIT LEUVEN, CELESTIJNENLAAN 200B - BOX 2400, 3001 LEUVEN, BELGIUM  
*E-mail address:* [elisa.postinghel@wis.kuleuven.be](mailto:elisa.postinghel@wis.kuleuven.be)