

Networks of $M/G/\infty$ queues with shot-noise-driven arrival intensities

Citation for published version (APA):

Koops, D. T., Boxma, O. J., & Mandjes, M. R. H. (2016). *Networks of $M/G/\infty$ queues with shot-noise-driven arrival intensities*. (Report Eurandom; Vol. 2016010). Eurandom.

Document status and date:

Published: 30/09/2016

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EURANDOM PREPRINT SERIES

2016-010

September 30, 2016

**Networks of $\cdot/G/\infty$ Queues
with Shot-Noise-Driven Arrival Intensities**

D. Koops, O. Boxma, M. Mandjes
ISSN 1389-2355

Networks of $\cdot/G/\infty$ Queues with Shot-Noise-Driven Arrival Intensities

D.T. Koops¹, O.J. Boxma², and M.R.H. Mandjes¹

¹*Korteweg-de Vries Institute, University of Amsterdam*

²*Eurandom and Department of Mathematics and Computer Science, Eindhoven
University of Technology*

September 30, 2016

Abstract

We study infinite-server queues in which the arrival process is a Cox process (or doubly stochastic Poisson process), of which the arrival rate is given by shot noise. A shot-noise rate emerges naturally in cases where the arrival rate tends to exhibit sudden increases (or: shots) at random epochs, after which the rate is inclined to revert to lower values. Exponential decay of the shot noise is assumed, so that the queueing systems are amenable to analysis. In particular, we perform transient analysis on the number of customers in the queue jointly with the value of the driving shot-noise process. Additionally, we derive heavy-traffic asymptotics for the number of customers in the system by using a linear scaling of the shot intensity. First we focus on a one dimensional setting in which there is a single infinite-server queue, which we then extend to a network setting.

1 Introduction

In the queueing literature one has traditionally studied queues with Poisson input. The Poisson assumption typically facilitates explicit analysis, but it does not always align well with actual data, see e.g. [11] and references therein. More specifically, statistical studies show that in many practical situations, Poisson processes underestimate the variability of the queue's input stream. This observation has motivated research on queues fed by arrival processes that better capture the burstiness observed in practice.

The extent to which burstiness takes place, can be measured by the dispersion index, i.e. the ratio of the variance of the number of arrivals in a given interval, and the corresponding expected value. In arrival streams that display burstiness, the dispersion index is

larger than unity (as opposed to Poisson processes, for which it is equal to unity), a phenomenon that is usually referred to as *overdispersion*. It is desirable that the arrival process of the queueing model can take the observed overdispersion into account. One way to achieve this, is to make use of *Cox processes*, which are Poisson processes, conditional on the stochastic time-dependent intensity. It is an immediate consequence of the law of total variance, that Cox processes *do* have a dispersion index larger than unity. Therefore, this class of processes makes for a good candidate to model overdispersed input processes.

In this paper we contribute to the development of queueing models fed by input streams that exhibit overdispersion. We analyze infinite-server queues driven by a particular Cox process, in which the rate is a (stochastic) shot-noise process.

The shot-noise process that we use, is one in which there are only upward jumps (or: shots), that arrive according to a homogeneous Poisson process. Furthermore, we employ an exponential ‘response’ or ‘decay’ function, in which it is encoded how quickly the process will decline after a jump. In this case, the shot-noise process is a Markov process, see [16, p. 393]. There are several variations on shot-noise processes; see e.g. [10] for a comprehensive overview.

It is not a novel idea to use a shot-noise process as stochastic intensity. For instance, in insurance mathematics, the authors of [5] use a shot-noise-driven Cox process to model the claim count. They assume that disasters happen according to a Poisson process, and each disaster can induce a cluster of arriving claims. The disaster corresponds to a shot upwards in the claim intensity. As time passes, the claim intensity process decreases, as more and more claims are settled. In our queueing model, there could be similar underlying dynamics causing a shot-noise arrival rate. For instance, it could be that many people go to a website after a certain commercial has been on television.

Another example of shot-noise arrival processes is found in the famous paper [13], where it is used to model the occurrences of earthquakes. The arrival process considered in this paper has one crucial difference with the one used in the present paper: it makes use of Hawkes processes [9], which do have a shot-noise structure, but have the special feature that they are *self-exciting*. More specifically, in Hawkes processes, an arrival can induce a shot in the arrival rate, whereas in our shot-noise-driven Cox model these shots are merely exogenous. The Hawkes process is less tractable than the shot-noise-driven Cox process. A very recent effort to analyze $M/G/\infty$ queues that are driven by a Hawkes process has been made in [8], where a functional central limit theorem is derived for the number of customers in the system. In this model, it is much harder (or impossible) to obtain explicit results (in a non-asymptotic setting), as we are able to do in the shot-noise-driven Cox variant.

In order to successfully implement a theoretical model, it is crucial to have methods to estimate its parameters from data. The shot-noise-driven Cox process is attractive since it

has this property. Statistical methods that filter the unobservable intensity process, based on Markov Chain Monte Carlo (MCMC) techniques, have been developed; see [3] and references therein. By filtering, they refer to the estimation of the intensity process in a given time interval, given a realized arrival process. Subsequently, given a sample path of the intensity process, the parameters of the shot-noise process can be estimated by a Monte Carlo version of the expectation maximization (EM) method. Furthermore, the shot-noise-driven Cox process can also be easily simulated; see e.g. the thinning procedure described in [12].

In this paper we study networks of infinite-server queues with shot-noise-driven Cox input. We assume that the service times at a given node are i.i.d. samples from a general distribution. The output of a queue is routed to a next queue, or leaves the network. Infinite-server queues have the inherent advantage that customers do not interfere with one another, which considerably simplifies the analysis. Furthermore, infinite-server systems are frequently used to produce approximations for corresponding finite-server systems. In the network setting, we can model queueing systems that are driven by correlated shot-noise arrival processes. With regards to applications, such a system could represent the call centers of a fire department and police department in the same town.

The contributions and organization of this paper are as follows. In this paper we derive exact and asymptotic results. The main result of the exact analysis is Thm. 4.6, where we find the joint Laplace transform of the numbers of customers in the queues of a feed-forward network, jointly with the shot-noise-driven arrival rates. We build up towards this result along the following route. In Section 2 we introduce notation and we state the important Lemma 2.1 that we repeatedly rely on. Then we commence the analysis by deriving exact results for the single infinite-server queue, with a shot-noise arrival rate, in Section 3.1. Subsequently, in Section 3.2, we show that after an appropriate scaling the number of customers in the system satisfies a functional central limit theorem (Thm. 3.4); the limiting process is an Ornstein-Uhlenbeck (OU) process driven by a superposition of a Brownian motion and an integrated OU process. We then extend the theory to a network setting in Section 4. Before we consider full-blown networks, we first consider a tandem system consisting of an arbitrary number of infinite-server queues in Section 4.1. Then it is argued in Section 4.2 that a feedforward network can be seen as a number of tandem queues in parallel. We analyze two different ways in which dependency can enter the system through the arrival process. Firstly, in Model (M1), parallel service facilities are driven by a multidimensional shot-noise process in which the *shots* are simultaneous (which includes the possibility that all shot-noise processes are equal). Secondly, in Model (M2), we assume that there is one shot-noise arrival intensity that generates simultaneous *arrivals* in all tandems. In Section 5 we finish with some concluding remarks.

2 Notation and preliminaries

Let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ be a probability space, in which the filtration $\{\mathcal{F}_t\}_{t \geq 0}$ is such that $\Lambda(\cdot)$ is adapted to it. A shot-noise process is a process that has random jumps at Poisson epochs, and a deterministic ‘response’ or ‘decay’ function, which governs the behavior of the process. See [16, Section 8.7] for a brief account of shot-noise processes. The shot noise that we use in this paper has the following representation:

$$\Lambda(t) = \Lambda(0)e^{-rt} + \sum_{i=1}^{Y(t)} B_i e^{-r(t-t_i)}, \quad (1)$$

where the $B_i \geq 0$ are i.i.d. shots from a general distribution, the decay function is exponential with rate $r > 0$, Y is a homogeneous Poisson process with rate ν , and the epochs of the shots, that arrived before time t , are labelled $t_1, t_2, \dots, t_{Y(t)}$.

As explained in the introduction, the shot-noise process serves as a stochastic arrival rate to a queueing system. It is straightforward to simulate a shot-noise process; for an illustration of a sample path, consider Fig. 1. Using the thinning method for nonhomogeneous Poisson processes [12], and using the sample path of Fig. 1 as the arrival rate, one can generate a corresponding sample path for the arrival process, as is displayed in Fig. 2. Typically, most arrivals occur shortly after peaks in the shot-noise process in Fig. 1, as expected.

We write Λ (i.e., without argument) for a random variable with distribution equal to $\lim_{t \rightarrow \infty} \Lambda(t)$. We now present well-known transient and stationary moments of the shot-noise process, see Appendix B and e.g. [16]: with B distributed as B_1 ,

$$\begin{aligned} \mathbb{E} \Lambda(t) &= \Lambda(0)e^{-rt} + \frac{\nu \mathbb{E} B}{r}(1 - e^{-rt}), & \mathbb{E} \Lambda &= \frac{\nu \mathbb{E} B}{r}, \\ \text{Var} \Lambda(t) &= \frac{\nu \mathbb{E} B^2}{2r}(1 - e^{-2rt}), & \text{Var} \Lambda &= \frac{\nu \mathbb{E} B^2}{2r}, \\ \text{Cov}(\Lambda(t), \Lambda(t + \delta)) &= e^{-r\delta} \text{Var} \Lambda(t). \end{aligned} \quad (2)$$

We remark that, for convenience, we throughout assume $\Lambda(0) = 0$. The results can be extended to $\Lambda(0) > 0$ in a straightforward fashion, at the cost of somewhat more cumbersome notation.

In the one-dimensional case, we denote $\beta(s) = \mathbb{E} e^{-sB}$, and in the multidimensional case, where $s = (s_1, s_2, \dots, s_d)$, for some integer $d \geq 2$, now denotes a vector, we write

$$\beta(s) = \mathbb{E} e^{-\sum_i s_i B_i}.$$

In the remainder of the paper we often use the following lemma.




Figure 1: Sample path of shot-noise process




Figure 2: A realization of arrival process corresponding to the sample path of the arrival rate process in Fig. 1

Lemma 2.1. Let $\Lambda(\cdot)$ be a shot-noise process. Let $f : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a function which is piecewise continuous in its first argument, with at most a countable number of discontinuities. Then it holds that

$$\begin{aligned} & \mathbb{E} \exp \left(\int_0^t f(u, z) \Lambda(u) \, du - s \Lambda(t) \right) \\ &= \exp \left(\nu \int_0^t \left(\beta \left(s e^{-r(t-v)} - e^{rv} \int_v^t f(u, z) e^{-ru} \, du \right) - 1 \right) \, dv \right). \end{aligned}$$

Proof. See appendix A. □

3 A single infinite-server queue

In this section we study the $M_S/G/\infty$ queue. This is a single infinite-server queue, of which the arrival process is a Cox process driven by the shot-noise process $\Lambda(\cdot)$, as defined in Section 2. First we derive exact results in Section 3.1, where we find the joint transform

of the number of customers in the system and the shot-noise rate, and derive expressions for the expected value and variance. Subsequently, in Section 3.2, we derive a functional central limit theorem for this model.

3.1 Exact analysis

We let J_i denote the service requirement of the i -th customer, where J_1, J_2, \dots are assumed to be i.i.d.; in the sequel J denotes a random variable that is equal in distribution to J_1 . Our first objective is to find the distribution of the number of customers in the system at time t , in the sequel denoted by $N(t)$. This can be found in several ways; because of the appealing underlying intuition, we here provide an argument in which we approximate the arrival rate on intervals of length Δ by a constant, and then let $\Delta \downarrow 0$.

This procedure works as follows. We let $\Lambda(t) = \Lambda(\omega, t)$ be an arbitrary sample path of the driving shot-noise process. Given $\Lambda(t)$, the number of customers that arrived in the interval $[k\Delta, (k+1)\Delta)$ and are still in the system at time t , has a Poisson distribution with parameter $\mathbb{P}(J > t - (k\Delta + \Delta U_k)) \cdot \Delta\Lambda(k\Delta) + o(\Delta)$, where U_1, U_2, \dots are i.i.d. standard uniform random variables. Summing over k yields that the number of customers in the system at time t has a Poisson distribution with parameter

$$\sum_{k=1}^{t/\Delta} \mathbb{P}(J > t - (k\Delta + \Delta U_k)) \Delta\Lambda(k\Delta) + o(\Delta),$$

which converges, as $\Delta \downarrow 0$, to

$$\int_0^t \mathbb{P}(J > t - u) \Lambda(u) du. \quad (3)$$

The argument above is not new: a similar observation was mentioned in e.g. [6], for deterministic rate functions. Since $\Lambda(\cdot)$ is actually a stochastic process, we conclude that the number of customers has a mixed Poisson distribution, with the expression in Eqn. (3) as random parameter. As a consequence, we find by conditioning on \mathcal{F}_t ,

$$\begin{aligned} \xi(t, z, s) &:= \mathbb{E} z^{N(t)} e^{-s\Lambda(t)} = \mathbb{E} \left(e^{-s\Lambda(t)} \mathbb{E} \left(z^{N(t)} \mid \mathcal{F}_t \right) \right) \\ &= \mathbb{E} \exp \left(\int_0^t (z - 1) \mathbb{P}(J > t - u) \Lambda(u) du - s\Lambda(t) \right). \end{aligned} \quad (4)$$

We have found the following result.

Theorem 3.1. *Let $\Lambda(\cdot)$ be a shot-noise process. Then*

$$\log \xi(t, z, s) = \nu \int_0^t \left(\beta \left((1 - z) e^{rv} \int_v^t \mathbb{P}(J > t - u) e^{-ru} du + s e^{-r(t-v)} \right) - 1 \right) dv. \quad (5)$$

Proof. The result follows directly from Lemma 2.1 and Eqn. (4). \square

In Thm. 3.1 we found that $N(t)$ has a Poisson distribution with the random parameter given in Eqn. (3). This leads to the following expression for the expected value

$$\mathbb{E} N(t) = \int_0^t \mathbb{E} \Lambda(u) \mathbb{P}(J > t - u) du. \quad (6)$$

In addition, by the law of total variance we find

$$\text{Var} N(t) = \text{Var} \left(\int_0^t \Lambda(u) \mathbb{P}(J > t - u) du \right) + \mathbb{E} \left(\int_0^t \Lambda(u) \mathbb{P}(J > t - u) du \right). \quad (7)$$

The latter expression we can further evaluate, using an approximation argument that resembles the one we used above. Using a Riemann sum approximation, we find

$$\begin{aligned} \text{Var} \left(\int_0^t \Lambda(u) \mathbb{P}(J > t - u) du \right) &= \lim_{\Delta \downarrow 0} \text{Var} \left(\sum_{i=0}^{t/\Delta} \Lambda(i\Delta) \mathbb{P}(J > t - i\Delta) \Delta \right) \\ &= 2 \lim_{\Delta \downarrow 0} \sum_{i=0}^{t/\Delta} \sum_{j>i}^{t/\Delta} \text{Cov}(\Lambda(i\Delta) \mathbb{P}(J > t - i\Delta) \Delta, \Lambda(j\Delta) \mathbb{P}(J > t - j\Delta) \Delta) \\ &= 2 \int_0^t \int_v^t \text{Cov}(\Lambda(u), \Lambda(v)) \mathbb{P}(J > t - u) \mathbb{P}(J > t - v) du dv. \end{aligned}$$

Assuming that $u \geq v$, we know that $\text{Cov}(\Lambda(u), \Lambda(v)) = e^{-r(u-v)} \text{Var} \Lambda(v)$ (cf. Lemma B.1 in the appendix for a proof). We thus have that (7) equals

$$2 \int_0^t \int_v^t e^{-r(u-v)} \text{Var} \Lambda(v) \mathbb{P}(J > t - u) \mathbb{P}(J > t - v) du dv + \int_0^t \mathbb{E} \Lambda(u) \mathbb{P}(J > t - u) du.$$

We can make this more explicit using the corresponding formulas in (2).

Example 3.2 (Exponential case). Consider the case in which J is exponentially distributed with mean $1/\mu$ and $\Lambda(0) = 0$. Then we can calculate the mean and variance explicitly. For $\mu \neq r$,

$$\mathbb{E} N(t) = \frac{\mathbb{E} \Lambda}{\mu} h_{r,\mu}(t),$$

where the function $h_{r,\mu}(\cdot)$ is defined by

$$t \mapsto \begin{cases} \frac{\mu(1 - e^{-rt}) - r(1 - e^{-\mu t})}{\mu - r} & \text{if } \mu \neq r, \\ 1 - e^{-rt} - r t e^{-rt} & \text{if } \mu = r. \end{cases}$$

For the variance, we thus find for $\mu \neq r$

$$\begin{aligned} \text{Var} N(t) &= \frac{\nu \mathbb{E} B^2}{2r} \frac{r^2(1 - e^{-2\mu t}) + \mu^2(1 - e^{-2rt}) + \mu r(4e^{-t(\mu+r)} - e^{-2\mu t} - e^{-2rt} - 2)}{\mu(\mu - r)^2(\mu + r)} \\ &\quad + \frac{\nu \mathbb{E} B}{r} h_{r,\mu}(t), \end{aligned}$$

and for $\mu = r$

$$\mathbb{V}\text{ar } N(t) = \frac{\nu \mathbb{E} B^2}{4r^3} \left(1 - e^{-2rt} - 2rt(1 + rt)e^{-2rt} \right) + \frac{\nu \mathbb{E} B}{r} h_{r,r}(t).$$

3.2 Asymptotic analysis

This subsection focuses on deriving a functional central limit theorem (FCLT) for the model under study, after having appropriately scaled the shot rate of the shot-noise process. In the following we assume that the service requirements are exponentially distributed with rate μ , and we point out how it can be generalized to a general distribution in Remark 3.6 below. We follow the standard approach to derive the FCLT for infinite-server queueing systems; we mimic the argumentation used in e.g. [1, 14]. As the proof has a relatively large number of standard elements, we restrict ourselves to the most important steps.

We apply a linear scaling to the shot rate of the shot-noise process, i.e. $\nu \mapsto n\nu$. It is readily checked that under this scaling, the steady-state level of the shot-noise process, as well as the steady-state number of customers in the queue, blow up by a factor n , too. It is our objective to prove that, after appropriate centering and normalization, the process recording the number of customers in the system converges to a Gaussian process.

In the n -th scaled model, the number of customers in the system at time t , denoted by $N_n(t)$, has the following (obvious) representation: with $A_n(t)$ denoting the number of arrivals in $[0, t]$, and $D_n(t)$ the number of departures,

$$N_n(t) = N_n(0) + A_n(t) - D_n(t). \quad (8)$$

Here, $A_n(t)$ corresponds to a Cox process with a shot-noise-driven rate, and therefore we have, with $\Lambda_n(s)$ the shot-noise in the scaled model at time s and $S_A(\cdot)$ a unit-rate Poisson process,

$$A_n(t) = S_A \left(\int_0^t \Lambda_n(u) du \right);$$

in line with our previous assumptions, we put $\Lambda_n(0) = 0$. For our infinite-server model the departures $D_n(t)$ can be written as, with $S_D(\cdot)$ a unit-rate Poisson process (independent of $S_A(\cdot)$),

$$D_n(t) = S_D \left(\int_0^t \mu N_n(u) du \right).$$

We start by identifying the average behavior of the process $N_n(t)$. Following the reasoning of [1], assuming that $N_n(0)/n \Rightarrow \rho(0)$, $N_n(t)/n$ converges almost surely to the solution of

$$\rho(t) = \rho(0) + \int_0^t \mathbb{E} \Lambda(u) du - \int_0^t \mu \rho(u) du. \quad (9)$$

This equation is solved by $\rho(t) = \mathbb{E} N(t)$, with $\mathbb{E} N(t)$ provided in Example 3.2. Now we move to the derivation of the FCLT. Following the approach used in [1], we proceed by studying an FCLT for the input rate process. To this end, we first define

$$\hat{\Lambda}_n(t) := \sqrt{n} \left(\frac{1}{n} \Lambda_n(t) - \mathbb{E} \Lambda(t) \right); \quad \hat{K}_n(t) := \int_0^t \hat{\Lambda}_n(u) du.$$

The following lemma states that $\hat{K}_n(\cdot)$ converges to an integrated Ornstein-Uhlenbeck (OU) process, corresponding to an OU process $\hat{\Lambda}(\cdot)$ with a speed of mean reversion equal to r , long-run equilibrium level 0, and variance $\sigma_\Lambda^2 := \nu \mathbb{E} B^2 / (2r)$. We let ' \Rightarrow ' denote weak convergence.

Lemma 3.3. *Assume that for a shot B it holds that $\mathbb{E} B, \mathbb{E} B^2 < \infty$. Then $\hat{K}_n(\cdot) \Rightarrow \hat{K}(\cdot)$ as $n \rightarrow \infty$, where*

$$\hat{K}(t) = \int_0^t \hat{\Lambda}(u) du, \tag{10}$$

in which $\hat{\Lambda}$ satisfies, with $W_1(\cdot)$ a standard Brownian motion,

$$\hat{\Lambda}(t) = \sigma_\Lambda W_1(t) - r \int_0^t \hat{\Lambda}(u) du. \tag{11}$$

Proof. This proof is standard; for instance from [2, Prop. 3], by putting the λ_d in that paper to zero, it follows that $\hat{\Lambda}_n(\cdot) \Rightarrow \hat{\Lambda}(\cdot)$. This implies $\hat{K}_n(\cdot) \Rightarrow \hat{K}(\cdot)$, using (10) together with the continuous mapping theorem. \square

Interestingly, the above result entails that the rate process displays mean-reverting behavior. This also holds for infinite-server queues in general. In other words, the number of customer process, in the queueing system we are studying, can be considered as the composition of two mean-reverting processes. We make this more precise in the following.

From now on we consider the following centered and normalized version of the number of customers in the system:

$$\hat{N}_n(t) := \sqrt{n} \left(\frac{1}{n} N_n(t) - \rho(t) \right).$$

We assume that $\hat{N}_n(0) \Rightarrow \hat{N}(0)$ as $n \rightarrow \infty$. To prove the FCLT, we rewrite $\hat{N}_n(t)$ in a convenient form. Mimicking the steps performed in [1] or [14], with $\bar{S}_A(t) := S_A(t) - t$, $\bar{S}_D(t) := S_D(t) - t$,

$$R_n(t) := \bar{S}_A \left(\int_0^t \Lambda_n(u) du \right) - \bar{S}_D \left(\mu \int_0^t N_n(u) du \right),$$

and using the relation (9), we eventually obtain

$$\hat{N}_n(t) = \hat{N}_n(0) + \frac{R_n(t)}{\sqrt{n}} + \hat{K}_n(t) - \mu \int_0^t \hat{N}_n(u) du.$$

Our next goal is to apply the martingale FCLT to the martingales $R_n(t)/\sqrt{n}$; see for background on the martingale FCLT for instance [7] and [17]. The quadratic variation equals

$$\left[\frac{R_n}{\sqrt{n}} \right]_t = \frac{1}{n} \left(S_A \left(\int_0^t \Lambda_n(u) du \right) + S_D \left(\mu \int_0^t N_n(u) du \right) \right),$$

which converges to $\int_0^t \mathbb{E} \Lambda(u) du + \mu \int_0^t \rho(u) du$. Appealing to the martingale FCLT, the following FCLT is obtained.

Theorem 3.4. *The centered and normalized version of the number of customers in the queue satisfies an FCLT: $\hat{N}_n(\cdot) \Rightarrow \hat{N}(\cdot)$ as $n \rightarrow \infty$, where $\hat{N}(t)$ solves the stochastic integral equation*

$$\hat{N}(t) = \hat{N}(0) + \int_0^t \sqrt{\mathbb{E} \Lambda(u) + \mu \rho(u)} dW_2(u) + \hat{K}(t) - \mu \int_0^t \hat{N}(u) du,$$

with $W_2(\cdot)$ a standard Brownian motion that is independent of the Brownian motion $W_1(\cdot)$ we introduced in the definition of $\hat{K}(\cdot)$.

Remark 3.5. In passing, we have proven that the arrival process as such obeys an FCLT. With

$$\hat{A}_n(t) := \sqrt{n} \left(\frac{1}{n} A_n(t) - \int_0^t \mathbb{E} \Lambda(u) du \right),$$

we find that $\hat{A}_n(t) \Rightarrow \hat{A}(t)$ as $n \rightarrow \infty$, where

$$\hat{A}(t) := \int_0^t \sqrt{\mathbb{E} \Lambda(u) + \mu \rho(u)} dW_2(u) + \hat{K}(t) = \int_0^t \sqrt{2\mu \rho(u) + \rho'(u)} dW_2(u) + \hat{K}(t);$$

the last equality follows from the fact that $\rho(\cdot)$ satisfies (9).

Remark 3.6. The FCLT can be extended to non-exponential service requirements, by making use of [15, Thm. 3.2]. Their approach relies on two assumptions:

- The arrival process should satisfy an FCLT;
- The service times are i.i.d. non-negative random variables with a general c.d.f. independent of the arrival process.

As noted in Remark 3.5, the first assumption is satisfied for the model in this paper. The second assumption holds as well. In the non-exponential case the results are less clean; in general, the limiting process can be expressed in terms of a Kiefer process, cf. e.g. [4].

4 Networks

Now that the reader is familiar with the one-dimensional setting, we extend this to networks. In this section, we mainly study feedforward networks in which each node corresponds to an infinite-server queue. Feedforward networks are defined as follows.




Figure 3: Since the customers are not interfering with each other, the network on the left is equivalent to the graph on the right. Node $3'$ is a copy of node 3: it works at the same speed and induces the same service requirements.

Definition 4.1 (feedforward network). Let $G = (V, E)$ be a directed graph with nodes V and edges E . The nodes represent infinite-server queues and the directed edges between the facilities demonstrate how customers move through the system. We suppose that there are no cycles in G , i.e. there is no sequence of nodes, starting and ending at the same node, with each two consecutive nodes adjacent to each other in the graph, consistent with the orientation of the edges.

We focus on feedforward networks to keep the notation manageable. In Thm. 4.6, we derive the transform of the numbers of customers in all nodes, jointly with the shot-noise process(es) for feedforward networks. Nonetheless, we provide Example 4.5, to show that analysis is in fact possible if there is a loop, but at the expense of more involved calculations.

Since all nodes represent infinite-server queues, one can see that whenever a node has multiple input streams, it is equivalent to multiple infinite-server queues that work independently from each other, but have the same service speed and induce the same service requirement for arriving jobs. Consider Fig. 3 for an illustration. The reason why this holds is that different customer streams move independently through the system, without creating waiting times for others. Therefore, merging streams do not increase the complexity of our network. The same holds for ‘splits’ in customer streams. By this we mean that after customers finished their service in a server, they move to server i with probability q_i (with $\sum_i q_i = 1$). Then, one can simply sample the entire path that the customer will take through the system, at the arrival instance at its first server.

If one recognizes the above, then all feedforward networks reduce to parallel tandem systems in which the first node in each tandem system is fed by external input. The procedure to decompose a network into parallel tandems, is to find all paths in the network that start from a node in which there is external input and end in a node of which the departures leave the system. Each of these paths will subsequently be considered as a tandem queue, which are then set in parallel.

To build up to the main result, we first study tandem systems in Section 4.1. Subsequently, we put the tandem systems in parallel in Section 4.2 and finally we present the main theorem and some implications in Section 4.3.

4.1 Tandem systems

As announced, we proceed by studying tandem systems. In Section 4.2 below, we study d parallel tandem systems, where $i = 1, \dots, d$. In this subsection we consider the i -th of these tandem systems.

Suppose that tandem i has S_i service facilities and the input process at the first node is Poisson, with a shot-noise arrival rate $\Lambda_i(\cdot)$. We assume that customers enter node $i1$. When they finish service, they enter node $i2$, etc., until they enter node iS_i after which they leave the system. We use ij as a subscript referring to node j in tandem system i and we refer to the node as node ij . Hence $N_{ij}(t)$ and J_{ij} denote the number of customers in node ij at time t , and a copy of a service requirement, respectively, where $j = 1, \dots, S_i$. Fix some time $t > 0$. Again we derive results by splitting time into intervals of length Δ . Denote $M_{ij}(k, \Delta)$ for the number of customers present in node ij at time t that have entered node $i1$ between time $k\Delta$ and $(k+1)\Delta$; as we keep t fixed we suppress it in our notation. Because customers are not interfering with each other in the infinite-server realm, we can decompose the transform of interest:

$$\mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} \right) = \lim_{\Delta \downarrow 0} \prod_{k=0}^{t/\Delta} \mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{M_{ij}(k, \Delta)} \right). \quad (12)$$

Supposing that the arrival rate is a deterministic function of time $\lambda_i(\cdot)$, by conditioning on the number of arrivals in the k -th interval,

$$\begin{aligned} \mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{M_{ij}(k, \Delta)} \right) &= \sum_{m=0}^{\infty} e^{-\lambda_i(k\Delta)\Delta} \frac{(\lambda_i(k\Delta)\Delta)^m}{m!} (f_i(k\Delta, z))^m \\ &= \exp \left(\Delta \lambda_i(k\Delta) (f_i(k\Delta, z) - 1) \right), \end{aligned}$$

where

$$f_i(u, z) := p_i(u) + \sum_{j=1}^{S_i} z_{ij} p_{ij}(u), \quad (13)$$

in which $p_i(u)$ ($p_{ij}(u)$, respectively) denotes the probability that the job that entered tandem i at time u has already left the tandem (is in node j , respectively) at time t . Note that

$$p_i(u) = \mathbb{P} \left(\sum_{\ell=1}^{S_i} J_{i\ell} < t - u \right), \quad p_{ij}(u) = \mathbb{P} \left(\sum_{\ell=1}^{j-1} J_{i\ell} < t - u, \sum_{\ell=1}^j J_{i\ell} > t - u \right). \quad (14)$$

Recognizing a Riemann sum and letting $\Delta \downarrow 0$, we conclude that Eqn. (12) takes the following form:

$$\mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} \right) = \exp \left(\int_0^t \lambda_i(u) (f_i(u, z) - 1) du \right).$$

In case of a stochastic rate process $\Lambda_i(\cdot)$, we obtain

$$\mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} \middle| \Lambda_i(\cdot) \right) = \exp \left(\int_0^t \Lambda_i(u) (f_i(u, z) - 1) du \right).$$

Therefore it holds that

$$\begin{aligned} \mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} e^{-s\Lambda_i(t)} \right) &= \mathbb{E} \left(\mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} e^{-s\Lambda_i(t)} \middle| \Lambda_i(\cdot) \right) \right) \\ &= \mathbb{E} \left(e^{-s\Lambda_i(t)} \mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} \middle| \Lambda_i(\cdot) \right) \right), \end{aligned}$$

and we consequently find

$$\mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} e^{-s\Lambda_i(t)} \right) = \mathbb{E} \exp \left(\int_0^t \Lambda_i(u) (f_i(u, z) - 1) du - s\Lambda_i(t) \right). \quad (15)$$

4.2 Parallel (tandem) systems

Now that the tandem case has been analyzed, the next step is to put the tandem systems as described in Section 4.1 in parallel. We assume that there are d parallel tandems. There are different ways in which dependence between the parallel systems can be created. Two relevant models are listed below, and illustrated in Fig. 4.

- (M1)** Let $\Lambda \equiv \Lambda(\cdot)$ be a d -dimensional shot-noise process $(\Lambda_1, \dots, \Lambda_d)$ where the shots in all Λ_i occur simultaneously (the shot distributions and decay rates may be different). The process Λ_i , for $i = 1, \dots, d$, corresponds to the arrival rate of tandem system i . Each tandem system has an arrival process, in which the Cox processes are independent given their shot-noise arrival rates.
- (M2)** Let $\Lambda \equiv \Lambda(\cdot)$ be the shot-noise rate of a Cox process. The corresponding Poisson process generates *simultaneous* arrivals in all tandems.




Figure 4: Model (M1) is illustrated on the left, and Model (M2) is illustrated on the right. The rectangles represent tandem systems, which consist of an arbitrary number of nodes in series.

Remark 4.2. The model in which there is essentially one shot-noise process that generates arrivals for all queues independently, is a special case of Model (M1). This can be seen by setting all components of $\Lambda = (\Lambda_1, \dots, \Lambda_d)$ equal, by letting the shots and decay rate be identical.

In Model (M1), correlation between the shot-noise arrival rates induces correlation between the numbers of customers in the different queues. In Model (M2), correlation clearly appears because all tandem systems have the same input process. Of course, the tandem systems will not behave identically because the customers may have different service requirements. In short, correlation across different tandems in Model (M1) is due to linked arrival rates, and correlation in Model (M2) is due to simultaneous arrival epochs. We feel that both versions are relevant, depending on the application, and hence we analyze both.

Analysis of (M1) — Suppose that the dependency is of the type as in (M1). This means that $\Lambda = (\Lambda_1, \dots, \Lambda_d)$ for which the disturbances in $\Lambda_1, \dots, \Lambda_d$ arrive simultaneously. Recall the definition of f_i as stated in Eqn. (13). It holds that

$$\begin{aligned}
\mathbb{E} \left(\prod_{i=1}^d e^{-s_i \Lambda_i(t)} \prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} \right) &= \mathbb{E} \left(\prod_{i=1}^d \mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} e^{-s_i \Lambda_i(t)} \mid \Lambda_i(\cdot) \right) \right) \\
&= \mathbb{E} \left(\prod_{i=1}^d \left(e^{-s_i \Lambda_i(t)} \mathbb{E} \left(\prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} \mid \Lambda_i(\cdot) \right) \right) \right) \\
&= \mathbb{E} \exp \left(\sum_{i=1}^d \left(\int_0^t \Lambda_i(u) (f_i(u, z) - 1) du - s_i \Lambda_i(t) \right) \right), \quad (16)
\end{aligned}$$

where the last equality holds due to (15).

Analysis of (M2) — Now suppose that the dependency in this model is of type (M2), i.e., there is one shot-noise process that generates simultaneous arrivals in the parallel tandem systems.

First we assume a deterministic arrival rate function $\lambda(\cdot)$. Let $M_{ij}(k, \Delta)$ be the number of jobs present in tandem system i at node j at time t that have arrived in the system between $k\Delta$ and $(k+1)\Delta$. Note that

$$\mathbb{E} \left(\prod_{i=1}^d \prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} \right) = \lim_{\Delta \downarrow 0} \prod_{k=1}^{t/\Delta} \mathbb{E} \left(\prod_{i=1}^d \prod_{j=1}^{S_i} z_{ij}^{M_{ij}(k, \Delta)} \right).$$

To further evaluate the right hand side of the previous display, we observe that we can write

$$\mathbb{E} \left(\prod_{i=1}^d \prod_{j=1}^{S_i} z_{ij}^{M_{ij}(k, \Delta)} \right) = \sum_{m=0}^{\infty} e^{-\lambda(k\Delta)\Delta} \frac{\lambda(k\Delta)\Delta^m}{m!} (f(k\Delta, z))^m = e^{\Delta\lambda(k\Delta)(f(k\Delta, z)-1)},$$

where

$$f(u, z) := \sum_{j=1}^d \sum_{\ell_j=1}^{S_{j+1}} p_{\ell_1, \dots, \ell_d} \prod_{i=1}^d z_{i\ell_i}; \quad (17)$$

in this definition $p_{\ell_1, \dots, \ell_d} \equiv p_{\ell_1, \dots, \ell_d}(u)$ equals the probability that a job that arrived at time u in tandem i is in node ℓ_i at time t (cf. Eqn. (14)). The situation that $\ell_i = S_i + 1$ means that the job left the tandem system; we define $z_{i, S_i+1} = 1$.

In a similar fashion as before, we conclude that

$$\mathbb{E} \left(\prod_{i=1}^d \prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} e^{-s\Lambda(t)} \right) = \mathbb{E} \exp \left(\int_0^t \Lambda(u)(f(u, z) - 1) du - s\Lambda(t) \right), \quad (18)$$

with f defined in Eqn. (17).

Example 4.3 (Two-node parallel system). In case we would be analysing a parallel system of simply two infinite-server queues, then $f(u, z)$ simplifies considerably to

$$f(u, z_{11}, z_{21}) = \sum_{\ell_1=1}^2 \sum_{\ell_2=1}^2 z_{1\ell_1} z_{2\ell_2} p_{\ell_1, \ell_2} = z_{11} z_{21} p_{11} + z_{21} p_{21} + z_{11} p_{12} + p_{22}.$$

Remark 4.4 (Routing). Consider a feedforward network with routing. As argued in the beginning of this section, the network can be decomposed as a parallel tandem system. In case there is splitting at some point, then one decomposes the network as a parallel system, in which each tandem i receives the customer with probability q_i , such that $\sum q_i = 1$. This can be incorporated simply by adjusting the probabilities contained in f_i in Eqn. (16), which are given in Eqn. (14), so that they include the event that the customer joined the tandem under consideration. For instance, the expression for $p_i(u)$ in the left equation in (14) would become

$$\mathbb{P} \left(Q = i, \sum_{\ell=1}^{S_i} J_{i\ell} < t - u \right),$$

where Q is a random variable with a generalized Bernoulli (also called ‘categorical’) distribution, where

$$\mathbb{P}(\text{customer is assigned to tandem } i) = \mathbb{P}(Q = i) = q_i, \quad \text{for } i = 1, \dots, d,$$

with $\sum q_i = 1$; the right equation in (14) is adjusted similarly. Other than that, the analysis is the same for the case of splits.

Remark 4.5 (Networks with loops). So far we only considered feedforward networks. Networks with loops can be analyzed as well, but the notation becomes quite cumbersome. To show the method in which networks with loops and routing can be analyzed, we consider a specific example. Suppose that arrivals enter node one, after which they enter node two. After they have been served in node two, they go back to node one with probability η , or leave the system with probability $1 - \eta$. Note that this network has a loop. In this case, with similar techniques as before, we can find

$$\mathbb{E} z_1^{N_1(t)} z_2^{N_2(t)} = \exp \left(\int_0^t \Lambda(u) (f(u, z_1, z_2) - 1) du \right),$$

with

$$f(u, z_1, z_2) = \mathbb{P}(\text{job}(u) \text{ left system}) + \sum_{i=1}^2 z_i \mathbb{P}(\text{job}(u) \text{ is in node } i),$$

in which $\text{job}(u)$ is the job that arrived at time u and we are examining the system at time t . Now, if we denote service times in the j -th node by $J^{(j)}$, then, at a specific time t ,

$$\mathbb{P}(\text{job}(u) \text{ left system}) = \sum_{k=0}^{\infty} \mathbb{P} \left(\sum_{i=1}^{k+1} (J_i^{(1)} + J_i^{(2)}) \leq t - u \right) \eta^k (1 - \eta).$$

Analogously, $\mathbb{P}(\text{job}(u) \text{ is in node 1})$ equals, by conditioning on the job having taken k loops,

$$\sum_{k=0}^{\infty} \eta^k \mathbb{P} \left(J_{k+1}^{(1)} + \sum_{i=1}^k (J_i^{(1)} + J_i^{(2)}) > t - u, \sum_{i=1}^k (J_i^{(1)} + J_i^{(2)}) \leq t - u \right);$$

likewise, $\mathbb{P}(\text{job}(u) \text{ is in node 2})$ equals

$$\sum_{k=0}^{\infty} \eta^k \mathbb{P} \left(\sum_{i=1}^{k+1} (J_i^{(1)} + J_i^{(2)}) > t - u, J_{k+1}^{(1)} + \sum_{i=1}^k (J_i^{(1)} + J_i^{(2)}) \leq t - u \right).$$

For example, in case all $J_i^{(j)}$ are independent and exponentially distributed with mean $1/\mu$, we can calculate those probabilities explicitly. Indeed, if we denote Y for a Poisson

process with rate μ , then e.g.,

$$\begin{aligned} \mathbb{P} \left(\sum_{i=1}^{k+1} (J_i^{(1)} + J_i^{(2)}) > t - u, J_{k+1}^{(1)} + \sum_{i=1}^k (J_i^{(1)} + J_i^{(2)}) \leq t - u \right) &= \mathbb{P}(Y(t - u) = 2k + 1) \\ &= e^{-\mu(t-u)} \frac{(\mu(t-u))^{2k+1}}{(2k+1)!} \end{aligned}$$

and thus

$$\mathbb{P}(\text{job}(u) \text{ is in node 2}) = \sum_{m=0}^{\infty} \eta^m e^{-\mu(t-u)} \frac{(\mu(t-u))^{2m+1}}{(2m+1)!}.$$

A similar calculation can be done for the probability that the job is in node one. Recalling that a sum of exponentials has a Gamma distribution, we can write

$$\begin{aligned} f(u, z_1, z_2) &= z_1 \sum_{m=0}^{\infty} \eta^m e^{-\mu(t-u)} \frac{(\mu(t-u))^{2m}}{(2m)!} + z_2 \sum_{m=0}^{\infty} \eta^m e^{-\mu(t-u)} \frac{(\mu(t-u))^{2m+1}}{(2m+1)!} \\ &\quad + \sum_{m=0}^{\infty} \eta^m (1 - \eta) F_{\Gamma(2m+2, \mu)}(t - u) \\ &= z_1 e^{-\mu(t-u)} \cosh(\mu\sqrt{\eta}(t-u)) + z_2 \frac{e^{-\mu(t-u)}}{\sqrt{\eta}} \sinh(\mu\sqrt{\eta}(t-u)) \\ &\quad + (1 - \eta) \sum_{m=0}^{\infty} \eta^m F_{\Gamma(2m+2, \mu)}(t - u), \end{aligned}$$

where $F_{\Gamma(2m+2, \mu)}$ denotes the distribution function of a Γ -distributed random variable with rate μ and shape parameter $2m + 2$.

4.3 Main result

In this subsection we summarize and conclude with the following main result. Recall Definition 4.1 of a feedforward network. In the beginning of Section 4 we argued that we can decompose a feedforward network into parallel tandems. In Section 4.2 we studied exactly those systems, leading up to the following results.

Theorem 4.6. *Suppose we have a feedforward network of infinite-server queues, where the input process is a Poisson process with shot-noise arrival rate. Then the network can be decomposed into parallel tandem systems. Under assumption (M1), it holds that*

$$\begin{aligned} \mathbb{E} \left(\prod_{i=1}^d \prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} e^{-s\Lambda_i(t)} \right) &= \mathbb{E} \exp \left(\sum_{i=1}^d \left(\int_0^t \Lambda_i(u) (f_i(u, z) - 1) du - s_i \Lambda_i(t) \right) \right) \\ &= \exp \left(\nu \int_0^t (\beta(g(s, v)) - 1) dv \right), \end{aligned}$$

with $f_i(\cdot, \cdot)$ as defined in Eqn. (13) and where $g(s, v)$ is a vector-valued function in which component i is given by

$$s_i e^{-r_i(t-v)} - e^{r_i v} \int_v^t (f_i(u, z) - 1) e^{-r_i u} du.$$

Furthermore, under assumption (M2),

$$\begin{aligned} \mathbb{E} \left(\prod_{i=1}^d \prod_{j=1}^{S_i} z_{ij}^{N_{ij}(t)} e^{-s\Lambda(t)} \right) &= \mathbb{E} \exp \left(\int_0^t \Lambda(u) (f(u, z) - 1) du - s\Lambda(t) \right) \\ &= \exp \left(\nu \int_0^t \left(\beta \left(s e^{-r(t-v)} - e^{r v} \int_v^t (f(u, z) - 1) e^{-r u} du \right) - 1 \right) dv \right), \end{aligned}$$

with $f(\cdot, \cdot)$ as defined in Eqn. (17).

Proof. These are Eqns. (16) and (18) to which we applied Lemma 2.1. \square

Next we calculate covariances between nodes in tandem and between nodes in parallel.

Covariance in Tandem System — Suppose that we have a tandem system consisting of two nodes and we want to analyze the covariance between the numbers of customers in both nodes. Dropping the index of the tandem system, we denote $N_1(\cdot)$ and $N_2(\cdot)$ for the numbers of customers in node 1 and 2, respectively. From Eqn. (16),

$$\mathbb{E} N_2(t) = \int_0^t \mathbb{P}(J_1 < t - u, J_1 + J_2 > t - u) \mathbb{E} \Lambda(u) du$$

and

$$\mathbb{E} N_1(t) N_2(t) = \mathbb{E} \left(\int_0^t \mathbb{P}(J_1 < t - u, J_1 + J_2 > t - u) \Lambda(u) du \int_0^t \mathbb{P}(J_1 > t - v) \Lambda(v) dv \right)$$

so that

$$\begin{aligned} \text{Cov}(N_1(t), N_2(t)) &= \text{Cov} \left(\int_0^t \mathbb{P}(J_1 < t - u, J_1 + J_2 > t - u) \Lambda(u) du, \int_0^t \mathbb{P}(J_1 > t - v) \Lambda(v) dv \right) \\ &= 2 \int_0^t \int_v^t \mathbb{P}(J_1 < t - u, J_1 + J_2 > t - u) \mathbb{P}(J_1 > t - v) \text{Cov}(\Lambda(u), \Lambda(v)) du dv \\ &= 2 \int_0^t \int_v^t \mathbb{P}(J_1 < t - u, J_1 + J_2 > t - u) \mathbb{P}(J_1 > t - v) e^{-r(u-v)} \text{Var} \Lambda(u) du dv, \end{aligned}$$

cf. Lemma B.1 in Appendix B for the last equality.

Covariance parallel (M1) — Suppose that we have a parallel system consisting of two nodes only. By making use of Eqn. (16), we find

$$\mathbb{E} N_1(t)N_2(t) = \mathbb{E} \left(\int_0^t \Lambda_1(u) \mathbb{P}(J_1 > t - u) du \int_0^t \Lambda_2(v) \mathbb{P}(J_2 > t - v) dv \right).$$

This implies

$$\begin{aligned} \text{Cov}(N_1(t), N_2(t)) &= \text{Cov} \left(\int_0^t \Lambda_1(u) \mathbb{P}(J_1 > t - u) du, \int_0^t \Lambda_2(v) \mathbb{P}(J_2 > t - v) dv \right) \\ &= 2 \int_0^t \int_v^t \text{Cov}(\Lambda_1(u), \Lambda_2(v)) \mathbb{P}(J_1 > t - u) \mathbb{P}(J_2 > t - v) du dv \\ &= 2 \int_0^t \int_v^t \frac{\nu \mathbb{E} B_{11} B_{12}}{r_1 + r_2} \left(1 - e^{-(r_1+r_2)u} \right) e^{-r_2(u-v)} \mathbb{P}(J_1 > t - u) \mathbb{P}(J_2 > t - v) du dv \end{aligned}$$

where we made use of the fact that, for $u \geq v$,

$$\text{Cov}(\Lambda_1(u), \Lambda_2(v)) = \text{Cov}(\Lambda_1(u), \Lambda_2(u)) e^{-r_2(u-v)} = \frac{\nu \mathbb{E} B_{11} B_{12}}{r_1 + r_2} \left(1 - e^{-(r_1+r_2)u} \right) e^{-r_2(u-v)},$$

cf. Appendix B.

Covariance parallel (M2) — From Eqn. (18) we derive

$$\begin{aligned} \mathbb{E} N_1(t)N_2(t) &= \mathbb{E} \left(\int_0^t \Lambda(u) \mathbb{P}(J_1 > t - u, J_2 > t - u) du \right) \\ &\quad + \mathbb{E} \left(\int_0^t \Lambda(u) \mathbb{P}(J_1 > t - u) du \int_0^t \Lambda(u) \mathbb{P}(J_2 > t - u) du \right). \end{aligned}$$

This implies

$$\begin{aligned} \text{Cov}(N_1(t), N_2(t)) &= \text{Cov} \left(\int_0^t \Lambda(u) \mathbb{P}(J_1 > t - u) du, \int_0^t \Lambda(u) \mathbb{P}(J_2 > t - u) du \right) \\ &\quad + \int_0^t \mathbb{E} \Lambda(u) \mathbb{P}(J_1 > t - u, J_2 > t - u) du. \end{aligned}$$

The following proposition compares the correlations present under assumptions (M1) and (M2). In the proposition we refer to the number of customers in queue j in model i at time t as $N_j^{(i)}(t)$. We find the anticipated result, that the correlation under assumption (M2) is stronger than under assumption (M1).

Proposition 4.7. *Let $\Lambda(\cdot)$ be the shot-noise process that generates simultaneous arrivals in both queues and let $\Lambda_1(\cdot), \Lambda_2(\cdot)$ be processes that have simultaneous jumps and generate arrivals in both queues independently. Suppose that $\Lambda_1(t) \stackrel{d}{=} \Lambda_2(t) \stackrel{d}{=} \Lambda(t)$, for $t \geq 0$. Then, for any $t \geq 0$,*

$$\text{Corr}(N_1^{(1)}(t), N_2^{(1)}(t)) \leq \text{Corr}(N_1^{(2)}(t), N_2^{(2)}(t)).$$

Proof. Because of the assumption $\Lambda_1(t) \stackrel{d}{=} \Lambda_2(t) \stackrel{d}{=} \Lambda(t)$, we have that, for all combinations $i, j \in \{1, 2\}$, the $N_i^{(j)}(t)$ are equal in distribution. Therefore it is sufficient to show that

$$\mathbb{Cov}(N_1^{(1)}(t), N_2^{(1)}(t)) \leq \mathbb{Cov}(N_1^{(2)}(t), N_2^{(2)}(t)).$$

The expressions for the covariances, which are derived earlier in this section, imply that

$$\mathbb{Cov}(N_1^{(2)}(t), N_2^{(2)}(t)) - \mathbb{Cov}(N_1^{(1)}(t), N_2^{(1)}(t)) = \mathbb{E} \int_0^t \Lambda(u) \mathbb{P}(J_1 > t - u, J_2 > t - u) du,$$

which is non-negative, as desired. \square

5 Concluding remarks

We have considered networks of infinite-server queues with shot-noise-driven Coxian input processes. For the single queue, we found explicit expressions for the Laplace transform of the joint distribution of the number of customers and the driving shot-noise arrival rate, as well as a functional central limit theorem of the number of customers in the system under a particular scaling. The results were then extended to a network context: we derived an expression for the joint transform of the numbers of customers in the individual queues, jointly with the values of the driving shot-noise processes.

We included the functional central limit theorem for the single queue, but it is anticipated that a similar setup carries over to the network context, albeit at the expense of considerably more involved notation. Our future research will include the study of the departure process of a single queue; the output stream should remain Coxian, but of another type than the input process.

Acknowledgements

The authors thank Marijn Jansen for insightful discussions about the functional central limit theorem in this paper. The research for this paper is partly funded by the NWO Gravitation Project NETWORKS, Grant Number 024.002.003. The research of Onno Boxma was also partly funded by the Belgian Government, via the IAP Bestcom Project.

Appendices

A Proof of Lemma 2.1

There are various ways to prove this result; we here include a procedure that intensively relies on the probabilistic properties of the shot-noise process involved. Observe that,

recognizing a Riemann sum,

$$\int_0^t f(u, z) \Lambda(u) \, du = \lim_{\Delta \downarrow 0} \Delta \sum_{k=1}^{t/\Delta} f(k\Delta, z) \Lambda(k\Delta). \quad (19)$$

With $P_B(t)$ a Poisson process with rate ν and the U_i i.i.d. samples from a uniform distribution on $[0, 1]$, it holds that

$$\Lambda(k\Delta) = \sum_{\ell=1}^k \sum_{i=P_B((\ell-1)\Delta)+1}^{P_B(\ell\Delta)} B_i e^{-r\Delta U_i} e^{-r(k-\ell)\Delta}.$$

We thus obtain that the expression in (19) equals (where the equality follows by interchanging the order of the summations)

$$\begin{aligned} & \lim_{\Delta \downarrow 0} \Delta \sum_{k=1}^{t/\Delta} f(k\Delta, z) \sum_{\ell=1}^k \sum_{i=P_B((\ell-1)\Delta)+1}^{P_B(\ell\Delta)} B_i e^{-r\Delta U_i} e^{-r(k-\ell)\Delta} \\ &= \lim_{\Delta \downarrow 0} \Delta \sum_{\ell=1}^{t/\Delta} \sum_{i=P_B((\ell-1)\Delta)+1}^{P_B(\ell\Delta)} B_i e^{-r\Delta U_i} \sum_{k=\ell}^{t/\Delta} f(k\Delta, z) e^{-r(k-\ell)\Delta}, \end{aligned}$$

which behaves as

$$\lim_{\Delta \downarrow 0} \sum_{\ell=1}^{t/\Delta} e^{r\ell\Delta} \int_{\ell\Delta}^t f(z, u) e^{-ru} \, du \sum_{i=P_B((\ell-1)\Delta)+1}^{P(\ell\Delta)} B_i e^{-r\Delta U_i}.$$

Furthermore, we have the representation

$$\Lambda(t) = \lim_{\Delta \downarrow 0} \sum_{\ell=1}^{t/\Delta} \sum_{i=P_B((\ell-1)\Delta)+1}^{P_B(\ell\Delta)} B_i e^{-r(t-\ell\Delta)}.$$

We conclude that $\xi(t, z, s)$ (cf. Eqn. (4)) equals

$$\lim_{\Delta \downarrow 0} \mathbb{E} \exp \left(\sum_{\ell=1}^{t/\Delta} \sum_{i=P_B((\ell-1)\Delta)+1}^{P_B(\ell\Delta)} B_i \left(e^{-r\Delta U_i} e^{r\ell\Delta} \int_{\ell\Delta}^t f(u, z) e^{-ru} \, du - s e^{-r(t-\ell\Delta)} \right) \right). \quad (20)$$

Conditioning on the values of $P_B(\ell\Delta) - P_B((\ell - 1)\Delta)$, for $\ell = 1, \dots, t/\Delta$, and using that the B_i are i.i.d., we find that the expression in Eqn. (20) equals

$$\begin{aligned}
&= \lim_{\Delta \downarrow 0} \prod_{\ell=1}^{t/\Delta} e^{-\nu\Delta} \sum_{k_\ell=0}^{\infty} \frac{(\nu\Delta)^{k_\ell}}{k_\ell!} \left(\mathbb{E} \exp \left(B_1 \left(e^{-r\Delta U_i} e^{r\ell\Delta} \int_{\ell\Delta}^t f(u, z) e^{-ru} du - se^{-r(t-\ell\Delta)} \right) \right) \right)^{k_\ell} \\
&= \lim_{\Delta \downarrow 0} \prod_{\ell=1}^{t/\Delta} e^{-\nu\Delta} \exp \left(\nu\Delta \mathbb{E} \exp \left(B_1 \left(e^{-r\Delta U_i} e^{r\ell\Delta} \int_{\ell\Delta}^t f(u, z) e^{-ru} du - se^{-r(t-\ell\Delta)} \right) \right) \right) \\
&= \lim_{\Delta \downarrow 0} \exp \left(\sum_{\ell=1}^{t/\Delta} \left(\nu\Delta \mathbb{E} \exp \left(B_1 \left(e^{-r\Delta U_i} e^{r\ell\Delta} \int_{\ell\Delta}^t f(u, z) e^{-ru} du - se^{-r(t-\ell\Delta)} \right) \right) - \nu\Delta \right) \right),
\end{aligned}$$

which can be written as

$$\lim_{\Delta \downarrow 0} \exp \left(\nu\Delta \sum_{\ell=1}^{t/\Delta} \left(\beta \left(se^{-r(t-\ell\Delta)} - e^{-r\Delta U_i} e^{r\ell\Delta} \int_{\ell\Delta}^t f(u, z) e^{-ru} du \right) - 1 \right) \right).$$

By continuity of the exponent and the definition of the Riemann integral, this equals

$$\exp \left(\nu \int_0^t \left(\beta \left(se^{-r(t-v)} - e^{rv} \int_v^t f(u, z) e^{-ru} du \right) - 1 \right) dv \right).$$

B (Proof of) Lemma B.1

Lemma B.1. *Let $\Lambda_1(\cdot), \Lambda_2(\cdot)$ be shot-noise processes of which the jumps occur simultaneously according to a Poisson arrival process $P_B(\cdot)$ with rate ν . Let the decay be exponential with rate r_1, r_2 , respectively. Then it holds that, for $\delta > 0$,*

$$\text{Cov}(\Lambda_1(t), \Lambda_2(t + \delta)) = e^{-r_2\delta} \text{Cov}(\Lambda_1(t), \Lambda_2(t)),$$

which, in case $\Lambda_1 = \Lambda_2$, reduces to

$$\text{Cov}(\Lambda_i(t), \Lambda_i(t + \delta)) = e^{-r_i\delta} \text{Var} \Lambda_i(t), \quad \text{for } i = 1, 2,$$

corresponding to [16, p. 394].

Proof. Let $\mathcal{E}_{t,\delta}(n)$ be the event that $P_B(t + \delta) - P_B(t) = n$. By conditioning on the number of arrivals in the interval $(t, t + \delta]$, we find

$$\begin{aligned}
\mathbb{E} \Lambda_1(t) \Lambda_2(t + \delta) &= \sum_{n=0}^{\infty} \mathbb{E}(\Lambda_1(t) \Lambda_2(t + \delta) | \mathcal{E}_{t,\delta}(n)) \mathbb{P}(\mathcal{E}_{t,\delta}(n)) \\
&= \sum_{n=0}^{\infty} \mathbb{E}(\Lambda_1(t) \Lambda_2(t + \delta) | \mathcal{E}_{t,\delta}(n)) \frac{(\delta\nu)^n}{n!} e^{-\nu\delta}.
\end{aligned}$$

We proceed by rewriting the conditional expectation as

$$\mathbb{E}(\Lambda_1(t)\Lambda_2(t+\delta) \mid \mathcal{E}_{t,\delta}(n)) = \frac{1}{\delta^n} \int_t^{t+\delta} \dots \int_t^{t+\delta} \mathbb{E}(\Lambda_1(t)\Lambda_2(t+\delta) \mid \mathcal{F}_{t_1,\dots,t_n,\delta}(n)) dt_1 \dots dt_n,$$

denoting by $\mathcal{F}_{t_1,\dots,t_n,\delta}(n)$ the event $\mathcal{E}_{t,\delta}(n)$ and the arrival epochs are t_1, \dots, t_n . Note that we have, conditional on $\mathcal{F}_{t_1,\dots,t_n,\delta}(n)$, the distributional equality

$$\Lambda_2(t+\delta) = \Lambda_2(t)e^{-r_2\delta} + \sum_{i=1}^n B_{i2}e^{-r_2(t+\delta-t_i)}, \quad (21)$$

and consequently

$$\mathbb{E}(\Lambda_1(t)\Lambda_2(t+\delta) \mid \mathcal{F}_{t_1,\dots,t_n,\delta}(n)) = \mathbb{E}\Lambda_1(t)\Lambda_2(t)e^{-r_2\delta} + \mathbb{E}\Lambda_1(t) \sum_{i=1}^n \mathbb{E}B_{i2}e^{-r_2(t+\delta-t_i)}. \quad (22)$$

Note that for all $i = 1, \dots, n$ we have

$$\int_t^{t+\delta} \dots \int_t^{t+\delta} \int_t^{t+\delta} e^{-r_2(t+\delta-t_i)} dt_1 dt_2 \dots dt_n = \frac{1}{r_2} (1 - e^{-r_2\delta}) \delta^{n-1}. \quad (23)$$

After unconditioning Eqn. (22) with respect to the arrival epochs by integrating over all t_i from t to $t+\delta$ and dividing by δ^n , we thus obtain

$$\mathbb{E}(\Lambda_1(t)\Lambda_2(t+\delta) \mid \mathcal{E}_{t,\delta}(n)) = \mathbb{E}\Lambda_1(t) \mathbb{E}\Lambda_2(t)e^{-r_2\delta} + \mathbb{E}\Lambda_1(t) \frac{1}{r_2\delta} (1 - e^{-r_2\delta}) n \mathbb{E}B_{12}$$

and hence, denoting $\Lambda_i := \lim_{t \rightarrow \infty} \Lambda_i(t)$ for $i = 1, 2$,

$$\begin{aligned} \mathbb{E}\Lambda_1(t)\Lambda_2(t+\delta) &= \sum_{n=0}^{\infty} \left(\mathbb{E}\Lambda_1(t)\Lambda_2(t)e^{-r_2\delta} + \mathbb{E}\Lambda_1(t) \frac{1}{r_2\delta} (1 - e^{-r_2\delta}) n \mathbb{E}B_{12} \right) \frac{(\delta\nu)^n}{n!} e^{-\nu\delta} \\ &= \mathbb{E}\Lambda_1(t)\Lambda_2(t)e^{-r_2\delta} + \mathbb{E}\Lambda_1(t) \frac{\nu}{r_2} (1 - e^{-r_2\delta}) \mathbb{E}B_{12} \\ &= \mathbb{E}\Lambda_1(t) \mathbb{E}\Lambda_2 + e^{-r_2\delta} (\mathbb{E}\Lambda_1(t)\Lambda_2(t) - \mathbb{E}\Lambda_1(t) \mathbb{E}\Lambda_2), \end{aligned}$$

where we made use of $\mathbb{E}\Lambda_i = \nu \mathbb{E}B_{1i}/r_i$ in the last equality. It follows that

$$\begin{aligned} \text{Cov}(\Lambda_1(t), \Lambda_2(t+\delta)) &= \mathbb{E}\Lambda_1(t)\Lambda_2(t+\delta) - \mathbb{E}\Lambda_1(t) \mathbb{E}\Lambda_2(t+\delta) \\ &= \mathbb{E}\Lambda_1(t)\Lambda_2(t)e^{-r_2\delta} + (1 - e^{-r_2\delta}) \mathbb{E}\Lambda_1(t) \mathbb{E}\Lambda_2 \\ &\quad - \mathbb{E}\Lambda_1(t) (\mathbb{E}\Lambda_2(t)e^{-r_2\delta} + (1 - e^{-r_2\delta}) \mathbb{E}\Lambda_2), \end{aligned}$$

from which the lemma follows. \square

References

- [1] D. Anderson, J. Blom, M. Mandjes, H. Thorsdottir, and K. de Turck. A functional central limit theorem for a Markov-modulated infinite-server queue. *Methodology and Computing in Applied Probability*, 18:153–168, 2016.
- [2] S. Bar-Lev, O. Boxma, B. Mathijsen, and D. Perry. A blood bank model with perishable blood and impatience. *Eurandom Reports*, 2015.
- [3] S. Centanni and M. Minozzo. A Monte Carlo approach to filtering for a class of marked doubly stochastic Poisson processes. *Journal of the American Statistical Association*, 101(476):1582–1597, 2006.
- [4] M. Csörgő and P. Révész. *Strong Approximations in Probability and Statistics*. Academic Press, New York, 1981.
- [5] A. Dassios and J.-W. Jang. Pricing of catastrophe reinsurance and derivatives using the Cox process with shot noise intensity. *Journal of Finance and Stochastics*, 7:73–95, 2003.
- [6] S. Eick, W. Massey, and W. Whitt. The physics of the $M_t/G/\infty$ queue. *Management Science*, 39(2):241–252, 1993.
- [7] S. Ethier and T. Kurtz. *Markov Processes: Characterization and Convergence*. Wiley, 1986.
- [8] X. Gao and L. Zhu. A functional central limit theorem for stationary Hawkes processes and its application to infinite-server queues. *To appear*, 2016.
- [9] A.G. Hawkes. Point spectra of some mutually exciting point processes. *Journal of the Royal Statistical Society*, 33:438–443, 1971.
- [10] A. Iksanov and Z. Jurek. Shot noise distributions and selfdecomposability. *Stochastic Analysis and Applications*, 21(3):593–609, 2003.
- [11] S.-H. Kim and W. Whitt. Are call center and hospital arrivals well modeled by nonhomogeneous Poisson processes? *Manufacturing and Service Operations Management*, 16(3):464–480, 2014.
- [12] P.A. Lewis. Simulation of non-homogenous Poisson processes by thinning. *Navel Research Logistics Quarterly*, 26(3):403–413, 1979.
- [13] Y. Ogata. Statistical models for earthquake occurrences and residual analysis for point processes. *Journal of the American Statistical Association*, 83:9–27, 1988.
- [14] G. Pang, R. Talreja, and W. Whitt. Martingale proofs of many-server heavy-traffic limits for Markovian queues. *Probability Surveys*, 4:193–267, 2007.
- [15] G. Pang and W. Whitt. Two-parameter heavy-traffic limits for infinite-server queues. *Queueing Systems*, 65:325–364, 2010.
- [16] S.M. Ross. *Stochastic Processes*. Wiley, New York, 1996.
- [17] W. Whitt. Proofs of the Martingale FCLT. *Probability Surveys*, 4:268–302, 2007.