

## Thomson-Rayleigh scattering in expanding plasmas

**Citation for published version (APA):**

Sanden, van de, M. C. M., Regt, de, J. M., van den Bercken, R. E. J., Meulenbroeks, R. F. G., Beulens, J. J., Buuron, A. J. M., & Schram, D. C. (1992). Thomson-Rayleigh scattering in expanding plasmas. In *ESCAMPIG : 11th, St. Petersburg, Russia, 25-29 August 1992* (pp. 49-52)

**Document status and date:**

Published: 01/01/1992

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

# THOMSON-RAYLEIGH SCATTERING IN EXPANDING PLASMAS

M.C.M. van de Sanden, J.M. de Regt, R. van den Bercken,  
R.F.G. Meulenbroeks, J.J. Beulens, A.J.M. Buuron and D.C. Schram

Department of Physics, Eindhoven University of Technology  
P.O.Box 513, 5600 MB Eindhoven, The Netherlands

Expanding plasmas are interesting from a fundamental as well as from a technological point of view. Nowadays expanding plasmas are used in plasma processing, e.g. plasma deposition [1, 2] and plasma spraying. The fundamental interest in expanding plasmas is the different behavior of the electrons and heavy particles due to their large mass difference. As a consequence, if the expansion is strong, charge separation can occur, which leads to a generated electromagnetic field [3]. This electromagnetic field in turn influences the behavior of the charged particles in the expanding plasma. Another interesting feature is the coupling between the ionization-recombination phenomena and the transport properties as electron temperature and electron density. In this respect the study of the equilibrium departure is essential, as it can help in the understanding of the kinetic processes involved.




Figure 1: A measured Thomson-Rayleigh profile with least mean square fit and five times magnified residue.




Figure 2:  $n_e$  on the axis as a function of the axial position for different background pressures.  $I_{cas} = 45$  A, Ar - flow = 3500 ml/min.

The measurements discussed in this paper were performed on a plasma which expands freely from a sub atmospheric thermal plasma in argon (a cascaded arc,  $T \approx 1$  eV,  $n_e \approx 10^{22} \text{ m}^{-3}$  and  $n_0 \approx 10^{23} \text{ m}^{-3}$ ) into a low background pressure ( $p \approx 10$ -100 Pa) [1, 2]. The electron density and temperature and the neutral particle temperature in the expansion part are measured locally by means of Thomson and Rayleigh scattering [4]. The

main components of the scattering diagnostic are a frequency doubled Nd:YAG laser, a polychromator based on a holographic grating and a gateable light amplifier in combination with a linear photodiode array. Much attention is given to the suppression of stray light which is essential if one wants to measure Rayleigh scattering (stray light level 0.4 Pa argon at 300 K). For more details cf. Refs. [4, 5]. In fig. 1 a typical Thomson-Rayleigh profile is given, in which the different components can be distinguished: a broadened Thomson profile with a superimposed Rayleigh profile.

In figs. 2 and 3 the measured electron density and temperature and the neutral particle density on the axis are shown for different background pressures. The supersonic expansion can be clearly seen. After several mm, depending on the background pressure, a shock occurs, after which the plasma expands further subsonically. The behavior of the densities and the electron temperature in the first region will be treated here in more detail.




Figure 3:  $T_e$  (left) and  $n_0$  (right) as a function of the axial position (cf. fig. 2).

In figs. 4 the measured electron and neutral particle densities in the expansion are compared with the model of Ashkenas and Sherman [6] developed for an adiabatic supersonic expansion of a gas. Following this model the densities should scale for  $z > z_0$  as

$$n = \frac{n_{res} z_{ref}^2}{(z - z_0)^2} \quad (1)$$

Equation 1 is the well known source expansion with  $n_{res}$  the reservoir particle density and  $z_0$  the origin from which the particle trajectories seem to originate.  $z_{ref}$  is a reference length. As can be seen from figs. 4 the agreement with the measurements of  $n_0$  and  $n_e$  is excellent ( $z_0 \approx 1.5$  mm,  $n_e = 1.3 \cdot 10^{22} m^{-3}$  and  $n_0 = 3 \cdot 10^{23} m^{-3}$ ). Equation (1) can be interpreted as follows. During the expansion thermal energy is converted into directed kinetic energy. Since the thermal energy is limited by the total enthalpy in the reservoir the velocity  $v_{axis}$  along the axis saturates and becomes finally constant. From flux conservation this means that the product  $n v_{axis} A = constant$ . Since the surface  $A$  scales

as  $z^2$  for large  $z$ , the densities scale as  $z^{-2}$  (cf. (1)). But the flux of particles is only constant if recombination can be neglected. This means that from figs. (4), it can be concluded that argon does not recombine on the particle level during the strong expansion or to put it in other words, the recombination is smaller than can be measured using the Thomson-Rayleigh scattering diagnostic.




Figure 4:  $n_e$  and  $n_0$  on the axis as a function of the axial position for different background pressures compared with the model of Ashkenas and Sherman (lines) [6].  $I_{cas} = 45$  A,  $Ar - flow = 3500$  ml/min.

The recombination of argon can be better examined by combining measurements of the excited state densities with Thomson-Rayleigh measurements using the Saha equation. In fig. 5 the calculated  $b_p$ -factor [7, 8] is given using the locally measured  $\frac{n_p}{g_p}$  and  $n_e$  and  $T_e$ . The  $b_p$ -factor is defined as the ratio of the measured excited state density to the excited state density according to the Saha equation for the given  $n_e$  and  $T_e$ . If the  $b_p$ -factor is smaller than one the plasma is recombining. As can be seen from fig. 5, the  $b_p$ -factor approaches one for the highly excited states, revealing the fact that the highly excited states are collisionally dominated, and is much smaller than one for the lower excited states. In fact from fig. 5 one can conclude that the plasma is recombining and from the gradient in the  $b_p$ -factor space the recombination rate can be calculated [5, 7].

Another way in which one can research the recombination phenomena is by means of the electron energy balance. In fig. 6  $n_e$  as a function of  $T_e$  is given in the expansion for different conditions. As can be seen a simple scaling law is found which reads  $n_e = C_0 T_e^\chi$  with  $\chi \approx 3.7$ . A similar result was found by Stevefelt and Cowins [9] for a laser produced carbon plasma. The explanation is the fact that although the three particle recombination can be neglected, it can *not* be neglected on the energy level, i.e. it has to be taken into account as a heat source for the electron gas. Therefore the electron gas does not expand adiabatically (corresponding with  $\chi = \frac{3}{2}$ ). Using a quasi-one dimensional model for the expansion and using the Ashkenas and Sherman result (cf. (1)), it can be shown that the determined  $\chi$  corresponds to a three particle recombination coefficient which reveals the




Figure 5:  $b_p$  as a function of the ionization potential  $I_p$  for different radial positions at  $z = 40$  mm.  $I_{cas} = 45$  A, Ar - flow = 3500 ml/min,  $p = 40$  Pa.




Figure 6:  $n_e$  as a function of  $T_e$  in the expansion for different conditions of the cascaded arc and background pressure.

well know  $T_e^{-9/2}$  dependence on the electron temperature.

We would like to thank M.J.F. van de Sande, H.M.M. de Jong and A.B.M. Hüsken for their skillful technical assistance.

## References

- [1] J.J. Beulens, A.J.M. Buuron, and D.C. Schram, *Surface and Coatings Techn.* **47** 401 (1991)
- [2] G.M.W. Kroesen, D.C. Schram and J.C.M. de Haas, *Plasma Chem. Plasma Proc.* **10** 531 (1990)
- [3] H.J.G. Gielen and D.C. Schram, *IEEE Trans. Plasma Sci.* **18** 127 (1990)
- [4] M.C.M. van de Sanden, J.M. de Regt, G.M. Janssen, D.C. Schram, J.A.M. van der Mullen, B. van der Sijde, accepted for publication in *Rev. Sci. Instr.* (june 1992)
- [5] M.C.M. van de Sanden, thesis University of Technology of Eindhoven, the Netherlands (1991)
- [6] H. Ashkenas and F.S. Sherman, *Rarefied Gasdynamics* **4**, Ed. J.H. de Leeuw (Academic Press, New York, 1966)
- [7] J.A.M. van der Mullen, *Phys. Rep.* **191** 109 (1990)
- [8] L.M. Bibermann, V.S. Vorob'ev and I.T. Yakubov, *Kinetics of Nonequilibrium Low Temperature Plasmas*, (Plenum, New York, 1987)
- [9] J. Stevefelt and C.B. Collins, *J. Phys. D* **24** 2149 (1991)