

A construction of generalized eigenprojections based on geometric measure theory

Citation for published version (APA):

Eijndhoven, van, S. J. L. (1985). *A construction of generalized eigenprojections based on geometric measure theory*. (Eindhoven University of Technology : Dept of Mathematics : memorandum; Vol. 8509). Technische Hogeschool Eindhoven.

Document status and date:

Published: 01/01/1985

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EINDHOVEN UNIVERSITY OF TECHNOLOGY

Department of Mathematics and Computing Science

Memorandum 85-09

A CONSTRUCTION OF GENERALIZED EIGENPROJECTIONS
BASED ON GEOMETRIC MEASURE THEORY

by

S.J.L. van Eijndhoven

Eindhoven University of Technology
P.O.Box 513, 5600 MB Eindhoven
The Netherlands

June 1985

A CONSTRUCTION OF GENERALIZED EIGENPROJECTIONS

BASED ON GEOMETRIC MEASURE THEORY

by

S.J.L. van Eijndhoven

Abstract

Let M denote a σ -compact locally compact metric space which satisfies certain geometrical conditions. Then for each σ -additive projection valued measure P on M there can be constructed a "canonical" Radon-Nikodym derivative $\pi: \alpha \mapsto \pi_\alpha$, $\alpha \in M$, with respect to a suitable basic measure ρ on M . The family $(\pi_\alpha)_{\alpha \in M}$ consists of generalized eigenprojections related to the commutative von Neumann algebra generated by the projections $P(\Delta)$, Δ a Borel set of M .

A.M.S. Classifications 46 F 10, 47 A 70.

I.

In this paper M denotes a σ -compact locally compact (and hence separable) metric space. It follows that any positive Borel measure on M is regular (cf. [3], p. 162). In the monograph [2], certain geometrical conditions on M are introduced, which lead to the following result.

0. Theorem

Let μ denote a positive Borel measure on M with the property that bounded Borel sets of M have finite μ -measure, and let f denote a Borel function which is μ -integrable on bounded Borel sets. Then there exists a μ -null set N_f such that for all $\alpha \in M \setminus N_f$ both $\mu(B(\alpha, r)) > 0$, and the limit

$$\tilde{f}(\alpha) = \lim_{r \downarrow 0} \mu(B(\alpha, r))^{-1} \int_{B(\alpha, r)} f \, d\mu$$

exists. We have $f = \tilde{f}$ μ -almost everywhere.

($B(\alpha, r)$ denotes the closed ball with radius r and centre α .)

Remark: In the previous theorem, the Borel function f can be replaced by a Borel measure ν with the property that bounded Borel sets of M have finite ν -measure. Then a "canonical" Radon-Nikodym derivative $\frac{d\nu}{d\mu}$ is obtained, which satisfies

$$\frac{d\nu}{d\mu}(\alpha) = \lim_{r \downarrow 0} \frac{\nu(B(\alpha, r))}{\mu(B(\alpha, r))}$$

μ -almost everywhere.

In the sequel we assume that M also satisfies Federer's geometrical conditions. As examples of such spaces M we mention

- finite dimensional vector spaces with metric $d(x,y) = v(x - y)$ where v is any norm,
- Riemannian manifolds (of class ≥ 2) with their usual metric.

Let X denote a separable Hilbert space with inner product (\cdot, \cdot) and let there be given a σ -additive projection valued set function P on M .

So for all Borel sets $\Delta \subset M$, $P(\Delta)$ is an orthogonal projection on X .

Moreover, if Δ is the disjoint union $\bigcup_{j=1}^{\infty} \Delta_j$, then $P(\Delta) = \sum_{j=1}^{\infty} P(\Delta_j)$. In particular $\sum_{j=1}^{\infty} P(\Delta_j) = I$ if $\bigcup_{j=1}^{\infty} \Delta_j = M$.

Now let R denote a positive bounded linear operator on X with the property that for each bounded Borel set Δ the positive operator $RP(\Delta)R$ is trace class. E.g. for R any positive Hilbert-Schmidt operator can be taken.

For each bounded Borel set Δ we define $\rho(\Delta) = \text{trace}(RP(\Delta)R)$. In a natural way, ρ becomes a σ -finite positive Borel measure on M . Each bounded Borel set of M has a finite ρ -measure.

We take a fixed orthonormal basis $(v_k)_{k \in \mathbb{N}}$ in X , and for each $k, \ell \in \mathbb{N}$ we define the set function

$$\Phi_{k\ell} : \Delta \mapsto (RP(\Delta)R v_\ell, v_k), \quad \Delta \text{ Borel.}$$

The set functions $\Phi_{k\ell}$ are absolutely continuous with respect to ρ .

By Theorem 0, there exists a null set N_1 and there exist Borel functions $\hat{\Phi}_{k\ell}$

such that for all $k, \ell \in \mathbb{N}$ and all $\alpha \in M \setminus N_1$

$$\hat{\phi}_{k\ell}(\alpha) = \lim_{r \rightarrow 0} \left\{ \frac{\Phi_{k\ell}(B(\alpha, r))}{\rho(B(\alpha, r))} \right\} .$$

1. Lemma

Let $\alpha \in M \setminus N_1$. Then for all $k, \ell \in \mathbb{N}$

$$|\hat{\phi}_{k\ell}(\alpha)|^2 \leq \hat{\phi}_{kk}(\alpha) \hat{\phi}_{\ell\ell}(\alpha) .$$

Proof. Consider the estimation,

$$\begin{aligned} |\hat{\phi}_{k\ell}(\alpha)|^2 &= \lim_{r \rightarrow 0} \left| \frac{\Phi_{k\ell}(B(\alpha, r))}{\rho(B(\alpha, r))} \right|^2 \leq \\ &\leq \lim_{r \rightarrow 0} \left\{ \frac{\Phi_{kk}(B(\alpha, r))}{\rho(B(\alpha, r))} \right\} \lim_{r \rightarrow 0} \left\{ \frac{\Phi_{\ell\ell}(B(\alpha, r))}{\rho(B(\alpha, r))} \right\} = \\ &= \hat{\phi}_{kk}(\alpha) \hat{\phi}_{\ell\ell}(\alpha) . \quad \square \end{aligned}$$

The function $\sum_{k=1}^{\infty} \hat{\phi}_{kk}$ is Borel, and the functions $\hat{\phi}_{kk}$ are positive. So for each bounded Borel set Δ , we have

$$\int_{\Delta} \left(\sum_{k=1}^{\infty} \hat{\phi}_{kk} \right) d\rho = \sum_{k=1}^{\infty} \Phi_{kk}(\Delta) = \rho(\Delta) .$$

Then Theorem 0 yields a null set $N_2 \supset N_1$ such that for all $\alpha \in M \setminus N_2$,

$$\sum_{k=1}^{\infty} \hat{\phi}_{kk}(\alpha) = \lim_{r \rightarrow 0} \frac{\int_{B(\alpha, r)} \left(\sum_{k=1}^{\infty} \hat{\phi}_{kk} \right) d\rho}{\rho(B(\alpha, r))} = 1$$

2. Corollary

Let $\alpha \in M \setminus N_2$. Then $\sum_{k, \ell=1}^{\infty} |\hat{\phi}_{k\ell}(\alpha)|^2 < \infty$.

Proof. Consider the estimation

$$\sum_{k, \ell=1}^{\infty} |\hat{\phi}_{k\ell}(\alpha)|^2 \leq \sum_{k=1}^{\infty} \hat{\phi}_{kk}(\alpha) \sum_{\ell=1}^{\infty} \hat{\phi}_{\ell\ell}(\alpha) = 1 . \quad \square$$

3. Definition

The operators $B_\alpha : X \rightarrow X$, $\alpha \in M$, are defined by

$$\left[\begin{array}{l} B_\alpha = 0 \quad \text{for } \alpha \in N_2 , \\ B_\alpha x = \sum_{k, \ell=1}^{\infty} \hat{\phi}_{k\ell}(\alpha) (x, v_\ell) v_k , \quad x \in X, \quad \alpha \in M \setminus N_2 . \end{array} \right.$$

Observe that B_α is a Hilbert-Schmidt operator for each $\alpha \in M$.

The operators B_α are related to the set function P in the following way.

4. Lemma

Let $\alpha \in M \setminus N_2$. Then we have

$$\lim_{r \rightarrow 0} \left\| B_\alpha - \frac{RP(B(\alpha, r))R}{\rho(B(\alpha, r))} \right\|_{HS} = 0$$

with $\|\cdot\|_{HS}$ the Hilbert-Schmidt norm.

Proof

For all $r > 0$,

$$\left\| B_\alpha - \frac{RP(B(\alpha, r))R}{\rho(B(\alpha, r))} \right\|^2 = \sum_{k, \ell=1}^{\infty} \left| \hat{\phi}_{k\ell}(\alpha) - \frac{\Phi_{k\ell}(B(\alpha, r))}{\rho(B(\alpha, r))} \right|^2$$

Let $\varepsilon > 0$. Take a fixed $A \in \mathbb{N}$ so large that

$$(*) \quad \sum_{k=A+1}^{\infty} \hat{\phi}_{kk}(\alpha) < \varepsilon^2/4.$$

Next, take $r_0 > 0$ so small that for all r , $0 < r < r_0$, and all $k, \ell \in \mathbb{N}$ with $k, \ell \leq A$.

$$(**) \quad \left| \hat{\phi}_{k\ell}(\alpha) - \frac{\Phi_{k\ell}(B(\alpha, r))}{\rho(B(\alpha, r))} \right| < \varepsilon/A$$

and also

$$(***) \quad \sum_{k=A+1}^{\infty} \frac{\Phi_{kk}(B(\alpha, r))}{\rho(B(\alpha, r))} < \varepsilon^2.$$

Then we obtain the following estimation

$$\begin{aligned} & \left(\sum_{k=1}^A \sum_{\ell=1}^A + 2 \sum_{k=A+1}^{\infty} \sum_{\ell=1}^{\infty} \right) \left| \hat{\phi}_{k\ell}(\alpha) - \frac{\Phi_{k\ell}(B(\alpha, r))}{\rho(B(\alpha, r))} \right|^2 \\ & \leq \varepsilon^2 + 4 \sum_{k=A+1}^{\infty} \sum_{\ell=1}^{\infty} \left(|\hat{\phi}_{k\ell}(\alpha)| + \frac{|\Phi_{k\ell}(B(\alpha, r))|^2}{\rho(B(\alpha, r))^2} \right). \end{aligned}$$

By (*)

$$\sum_{k=A+1}^{\infty} \sum_{\ell=1}^{\infty} |\hat{\phi}_{k\ell}(\alpha)|^2 \leq \sum_{k=A+1}^{\infty} \hat{\phi}_{kk}(\alpha) \leq \varepsilon^2/4$$

and by (***)

$$\sum_{k=A+1}^{\infty} \sum_{\ell=1}^{\infty} \frac{|\Phi_{k\ell}(B(\alpha, r))|^2}{\rho(B(\alpha, r))^2} \leq \sum_{k=A+1}^{\infty} \frac{\Phi_{kk}(B(\alpha, r))}{\rho(B(\alpha, r))} < \varepsilon^2.$$

Thus it follows that

$$\left\| B_{\alpha} - \frac{RP(B(\alpha, r))R}{\rho(B(\alpha, r))} \right\|_{HS} < \varepsilon\sqrt{6}$$

for all r with $0 < r < r_0$.

□

In a natural way, the projection valued set function \mathcal{P} can be linked to the function-algebra $L_{\infty}(M, \rho)$. To show this, let $x, y \in X$. Then the finite measure $\mu_{x,y}$ is defined by $\mu_{x,y}(\Delta) = (\mathcal{P}(\Delta)x, y)$ where Δ is any Borel set. We have $\int_M d\mu_{x,y} = (x, y)$. Clearly, $\mu_{x,y}$ is absolutely continuous with respect to ρ .

Let f denote a Borel function on M which is bounded on bounded Borel sets. Then we define the operator T_f by

$$D(T_f) = \{x \in X \mid \int_M |f|^2 d\mu_{x,x} < \infty\}$$

and for $x \in D(T_f)$

$$(T_f x, y) = \int_M f d\mu_{x,y}, \quad y \in X .$$

Observe that T_f is a normal operator in X . Since f is bounded on bounded Borel sets we derive for each $r > 0$, $\alpha \in M$ and $x \in X$,

$$\begin{aligned} |(T_f P(B(\alpha, r))x, x)| &\leq \int_M |f| \chi_{B(\alpha, r)} d\mu_{x,x} \leq \\ &\leq \left(\sup_{\lambda \in B(\alpha, r)} |f(\lambda)| \right) (P(B(\alpha, r))x, x) . \end{aligned}$$

So $RT_f P(B(\alpha, r))R$ is a trace class operator.

5. Lemma

There exists a null set N_3 such that for all $\alpha \in M \setminus N_3$

$$\lim_{r \rightarrow 0} \left\| f(\alpha) B_\alpha - \frac{RT_f P(B(\alpha, r))R}{\rho(B(\alpha, r))} \right\|_{HS} = 0 .$$

Proof

Following Lemma 4, we are ready if we can prove that there exists a null set

$N_3 \supset N_2$ such that for all $\alpha \in M \setminus N_3$

$$\lim_{r \rightarrow 0} \left\| f(\alpha) \frac{RP(B(\alpha, r))R}{\rho(B(\alpha, r))} - \frac{RT_f P(B(\alpha, r))R}{\rho(B(\alpha, r))} \right\|_{HS} = 0 .$$

Therefore we estimate as follows

$$\begin{aligned} & \sum_{k, \ell=1}^{\infty} \rho(B(\alpha, r))^{-2} \left| \int_{B(\alpha, r)} (f(\alpha) - f(\lambda)) d\mu_{Rv_k, Rv_\ell}(\lambda) \right|^2 \leq \\ & \leq \rho(B(\alpha, r))^{-2} \left(\int_{B(\alpha, r)} |f(\alpha) - f(\lambda)|^2 d\rho(\lambda) \right) \left(\int_{B(\alpha, r)} \left(\sum_{k, \ell=1}^{\infty} |\hat{\phi}_{k\ell}(\lambda)|^2 \right) d\rho(\lambda) \right) \\ & \leq \rho(B(\alpha, r))^{-1} \int_{B(\alpha, r)} |f(\alpha) - f(\lambda)|^2 d\rho(\lambda) \end{aligned}$$

because

$$\sum_{k, \ell=1}^{\infty} |\hat{\phi}_{k\ell}(\lambda)|^2 \leq \left(\sum_{k=1}^{\infty} \hat{\phi}_{kk}(\lambda) \right)^2 = 1 .$$

Now there exists a null set $N_3 \supset N_2$ such that the latter expression tends to zero as $r \rightarrow 0$ for all $\alpha \in M \setminus N_3$.

II.

In the second part of this paper we employ the above auxiliary results in the announced construction of generalized eigenprojections.

We consider the triple of Hilbert spaces

$$R(X) \subseteq X \subseteq R^{-1}(X) .$$

Here $R(X)$ is the Hilbert space with inner product $(\cdot, \cdot)_1$,

$$(u, w)_1 = (R^{-1}u, R^{-1}w), \quad u, w \in R(X),$$

and $R^{-1}(X)$ is the completion of X with respect to the inner product $(\cdot, \cdot)_{-1}$,

$$(x, y)_{-1} = (Rx, Ry).$$

The spaces $R(X)$ and $R^{-1}(X)$ are in duality through the pairing $\langle \cdot, \cdot \rangle$,

$$\langle w, G \rangle = (R^{-1}w, RG), \quad w \in R(X), G \in R^{-1}(X).$$

6. Definition

For each $\alpha \in M$, we define the operator $\pi_\alpha : R(X) \rightarrow R^{-1}(X)$ by

$$R\pi_\alpha w = B_\alpha R^{-1}w, \quad w \in R(X).$$

Cf. Definition 3.

Observe that $\pi_\alpha : R(X) \rightarrow R^{-1}(X)$ is continuous.

7. Theorem

I. For all $\alpha \in M \setminus N_2$ and for all $w \in R(X)$

$$\lim_{r \rightarrow 0} \left\| \pi_\alpha w - \frac{P(B(\alpha, r))}{\rho(B(\alpha, r))} w \right\|_{-1} = 0.$$

II. Let $f : M \rightarrow \mathbb{C}$ be a Borel function which is bounded on bounded Borel sets. Then there exists a null set $N_f \supset N_2$ such that for all $\alpha \in M \setminus N_f$ and all $w \in R(X)$

$$\lim_{r \rightarrow 0} \left\| f(\alpha) \pi_\alpha w - T_f \frac{P(B(\alpha, r))}{\rho(P(\alpha, r))} w \right\|_{-1} = 0.$$

Proof

The proof of I follows from Lemma 4 and the inequality

$$\left\| \pi_{\alpha} w - \frac{P(B(\alpha, r))}{\rho(B(\alpha, r))} w \right\|_{-1} \leq \left\| R\pi_{\alpha}R - \frac{RP(B(\alpha, r))R}{\rho(B(\alpha, r))} \right\|_{HS} \|R^{-1}w\| .$$

The proof of II follows from Lemma 5 and the inequality

$$\begin{aligned} & \left\| f(\alpha)\pi_{\alpha} w - \frac{T_f P(B(\alpha, r))w}{\rho(B(\alpha, r))} \right\|_{-1} \leq \\ & \leq \left\| f(\alpha)R\pi_{\alpha}R - \frac{RT_f P(B(\alpha, r))R}{\rho(B(\alpha, r))} \right\|_{HS} \|R^{-1}w\| . \end{aligned} \quad \square$$

8. Corollary

Let the operator $RT_f R^{-1}$ be closable in X . Then T_f is closable as an operator from $R^{-1}(X)$ into $R^{-1}(X)$. For its closure \bar{T}_f we have

$$\bar{T}_f \pi_{\alpha} w = f(\alpha)\pi_{\alpha} w$$

with $w \in R(X)$ and $\alpha \in M \setminus N_f$.

The results stated in Theorem 7 and Corollary 8 indicate that the mappings $\pi_{\alpha} : R(X) \rightarrow R^{-1}(X)$ give rise to ("candidate") generalized eigenspaces $\pi_{\alpha}R(X)$ for the commutative von Neumann algebra $\{T_f | f \in L_{\infty}(M, \rho)\}$.

Finally, we explain in which way the operators π_{α} , $\alpha \in M$, can be seen as generalized projections.

9. Lemma

Let $w \in R(X)$. Then in weak sense

$$w = \int_M \pi_\alpha w d\rho(\alpha) .$$

So for all $v \in R(X)$,

$$(v, w) = \int_M \langle v, \pi_\alpha w \rangle d\rho(\alpha) .$$

Proof. Let Δ be a bounded Borel set. For all $v \in R(X)$,

$$\begin{aligned} \sum_{k, \ell=1}^{\infty} |\phi_{k\ell}(\alpha)(R^{-1}v, v_\ell)(v_k, R^{-1}w)| &\leq \\ &\leq \left(\sum_{k, \ell=1}^{\infty} |\phi_{k\ell}(\alpha)|^2 \right)^{\frac{1}{2}} \|R^{-1}w\| \|R^{-1}v\| , \end{aligned}$$

and hence by Fubini's theorem

$$\begin{aligned} \int_{\Delta} \langle v, \pi_\alpha w \rangle d\rho(\alpha) &= \sum_{k, \ell=1}^{\infty} \Phi_{k\ell}(\Delta)(R^{-1}v, v_\ell)(v_k, R^{-1}w) = \\ &= (P(\Delta)v, w) . \end{aligned}$$

Since M can be written as the disjoint union of bounded Borel sets it follows that

$$\int_M \langle v, \pi_\alpha w \rangle d\rho(\alpha) = (v, w) .$$

□

Remark: If R is Hilbert-Schmidt, the integral $\int_M \pi_\alpha w d\rho(\alpha)$ exists in strong sense. So in addition we have

$$\int_M \|R\pi_\alpha w\| d\rho(\alpha) < \infty .$$

In $\pi_\alpha R(X)$ we define the sesquilinear form $(\cdot, \cdot)_\alpha$ by

$$(F, G)_\alpha = \langle v, \pi_\alpha w \rangle ,$$

where $F = \pi_\alpha v$, $G = \pi_\alpha w$. $(F, G)_\alpha$ does not depend on the choice of v and w .

It can be shown easily that $(\cdot, \cdot)_\alpha$ is a well-defined non-degenerate sesquilinear form in $\pi_\alpha R(X)$. By X_α we denote the completion of $\pi_\alpha R(X)$ with respect to this sesquilinear form.

10. Theorem

I. The Hilbert space X_α with inner product $(\cdot, \cdot)_\alpha$ is a Hilbert subspace of $R^{-1}(X)$. π_α maps $R(X)$ continuously into X_α .

II. Let f be a Borel function which is bounded on bounded Borel sets.

Suppose the operator T_f is closable in $R^{-1}(X)$ with closure \bar{T}_f . Then

there exists a null set N_f such that for each $\alpha \in M \setminus N_f$ and all

$G \in X_\alpha$ we have

$$\bar{T}_f G = f(\alpha)G .$$

Proof.

I. Let $G \in \pi_\alpha R(X)$, $G = \pi_\alpha w$. We estimate as follows

$$\begin{aligned} \|RG\|^2 &= \langle R^2 \pi_\alpha w, \pi_\alpha w \rangle \leq \\ &\leq \langle R^2 \pi_\alpha w, \pi_\alpha R^2 \pi_\alpha w \rangle^{\frac{1}{2}} \langle w, \pi_\alpha w \rangle^{\frac{1}{2}} \leq \\ &\leq \|R\pi_\alpha R\|^{\frac{1}{2}} \|R\pi_\alpha w\| \|\pi_\alpha w\|_\alpha . \end{aligned}$$

It follows that

$$\|G\|_{-1} \leq \|R\pi_\alpha R\|^{\frac{1}{2}} \|G\|_\alpha .$$

Hence X_α can be seen as a subspace of $R^{-1}(X)$.

II. By Corollary 8, there exists a null set N_f such that for all $\alpha \in M \setminus N_f$ and for all $w \in R(X)$

$$\bar{T}_f \pi_\alpha w = f(\alpha) \pi_\alpha w .$$

Let $\alpha \in M \setminus N_f$. Since $X_\alpha \hookrightarrow R^{-1}(X)$ and $\pi_\alpha R(X)$ is dense in X_α it follows that for all $G \in X_\alpha$, $G \in \text{Dom}(\bar{T}_f)$ and $\bar{T}_f G = f(\alpha)G_\alpha$. □

11. Corollary

Let $\pi_\alpha^+ : X_\alpha \rightarrow R^{-1}(X)$ denote the adjoint of π_α .

Then $\pi_\alpha^+ \pi_\alpha = \pi_\alpha$.

Proof

Let $w, v \in R(X)$. We have

$$\langle w, \pi_\alpha^+ v \rangle = (\pi_\alpha w, \pi_\alpha v)_\alpha = \langle w, \pi_\alpha^+ \pi_\alpha v \rangle . \quad \square$$

Let $(u_k)_{k \in \mathbb{N}}$ denote an orthonormal basis in X which is contained in $R(X)$. For each $\alpha \in M$, the sequence $(\pi_\alpha u_k)_{k \in \mathbb{N}}$ is total in X_α . So the spaces X_α , $\alpha \in M$, establish a measurable field of Hilbert spaces. Its field structure S is defined by

$\phi \in S \Leftrightarrow$ the functions $\alpha \mapsto (\phi(\alpha), \pi_\alpha u_k)_\alpha$ are Borel functions.

So the direct integral $H = \int_M^\oplus X_\alpha d\rho(\alpha)$ is well-defined.

(For the general theory of direct integrals, see [1], p. 161-172.)

The vector fields $\alpha \mapsto \pi_\alpha u_k$, $\alpha \in M$, $k \in \mathbb{N}$, give rise to an orthonormal system $(\phi_k)_{k \in \mathbb{N}}$ in H . (We recall that the elements of H are equivalence classes of square integrable vector fields.) We define the isometry

$U : X \rightarrow H$ by

$$Ux = \sum_{k=1}^{\infty} (x, u_k) \phi_k, \quad x \in X.$$

Then for all $x, y \in X$ we have

$$(x, y) = \int_M d\mu_{x, y} = \int_M ((Ux)(\alpha), (Uy)(\alpha))_\alpha d\rho(\alpha).$$

It follows that for all $x, y \in X$ and all $f \in L_\infty(M, \rho)$

$$(T_f x, y) = \int_M f d\mu_{x, y} = \int_M f(\alpha) ((Ux)(\alpha), (Uy)(\alpha))_\alpha d\rho(\alpha),$$

and hence we can write

$$UT_f x = \int_M^{\oplus} f(\alpha) (Ux)(\alpha) d\rho(\alpha) .$$

12. Lemma

The operator $U : X \rightarrow H$ is unitary.

Proof . We show that the set $U(\{T_f u_k | k \in \mathbb{N}, f \in L_\infty(M, \rho)\})$ is total in H .

Let ϕ be a square integrable vector field such that for all $f \in L_\infty(M, \rho)$ and all $k \in \mathbb{N}$

$$0 = (\phi, T_f u_k)_H = \int_M f(\alpha) (\phi(\alpha), \pi_\alpha u_k)_\alpha d\rho(\alpha) .$$

Since $f \in L_\infty(M, \rho)$ is arbitrary taken, $(\phi(\alpha), \pi_\alpha u_k)_\alpha$ vanishes except on a set \tilde{N}_k of measure zero. Taking $\tilde{N} = \bigcup_{k=1}^{\infty} \tilde{N}_k$ this yields $\phi(\alpha) = 0$ on $M \setminus \tilde{N}$, and hence

$$\int_M \|\phi(\alpha)\|_\alpha^2 d\rho(\alpha) = 0 . \quad \square$$

Now the mappings $\pi_\alpha, \alpha \in M$, can be seen as generalized projections as follows: Let $w \in \mathcal{R}(X)$. The vector field $\alpha \mapsto \pi_\alpha w$ is a representant of the class Uw . These representants $\alpha \mapsto \pi_\alpha w, w \in \mathcal{R}(X)$, are canonical. Indeed, there exists a null set $N (= N_2)$ such that for all $w \in \mathcal{R}(X)$, and for all $\alpha \in M \setminus N$,

$$\lim_{r \downarrow 0} \left\| \pi_\alpha w - \rho(B(\alpha, r))^{-1} \int_{B(\alpha, r)} \pi_\lambda w \, d\rho(\lambda) \right\|_{-1} = 0 .$$

(Cf. Theorem 7.)

So the family $(\pi_\alpha)_{\alpha \in M}$ selects a canonical representant out of each class $U_w, w \in R(X)$. In this sense, each π_α "projects" $R(X)$ densely into X_α .

References

1. Federer, H., Geometric measure theory.
Die Grundlehren der mathematischen Wissenschaften, Band 153,
Springer-Verlag, 1969, Berlin.
2. Dixmier, J., Von Neumann Algebras.
North-Holland Mathematical Library, Vol. 27, 1981, Amsterdam.
3. Weir, A.J., General integration and measure.
Cambridge University Press, 1974, Cambridge.

Eindhoven University of Technology

P.O. Box 513

5600 MB Eindhoven

The Netherlands.