

## Aymptotics in normal order statistics

***Citation for published version (APA):***

Brands, J. J. A. M. (1986). *Aymptotics in normal order statistics*. (Eindhoven University of Technology : Dept of Mathematics : memorandum; Vol. 8605). Technische Hogeschool Eindhoven.

***Document status and date:***

Published: 01/01/1986

***Document Version:***

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

***Please check the document version of this publication:***

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

***General rights***

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

***Take down policy***

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

EINDHOVEN UNIVERSITY OF TECHNOLOGY  
Department of Mathematics and Computing Science

Memorandum 1986-05

May 1986

ASYMPTOTICS IN NORMAL ORDER STATISTICS

by

J.J.A.M. Brands

Eindhoven University of Technology  
Department of Mathematics and Computing Science  
PO Box 513  
5600 MB Eindhoven  
The Netherlands

# ASYMPTOTICS IN NORMAL ORDER STATISTICS

by

J.J.A.M. BRANDS

Department of Mathematics, Eindhoven University of Technology,  
The Netherlands

## ABSTRACT

In order statistics certain integrals involving the standard normal distribution play an important role. The asymptotic behaviour with respect to a large parameter is studied.

## 1. INTRODUCTION

The expectation and variance of the maximum in a random sample of size  $n$  from the standard normal distribution involve some of the integrals

$$(1) \quad M_j(n) := \int_{-\infty}^{\infty} x^{j-1} \phi(x) (1 - \phi^{n-1}(x)) dx \quad (n \in \mathbb{N}, j \in \mathbb{N}) ;$$

where

$$(2) \quad \phi(x) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-s^2/2} ds .$$

Integration by parts gives

$$(3) \quad \mu_j(n) = j M_j(n) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x^j e^{-x^2/2} dx ,$$

where

$$(4) \quad \mu_j(n) := \frac{n}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x^j e^{-x^2/2} \phi^{n-1}(x) dx \quad (n \in \mathbb{N}, j \in \mathbb{N}) .$$

The problem posed by two colleagues <sup>\*</sup>) of the author is to determine the asymptotic behaviour for  $n \rightarrow \infty$  of  $\mu_j(n)$  for  $j = 1, 2, 3, 4$ . Moreover, they are interested especially in the asymptotic behaviour of

$$(\mu_3 - \mu_1 \mu_2)(\mu_2 - \mu_1^2)^{-\frac{1}{2}} \quad \text{and} \quad (\mu_4 - \mu_2^2)^{\frac{1}{2}} .$$

We remark that the differences

$$M_j(n+1) - M_j(n) = \int_{-\infty}^{\infty} x^{j-1} (1 - \phi(x)) \phi^n(x) dx$$

are just the integrals occurring in the coefficients of the asymptotic formulas in a previous paper [1] (where  $f$  is defined by  $f(x) = \phi(x\sqrt{2})$ ).

## 2. RESULTS

Let the asymptotic series of  $(1 - \phi(x))(\phi'(x))^{-1}$  be denoted by  $A$ , i.e.

$$A := \sum_{\ell=0}^{\infty} (-1)^\ell (2\ell-1)!! x^{-2\ell-1} \quad (x \rightarrow \infty), \quad ((-1)!! = 1) .$$

Formal differentiations of  $A$  are denoted by  $A'$ ,  $A''$ , etc. Let  $x_1 = x_1(n)$  be defined by  $\phi(x_1) = 1 - \frac{1}{n}$ . Then  $\mu_j(n)$  has an asymptotic expansion in powers of  $x_1^{-1}$ , i.e.

$$\mu_j(n) \approx \sum_{k=-j}^{\infty} C(j,k) x_1^{-k} \quad (n \rightarrow \infty),$$

which can be computed as follows:  $x$  is considered to be a function of a variable  $z$  and  $\frac{dx}{dz} \approx \frac{1}{2}A$ . Higher derivatives can be computed by means of the chain rule. For instance,  $\frac{d^2x}{dz^2} \approx \frac{1}{2}AA'$ . Then

$$\mu_j(n) \approx \sum_{\ell=0}^{\infty} \left( \frac{d^\ell x^j}{dz^\ell} \right)_1 \frac{(-2)^\ell \Gamma(\ell)(1)}{\ell!} \quad (n \rightarrow \infty),$$

where the subscript 1 means that the value at  $x = x_1$  has to be taken.  $x_1$  has the following asymptotic series:

---

<sup>\*</sup>) F.W. Steutel and D.A. Overdijk, Department of Mathematics, Eindhoven University of Technology, The Netherlands.

$$x_1 \approx z^{\frac{1}{2}} \left( 1 + \sum_{k=1}^{\infty} q_k(\log z) z^{-k} \right) \quad (n \rightarrow \infty)$$

where  $z = 2 \log \frac{n}{\sqrt{2\pi}}$  and the  $q_k$ 's are polynomials of degree  $k$ . A few  $q_k$ 's are

$$q_1(t) = -\frac{1}{2} t, \quad q_2(t) = -\frac{1}{8} t^2 + \frac{1}{2} t - 1,$$

$$q_3(t) = -\frac{1}{16} t^3 + \frac{1}{2} t^2 - \frac{3}{2} t + \frac{7}{2}.$$

The coefficients  $C(j,k)$  have the property that  $C(j,-j+s) = 0$  if  $s$  is odd and  $C(j,-j) = 1$ . A few more coefficients  $C(j,k)$  are:

$$C(1,1) = -\Gamma'(1), \quad C(1,3) = \Gamma'(1) - \frac{1}{2} \Gamma''(1),$$

$$C(1,5) = 3\Gamma'(1) + 2\Gamma''(1) - \frac{1}{2} \Gamma'''(1);$$

$$C(2,0) = -2\Gamma'(1), \quad C(2,2) = 2\Gamma'(1), \quad C(2,4) = -6\Gamma'(1) + 2\Gamma''(1),$$

$$C(2,6) = 30\Gamma'(1) - 14\Gamma''(1) + \frac{8}{3} \Gamma'''(1);$$

$$C(3,-1) = -3\Gamma'(1), \quad C(3,1) = 3\Gamma'(1) + \frac{3}{2} \Gamma''(1),$$

$$C(3,3) = -9\Gamma'(1) + \frac{1}{2} \Gamma'''(1),$$

$$C(3,5) = 45\Gamma'(1) - \frac{21}{2} \Gamma''(1) - \frac{1}{2} \Gamma'''(1) + \frac{3}{8} \Gamma^{(4)}(1);$$

$$C(4,-2) = -4\Gamma'(1), \quad C(4,0) = 4\Gamma'(1) + 4\Gamma''(1),$$

$$C(4,2) = -12\Gamma'(1) - 4\Gamma''(1), \quad C(4,4) = 60\Gamma'(1) - \frac{8}{3} \Gamma'''(1).$$

A routine computation shows that

$$\frac{\mu_3 - \mu_1 \mu_2}{(\mu_2 - \mu_1)^{\frac{1}{2}}} \approx d_0 + d_2 x_1^{-2} + d_4 x_1^{-4} + \dots \quad (n \rightarrow \infty)$$

$$(\mu_4 - \mu_2)^{\frac{1}{2}} \approx e_0 + e_2 x_1^{-2} + e_4 x_1^{-4} + \dots \quad (n \rightarrow \infty)$$

where

$$d_0 = e_0 = 2(\Gamma''(1) - (\Gamma'(1))^2)^{\frac{1}{2}}$$

and

$$d_2 = e_2 = -d_0 .$$

### 3. PROOF OF THE RESULTS

We transform the integral in (4) by putting

$$(5) \quad \phi(x) = 1 - \frac{s}{n} .$$

Then

$$(6) \quad \frac{dx}{ds} = - \frac{\sqrt{2\pi}}{n} e^{x^2/2} ,$$

whence

$$(7) \quad \mu_j(n) = \int_0^n x^j \left(1 - \frac{s}{n}\right)^{n-1} ds .$$

We observe that  $x = x(s)$  is monotonically decreasing on  $[0, \infty)$ , that  $x(s) \rightarrow \infty$  ( $s \downarrow 0$ ),  $x(\frac{1}{2}n) = 0$  and  $x(s) \rightarrow -\infty$  ( $s \uparrow n$ ).

Now we shall prove that

$$(8) \quad \mu_j(n) = \left(1 + O\left(\frac{\log^2 n}{n}\right)\right) \int_0^{\log n} x^j e^{-s} ds + O(n^{-1} (\log n)^{j/2}) \quad (n \rightarrow \infty) .$$

Since

$$\begin{aligned} \left| \int_{\pi/2}^n x^j \left(1 - \frac{s}{n}\right)^{n-1} ds \right| &= \frac{n}{\sqrt{2\pi}} \int_{-\infty}^0 |x|^j e^{-x^2/2} \phi^{n-1}(x) dx \leq \\ &\leq \frac{n2^{-n}}{\sqrt{2\pi}} \int_0^{\infty} x^j e^{-x^2/2} dx \end{aligned}$$

we can write

$$(9) \quad \mu_j(n) = \int_0^{\frac{1}{2}n} x^j \left(1 - \frac{s}{n}\right)^{n-1} ds + O(n2^{-n}) \quad (n \rightarrow \infty) .$$

Using

$$(10) \quad \Phi(x) \approx 1 - \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \sum_{\ell=0}^{\infty} \frac{(-1)^\ell (2\ell-1)!!}{x^{2\ell-1}} \quad (x \rightarrow \infty)$$

we derive easily that at  $s = \log n$

$$(11) \quad x \sim \sqrt{2 \log n} \quad (n \rightarrow \infty) .$$

Hence

$$(12) \quad \int_{\log n}^{n/2} x^j \left(1 - \frac{s}{n}\right)^{n-1} ds = O\left((\log n)^{j/2} \int_{\log n}^{n/2} \left(1 - \frac{s}{n}\right)^{n-1} ds\right) = \\ = O(n^{-1} (\log n)^{j/2}) \quad (n \rightarrow \infty) .$$

From (9) and (12) it follows that

$$(13) \quad \mu_j(n) = \int_0^{\log n} x^j \left(1 - \frac{s}{n}\right)^{n-1} ds + O(n^{-1} (\log n)^{j/2}) \quad (n \rightarrow \infty) .$$

Furthermore

$$(14) \quad \int_0^{\log n} x^j \left(1 - \frac{s}{n}\right)^{n-1} ds = \left(1 + O\left(\frac{\log^2 n}{n}\right)\right) \int_0^{\log n} x^j e^{-s} ds \quad (n \rightarrow \infty)$$

since

$$(15) \quad \left(1 - \frac{s}{n}\right)^{n-1} = e^{-s} \left(1 + O\left(\frac{\log^2 n}{n}\right)\right) \quad (0 \leq s \leq \log n, n \rightarrow \infty) .$$

Clearly (13) and (14) imply (8).

On the interval  $0 \leq s \leq \log n$ , corresponding with large values of  $x$ , we can use (10) in order to solve  $x$  from (5) as a function of  $s$ .

Introduction of

$$(16) \quad z = 2 \log \frac{n}{\sqrt{2\pi}} - 2 \log s$$

transforms (5) into

$$(17) \quad \Phi(x) = 1 - \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z} .$$

Clearly  $z \rightarrow \infty$  if  $n \rightarrow \infty$  and  $0 < s < \log n$ .

Using (10) and taking logarithms we get

$$(18) \quad z \approx x^2 + \log(x^2) + \frac{2}{x^2} - \frac{5}{x^4} + \frac{74/3}{x^6} \dots \quad (x \rightarrow \infty)$$

By asymptotic iteration we find

$$(19) \quad x^2 \approx z - \log z + \sum_{k=1}^{\infty} z^{-k} p_k(\log z) \quad (z \rightarrow \infty)$$

where the  $p_k$ 's are polynomials of degree  $k$ . A few polynomials  $p_k$  are

$$(20) \quad \begin{aligned} p_1(t) &= t - 2 \\ p_2(t) &= \frac{1}{2} t^2 - 3t + 7 \\ p_3(t) &= \frac{1}{3} t^3 - \frac{3}{2} t^2 + 17t - \frac{107}{3} . \end{aligned}$$

The asymptotic expansions for  $x^j$  have the form

$$(21) \quad x^j \approx z^{j/2} \left( 1 + \sum_{k=1}^{\infty} p_{jk}(\log z) z^{-k} \right) \quad (z \rightarrow \infty)$$

where the  $p_{jk}$  are polynomials of degree  $k$ .

The individual terms in the asymptotic expansions (21) have the following property: Let  $f(z)$  be such a term occurring in the right side of (21).

Then  $f(z)$  is of the form

$$f(z) = (\log^m z) z^{-k+\frac{1}{2}j} \quad \text{where } m \leq k .$$

Let  $z_1 := 2 \log \frac{n}{\sqrt{2\pi}}$ . Clearly  $z_1 \geq 2$  if  $n \geq 7$ . Let  $n \geq 7$ . Then the power-series about  $z = z_1$

$$(22) \quad f(z_1 + \varepsilon) = \sum_{\ell=0}^{\infty} \frac{f^{(\ell)}(z_1)}{\ell!} \varepsilon^\ell$$

is convergent for  $|\varepsilon| < z_1$ . Now it is easily seen that this powerseries has the property that for all  $N \in \mathbb{N}$ ,  $N \geq \frac{1}{2}j - k$

$$(23) \quad f(z_1 + \varepsilon) = \sum_{\ell=0}^N \frac{f^{(\ell)}(z_1)}{\ell!} \varepsilon^\ell + O((\log^m z_1) z_1^{-k+\frac{1}{2}j-N-1} \varepsilon^{N+1})$$

( $-\frac{1}{2}z_1 < \varepsilon < \infty$ ,  $z_1 \geq 2$ ) .

Then it follows that upon substitution  $\varepsilon = -2 \log s$  in (22) we get an asymptotic expansion for  $0 < s < \log n$ ,  $n \rightarrow \infty$ , i.e. for all  $N \geq \frac{1}{2}j-k$

$$(24) \quad f(z_1 - 2 \log s) = \sum_{\ell=0}^N \frac{f^{(\ell)}(z_1)}{\ell!} (-2 \log s)^\ell + \\ + O((\log^m z_1) z_1^{-k+\frac{1}{2}j-N-1} (\log s)^{N+1}) \quad (0 < s < \log n, n \rightarrow \infty)$$

since  $2 \log \log n < \frac{1}{2}z_1$  for  $n$  sufficiently large. The hidden constant in the 0-term is independent of  $n$ .

Further, for every  $\ell \in \mathbb{N}$ ,

$$(25) \quad \int_{\log n}^{\infty} \log^\ell s e^{-s} ds = O(n^{-1} (\log \log n)^\ell) \quad (n \rightarrow \infty).$$

Therefore we can proceed as follows: In the asymptotic expansion (21) of  $x^j$  we substitute  $z = z_1 - 2 \log s$  and we expand formally into a power-series about  $z_1$ . After multiplication with  $e^{-s}$  and integration over  $(0, \infty)$  we get an asymptotic expansion for  $\mu_j(n)$ .

Denoting the asymptotic expansion (21) of  $x^j$  by  $X_j$  and writing  $X_j^{(k)}$  for its formal derivatives we have proved that

$$(26) \quad \mu_j(n) \approx \sum_{k=0}^n X_j^{(k)}(z_1) (-2)^k \Gamma^{(k)}(1) (k!)^{-1} \quad (n \rightarrow \infty),$$

where we have used that

$$(27) \quad \int_0^{\infty} e^{-s} \log^k s ds = \Gamma^{(k)}(1).$$

If we carry out the above program then, for instance, we find

$$(28) \quad \mu_1 = z^{1/2} - \frac{1}{2} z^{-1/2} \log z + \gamma z^{-1/2} - \frac{1}{8} z^{-3/2} \log^2 z + \\ + \frac{1}{2} (1+\gamma) z^{-3/2} \log z - (1-\gamma - \frac{1}{2} \gamma^2 - \frac{1}{12} \pi^2) z^{-3/2} + \\ + O(z^{-5/2} \log^3 z) \quad (n \rightarrow \infty)$$

where  $z = z_1$ . We have used that  $\Gamma'(1) = \gamma$  (Euler's constant) and  $\Gamma''(1) = \gamma^2 + \frac{1}{6} \pi^2$ .

Of course we can also find asymptotic results for  $\mu_2$ ,  $\mu_3$  and  $\mu_4$ . We will not do so since there is a more convenient way to obtain asymptotic expansions for  $\mu_j(n)$ . We shall show that  $\mu_j(n)$  has an asymptotic power-series expansion in powers of  $x_1^{-1}$ , where  $x_1$  is the value of  $x$  at  $z = z_1 := 2 \log \frac{n}{\sqrt{2\pi}}$ , i.e. there are sequences  $(C(j,k))_{k=-j}^{\infty}$  of real numbers such that

$$(29) \quad \mu_j(n) \approx \sum_{k=-j}^{\infty} C(j,k) x_1^{-k} \quad (n \rightarrow \infty) .$$

Considering  $x$  as a function of  $z$  defined by (17) we can write the integral in (8) as

$$(30) \quad \int_0^{\log n} x^j(z_1 - 2 \log s) e^{-s} ds .$$

We shall prove that we can find the asymptotic expansion of (30) by term-wise integration of the formal powerseries expansion of  $x^j(z_1 - 2 \log s)$  about  $z_1$ . We shall give the details of the proof for the case  $j = 1$ . The other cases  $j > 1$  can be treated analogously. So for the moment being we suppose  $j = 1$ . Obviously we are done with the problem if we have proved that

$$(31) \quad \forall_{k \in \mathbb{N}} \left( \frac{d^k x}{dz^k} \right)_1 \text{ has an asymptotic powerseries in } x_1^{-1} .$$

$$(32) \quad \forall_{k \in \mathbb{N}} \left( \frac{d^{k+1} x}{dz^{k+1}} \right)_1 = \sigma \left( \left( \frac{d^k x}{dz^k} \right)_1 \right) \quad (n \rightarrow \infty)$$

$$(33) \quad \forall_{N \in \mathbb{N}} \exists_{A > 0} x(z_1 - 2 \log s) = x_1 + \sum_{k=1}^N \left( \frac{d^k x}{dz^k} \right)_1 \frac{(-2 \log s)^k}{k!} + \\ + \mathcal{O} \left( \left( \frac{d^{N+1} x}{dz^{N+1}} \right)_1 (\log s)^{N+1} \right) \quad (0 < s < \log n, n > A)$$

where the subscript 1 in  $( )_1$  means the value at  $z_1$ .

From (17) it follows that

$$(34) \quad \frac{dx}{dz} = \frac{1}{2}a(x) ,$$

where

$$(35) \quad a(x) = \frac{1 - \Phi(x)}{\Phi'(x)} .$$

From (10) we see that

$$(36) \quad a(x) \approx \sum_{\ell=0}^{\infty} (-1)^{\ell} (2\ell-1)!! x^{-2\ell-1} \quad (x \rightarrow \infty) .$$

From (35) we derive that

$$(37) \quad \frac{da}{dx} = xa - 1 .$$

Clearly (36) and (37) imply that all derivatives  $d^k a/dx^k$  have asymptotic powerseries in  $x^{-1}$  which can be obtained by formal differentiation of the asymptotic series in (36). By means of the chain rule we can compute from (34) all derivatives  $dx^k/dz^k$ ; clearly,  $dx^k/dz^k$  is a sum of products involving  $a(x)$  and its derivatives  $d^{\ell} a/dx^{\ell}$ ,  $\ell = 1, 2, \dots, k-1$ . It follows that  $d^k x/dz^k$  has an asymptotic expansion in powers of  $x^{-1}$  for  $x \rightarrow \infty$ .

Using that

$$\frac{d^{\ell} a}{dx^{\ell}} \sim \frac{(-1)^{\ell} \ell!}{x^{\ell+1}} \quad (x \rightarrow \infty)$$

we easily derive that for  $k \geq 2$

$$(38) \quad \frac{d^k x}{dz^k} \sim (-1)^{k+1} (2k-3)!! 2^{-k} x^{-2k+1} \quad (x \rightarrow \infty) .$$

Thus we have proved (31) and (32).

Let  $N \in \mathbb{N}$ . Let  $h \in \mathbb{R}$ . Then there is a number  $\theta \in (0, 1)$  such that

$$(39) \quad x(z_1 + h) = x_1 + \frac{h}{1!} \left( \frac{dx}{dz} \right)_1 + \dots + \frac{h^N}{N!} \left( \frac{d^N x}{dz^N} \right)_1 + R_N$$

where

$$(40) \quad R_N = \frac{h^{N+1}}{(N+1)!} \frac{d^{N+1} x}{dz^{N+1}} (z_1 + \theta h) .$$

Restricting ourselves to  $h > -\frac{1}{2}z_1$ , we only have to prove that there is a number  $A > 0$  and a number  $K > 0$  such that for all  $z > A$

$$(41) \quad \left| \frac{d^{N+1}x}{dz^{N+1}}(\eta) \right| < K \left| \frac{d^{N+1}x}{dz^{N+1}}(z) \right| \quad (\eta > \frac{1}{2}z) .$$

On account of (38) and the fact that  $x(z) \sim z^{\frac{1}{2}}$  ( $z \rightarrow \infty$ ), (41) is obviously true. Hence (33) holds, since  $z_1 - 2 \log s > \frac{1}{2}z_1$  for  $0 < s < \log n$  and  $n$  sufficiently large.

Now using (33) in (30) for  $j = 1$  we get an asymptotic series in powers of  $x_1^{-1}$  for  $\mu_1(n)$ .

Analogously we can find asymptotic series for  $\mu_j(n)$ ,  $j = 2, 3, \dots$ . Only small adaptations are necessary; for instance in (33) we have to change  $\forall N \in \mathbb{N}$  into  $\forall N \in \mathbb{N}, N \geq \frac{1}{2}j$ .

REMARK. Especially for  $(\mu_3 - \mu_1\mu_2)(\mu_2 - \mu_1^2)^{-\frac{1}{2}}$  and  $(\mu_4 - \mu^2)^{\frac{1}{2}}$  the computations are most easily done if one postpones the replacement of  $d^k x^j / dz^k$  by its asymptotic series as long as possible. For instance, in this way we get

$$\mu_4 - \mu_2^2 = (8B - 4A^2)x^2(x')^2 + (24C - 8AB)(x^2x'x'' + x(x')^3) + \mathcal{O}(x^4)$$

where

$$A = -2\Gamma'(1), \quad B = 2\Gamma''(1) \quad \text{and} \quad C = -\frac{4}{3}\Gamma'''(1),$$

and  $x'$ ,  $x''$  denote first and second derivatives with respect to  $z$ . Now using the asymptotic series

$$x' = \frac{1}{2x} - \frac{1}{2x^3} + \dots, \quad x'' = -\frac{1}{4x^3} + \frac{1}{x^5} + \dots$$

we see that

$$x^2x'x'' + x(x')^3 = \mathcal{O}(x^{-4}) \quad \text{and} \quad x^2(x')^2 = \frac{1}{4} - \frac{1}{2x^2} + \dots$$

We get

$$\mu_4 - \mu_2^2 = d_0^2 - 2d_0^2x^{-2} + \mathcal{O}(x^{-4}),$$

from which we can easily derive the result in section 2.

## REFERENCE

- 1 BRANDS, J.J.A.M., Asymptotics in Poisson order statistics,  
Memorandum 86-03, Department of Mathematics,  
Eindhoven University of Technology,  
The Netherlands.