

On boundary value problems for ode with parameters

Citation for published version (APA):

Mattheij, R. M. M. (1987). *On boundary value problems for ode with parameters*. (Eindhoven University of Technology : Dept of Mathematics : memorandum; Vol. 8702). Technische Hogeschool Eindhoven.

Document status and date:

Published: 01/01/1987

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

EINDHOVEN UNIVERSITY OF TECHNOLOGY

Department of Mathematics and Computing Science

Memorandum 1987-02

October 1987

ON BOUNDARY VALUE PROBLEMS FOR ODE
WITH PARAMETERS

by

R.M.M. Mattheij

Eindhoven University of Technology
Department of Mathematics and Computing Science
P.O. Box 513
5600 MB Eindhoven
The Netherlands

ON BOUNDARY VALUE PROBLEMS FOR ODE WITH PARAMETERS

R.M.M. Mattheij

Eindhoven University of Technology
Department of Mathematics and Computing Science
P.O. Box 513
5600 MB Eindhoven
The Netherlands

ABSTRACT

When solving a boundary value problem for an ordinary differential equation with additional parameters, the usual stability and conditioning concepts induce a more complicated structure; in particular dichotomy of the (linearized) ODE is not necessarily implied. By using the adjoint problem and viewing this as an integral BVP it is shown that the so-called polychotomy of the latter induces a structure for the solution space of the original problem.

1. Introduction

In some applications problems of the following form are met: Let

$$\frac{dx}{dt} = f(x, t, \lambda), \quad (1.1)$$

$$\text{with } x : (0,1) \rightarrow \mathbb{R}^n,$$

be an ODE depending on a vector of parameters $\lambda := (\lambda_1, \dots, \lambda_m)^T$. For the $m + n$ unknowns there are $m + n$ boundary conditions (BC) specified:

$$g(x(0), x(1), \lambda) = 0, \quad (1.2)$$

$$\text{with } g : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^{n+m}.$$

Such problems may e.g. arise in chemistry where the unknown parameters are e.g. reaction rates or in geophysics, where they are wave speeds in different media. Often the BC are even

overdetermined, calling for a least squares solution, cf. [1]. Here we shall confine ourselves to the regular case, i.e. we assume that (1.1), (1.2) has an isolated solution. It is natural to think of the solution as the vector y , defined by

$$y(t) := \begin{bmatrix} x(t) \\ \lambda \end{bmatrix}. \quad (1.3)$$

Hence we may augment the ODE (1.1) by

$$\dot{\lambda} = 0. \quad (1.4)$$

In numerical computations one has to linearize the system (1.1) + (1.4), (1.2) at some stage of a practical algorithm (i.e. before or after discretization); hence one has to solve a sequence of linear BVP for which it makes sense to have knowledge about its conditioning and further structure. As has been shown in [2], linear two point BVP involve ODE having a dichotomic fundamental solution; for multipoint BVP a more complicated structure is present, cf. [4]. This has consequences for a numerical algorithm, in particular for the decoupling of the modes according to their growth behaviour in order to control the stability of the algorithm, cf. [3, 5]. In the sequel we shall show that parameter problems allow for yet another structure. As a consequence numerical methods for such problems should be based on a proper (different) decoupling technique.

2. Structure of the augmented system

When we linearize the ODE (1.1), (1.4) we obtain the following system

$$\frac{dx}{dt} = Lx + C\lambda + r(t) \quad (2.1a)$$

$$\frac{d\lambda}{dt} = 0, \quad (2.1b)$$

or equivalently using (1.3)

$$\frac{dy}{dt} = \hat{L}y + \hat{r}, \quad (2.1c)$$

Here $L(t) \in \mathbb{R}^{n \times n}$, $C(t) \in \mathbb{R}^{n \times m}$, $\hat{L}(t) \in \mathbb{R}^{(m+n) \times (m+n)}$, $r(t) \in \mathbb{R}^n$, $\hat{r}(t) \in \mathbb{R}^{n+m}$, are all assumed to be continuous at least.

The linearized BC can be written as

$$B_0x(0) + B_1x(1) + \tilde{B}_1\lambda = b \quad (2.2a)$$

$$K_0x(0) + K_1x(1) + \tilde{K}_1\lambda = c, \quad (2.2b)$$

or equivalently

$$\hat{B}_0 y(0) + \hat{B}_1 y(1) = \hat{b} . \quad (2.2c)$$

Here $B_0, B_1 \in \mathbb{R}^{n \times n}$, $K_i \in \mathbb{R}^{m \times n}$, $\bar{B}_1 \in \mathbb{R}^{n \times m}$, $\bar{K}_1 \in \mathbb{R}^{m \times m}$, $\hat{B}_0, \hat{B}_1 \in \mathbb{R}^{(n+m) \times (n+m)}$. Note that it is not restrictive to assume the last m columns of \hat{B}_0 to be zero. Finally, $b \in \mathbb{R}^n$, $c \in \mathbb{R}^m$, $\hat{b} \in \mathbb{R}^{n+m}$.

As is well known, for a well-posed problem, there exist constants κ_1, κ_2 , such that

$$\|y\| \leq \kappa_1 \|\hat{b}\| + \kappa_2 \|\hat{r}\| , \quad (2.3)$$

for appropriate norms (both vector- and function-). For *numerical relevance* these numbers κ_1, κ_2 should actually be *fairly moderate*. We can give an explicit expression for κ_2 based on the Green's function $\hat{G}(t, s)$.

Let $\hat{\Phi}$ be a fundamental solution of (2.1c), and define

$$\hat{Q} := \hat{B}_0 \hat{\Phi}(0) + \hat{B}_1 \hat{\Phi}(1) , \quad (2.4)$$

then

$$\hat{G}(t, s) = \hat{\Phi}(t) \hat{Q}^{-1} \hat{B}_0 \hat{\Phi}(0) \hat{\Phi}^{-1}(s) , \quad t > s ; \quad (2.5a)$$

$$\hat{G}(t, s) = -\hat{\Phi}(t) \hat{Q} \hat{B}_1 \hat{\Phi}(1) \hat{\Phi}^{-1}(s) , \quad t < s . \quad (2.5b)$$

If e.g. the BC are separated, i.e.

$$(2.6) \quad \hat{B}_0 = \begin{array}{|c|} \hline \text{diagonal lines} \\ \hline \emptyset \\ \hline \end{array} \quad \Downarrow I \quad \hat{B}_1 = \begin{array}{|c|} \hline \emptyset \\ \hline \text{diagonal lines} \\ \hline \end{array} \quad \Downarrow I$$

then the normalisation $\hat{Q} = I$ induces a projection P

$$P = \hat{B} \hat{\Phi}(0) = \begin{bmatrix} I_l & \emptyset \\ \emptyset & \emptyset \end{bmatrix} . \quad (2.7)$$

Hence well-conditioning implies

$$\|\hat{\Phi}(t) P \hat{\Phi}^{-1}(s)\| \leq \kappa_2 , \quad t > s \quad (2.8a)$$

$$\|\hat{\Phi}(t) (I - P) \hat{\Phi}^{-1}(s)\| \leq \kappa_2 , \quad t > s , \quad (2.8b)$$

i.e. the fundamental solution is dichotomic.

Remark 2.9.

In order to appreciate this result one has to realize that κ_2 is moderate indeed. On the other hand, by requiring such a constant κ_2 to be valid uniformly for an entire family of BVP (as in a singular perturbation context), it is clear that such statements can be put in a more mathematically precise framework, like by using asymptotics.

More generally we have

Theorem 2.10.

If the BVP (2.1c), (2.2c) is well-conditioned then there exists a dichotomic fundamental solution, i.e. satisfying (2.8) for some suitable projection P and moderate constant κ_2 .

Realizing that $\hat{\Phi}$ can be written as

$$\hat{\Phi}(t) = \begin{bmatrix} \Phi(t) & \emptyset \\ \int_0^t \Phi(s)ds & I \end{bmatrix} * E , \tag{2.11}$$

where Φ is a fundamental solution of E and E some nonsingular matrix, we see that dichotomy of $\hat{\Phi}$ does not imply dichotomy as such for Φ . We shall show that this is not true in general either.

3. Integral boundary conditions

In the next section we shall employ results that are known for general BC, including integral BC. As an upshot consider the multipoint BC

$$B_0 x(t_0) + B_1 x(t_1) + \dots + B_p x(t_p) = b , \quad t_0 = 0, \quad t_p = 1 . \tag{3.1}$$

By similar arguments as used in the two point case it is straightforward to see that separated BC induce projections and have a structure in the fundamental solution.

In particular we have in a well-conditioned situation that for some fundamental solution Φ and projections P_0, \dots, P_p , with $\sum_{j=0}^p P_j = I$, there exists a moderate constant κ such that

$$\| \Phi(t) \left[\sum_{j=0}^l P_j \right] \Phi^{-1}(s) \| \leq \kappa , \quad t_l < s < t_{l+1}, \quad t > s , \tag{3.2a}$$

$$\| \Phi(t) \left[\sum_{j=l+1}^p P_j \right] \Phi^{-1}(s) \| \leq \kappa , \quad t_l < s < t_{l+1}, \quad t < s . \tag{3.2b}$$

In [4] such a fundamental solution was called *polychotomic*.

Note that Φ is dichotomic intervalwise, but that this dichotomy may differ per interval. Such a polychotomy holds for fairly general BC, cf. [4], such as integral BC

$$\int_0^1 B(t) x(t) dt = b , \tag{3.3a}$$

or a combination

$$\int_0^1 B(t)x(t) dt + B_0x(0) + B_1x(1) = b . \quad (3.3b)$$

For interest of us is the form of the Green's function for (3.3b). Given the ODE

$$\frac{dx}{dt} = Lx , \quad (3.4)$$

we have

$$G(t,s) = \Phi(t)Q^{-1} \left\{ B_0\Phi(0) + \int_0^s B(\tau)\Phi(\tau) d\tau \right\} \Phi^{-1}(s) , \quad t > s \quad (3.5a)$$

$$G(t,s) = -\Phi(t)Q^{-1} \left\{ B_1\Phi(1) + \int_s^1 B(\tau)\Phi(\tau) d\tau \right\} \Phi^{-1}(s) , \quad t < s , \quad (3.5b)$$

where

$$Q := B_0\Phi(0) + B_1\Phi(1) + \int_0^1 B(\tau)\Phi(\tau) d\tau . \quad (3.6)$$

4. The adjoint of the parameter problem

As will turn out we can write the adjoint of our problem (2.1), (2.2) as a BVP with integral BC and hence relate its structure to that of our original problem. To see this we first consider the BC matrices \hat{B}_0, \hat{B}_1 and write them as an $(n+m) \times (2(n+m))$ matrix:

$$(4.1) \quad \begin{array}{|c|c|c|c|} \hline B_0 & \emptyset & B_1 & \tilde{B}_1 \\ \hline K_0 & \emptyset & K_1 & \tilde{K}_1 \\ \hline \end{array} .$$

It is not restrictive to assume that $[\hat{B}_0 | \hat{B}_1]$ has orthonormal rows. Hence we can construct matrices \hat{M}_0 and \hat{M}_1 such that

$$\begin{bmatrix} \hat{B}_0 & \hat{B}_1 \\ \hat{M}_0 & \hat{M}_1 \end{bmatrix} \quad (4.2a)$$

is orthogonal. It follows that $[\hat{M}_0 | \hat{M}_1]$ has the structure (where we have collected systematic zeros in a manner that suits our purposes):

$$(4.2b) \quad [\hat{M}_0 | \hat{M}_1] = \begin{array}{|c|c|c|c|} \hline M_0 & \emptyset & M_1 & L_1 \\ \hline \emptyset & L_0 & \emptyset & \emptyset \\ \hline \end{array} .$$

The (homogeneous) adjoint problem of (2.1c), (2.2c) then reads

$$\frac{dz}{dt} = -\hat{L}^T z \tag{4.3}$$

$$\hat{M}_0 z(0) - \hat{M}_1 z(1) = \hat{d} \quad (\text{for some } \hat{d} \in \mathbb{R}^{n+m}) . \tag{4.4}$$

Partitioning z and \hat{d} as

$$z = \begin{bmatrix} u \\ v \end{bmatrix} \begin{array}{l} \updownarrow n \\ \updownarrow m \end{array}, \quad \hat{d} = \begin{bmatrix} d_1 \\ d_2 \end{bmatrix} \begin{array}{l} \updownarrow n \\ \updownarrow m \end{array}, \tag{4.5}$$

we see that the system (4.3) is equivalent to

$$\frac{du}{dt} = -L^T u \tag{4.6a}$$

$$\frac{dv}{dt} = -C^T u . \tag{4.6b}$$

The latter ODE is in fact an integral BC for u :

$$v(1) = - \int_0^1 C(\tau)^T u(\tau) d\tau + v(0) . \tag{4.6c}$$

The BC (4.4) gives

$$L_0 v(0) = d_2 . \tag{4.7a}$$

Since L_0 must be nonsingular, we find via (4.6c) and (4.7a) the following BC for (4.6a):

$$M_0 u(0) - M_1 u(1) + \int_0^1 L_1 C^T(\tau) u(\tau) d\tau = d_1 + L_1 L_0^{-1} d_2 , \tag{4.7b}$$

a BC as in (3.3b).

We now first show that (4.6a), (4.7b) is well-conditioned if (4.3), (4.4) is:

Let Ψ be a fundamental solution of (4.6a), and let Ω be such that

$$\frac{d\Omega}{dt} = -C^T \Psi, \quad \Omega(0) = 0 . \tag{4.8}$$

Then obviously $\hat{\psi}$ defined by

$$\hat{\Psi} = \begin{bmatrix} \Psi & \emptyset \\ \Omega & I \end{bmatrix} \quad (4.9)$$

is a fundamental solution of (4.3). We obtain (cf. (2.4), (4.4))

$$(4.10) \quad \hat{Q} := \left[\begin{array}{c|c} M_0 \Psi(0) - M_1 \Psi(1) - L_1 \Omega(1) & -L_1 \\ \hline \emptyset & L_0 \end{array} \right] =: \begin{bmatrix} Q & -L_1 \\ \emptyset & L_0 \end{bmatrix}.$$

Denote the Green's function of (4.3), (4.4) by $\hat{H}(t, s)$; then it follows that the $n \times n$ left upper block, $H(t, s)$ say, equals

$$H(t, s) = \Psi(t) [Q^{-1} | Q^{-1} L_1 L_0^{-1}] \begin{bmatrix} M_0 \Psi(0) \Psi^{-1}(s) \\ L_0 \Omega(s) \Psi^{-1}(s) \end{bmatrix}, \quad (4.11a)$$

$$= \Psi(t) Q^{-1} \left\{ M_0 \Psi(0) + \int_0^s L_1 C^T(\tau) \Psi(\tau) d\tau \right\} \Psi^{-1}(s), \quad t > s,$$

$$H(t, s) = -\Psi(t) Q^{-1} \left\{ M_1 \Psi(1) + \int_s^1 L_1 C^T(\tau) \Psi(\tau) d\tau \right\} \Psi^{-1}(s), \quad t < s. \quad (4.11b)$$

Comparing (4.11) to (3.5) learns that $H(t, s)$ is precisely the Green's function of (4.6a), (4.7b). Now we have

$$\|H(t, s)\| \leq \|\hat{H}(t, s)\| \leq \max_{t,s} \|\hat{H}(t, s)\| \leq \max_{t,s} \|\hat{G}(t, s)\| \leq \kappa_2. \quad (4.12)$$

We therefore conclude from § 3 that Ψ is polychotomic.

We have

Property 4.13.

Let Φ, Ψ be fundamental solutions of (2.1a) and (4.6a) respectively. Then $\Psi(t)^T \Phi(t)$ is constant.

It is not restrictive to choose Φ such that

$$\Phi(t)^T \Psi(t) = I. \quad (4.14)$$

Property 4.15.

Let P_0, \dots, P_p be projections induced by the polychotomy of Ψ (cf. (3.2)). Then

$$\|\Phi(t) \left[\sum_{j=0}^l P_j \right] \Phi^{-1}(s)\|_2 = \|\Psi(s) \sum_{j=0}^l P_j \Psi^{-1}(t)\|_2.$$

Proof.

$$\Phi(t) \left[\sum_{j=0}^l P_j \right] \Phi^{-1}(s) = \Psi^{-T}(t) \left[\sum_{j=0}^l P_j \right] \Psi^{-T}(s).$$

Using the property $\|A^T\|_2 = \|A\|_2$, the result follows. □

Theorem 4.16.


If a linear problem with parameters is well-conditioned then there exist projections P_0, \dots, P_p and points $0 = t_0, \dots, t_p = 1$, such that for some moderate constant κ

$$\|\Phi(t) \left[\sum_{j=0}^l P_j \right] \Phi^{-1}(s)\| \leq \kappa, \quad t_l < s < t_{l+1}, \quad t < s, \tag{4.16a}$$

$$\|\Phi(t) \left[\sum_{j=l+1}^p P_j \right] \Phi^{-1}(s)\| \leq \kappa, \quad t_l < s < t_{l+1}, \quad t > s. \tag{4.16b}$$

We may call the property in (4.16) *skew polychotomy*. Roughly speaking it says that the solution space may be split into subspaces of modes that are nonincreasing till some point t_j after which they become nondecreasing. For clarity one may read for nonincreasing (nondecreasing): decreasing (increasing).

Below we summarize the three types of BVP and the solution types allowed by them:



References

- [1] H.G. Bock (1983), Recent advances in parameter identification techniques for ODE, in: Numerical Treatment of Inverse Problems in Differential and Integral Equations, (P. Deuffhard, E. Hairer eds.), Birkhäuser Boston, p. 99-121.
- [2] F.R. de Hoog, R.M.M. Mattheij (1987), On dichotomy and well-conditioning in BVP, SIAM J. Numer. Anal. 24, 89-105.
- [3] F.R. de Hoog, R.M.M. Mattheij (1987), An algorithm for solving multi-point boundary value problems, Computing 38, 219-234.
- [4] F.R. de Hoog, R.M.M. Mattheij (1987), On the conditioning of multipoint and integral boundary value problems, submitted for publication.
- [5] R.M.M. Mattheij (1985), Decoupling and stability of BVP algorithms, SIAM Rev. 27, 1-44.