

Decay of metastable Ne ($3P/2$)-atoms

Citation for published version (APA):

Dielis, J. W. H., Hoog, de, F. J., & Schram, D. C. (1979). Decay of metastable Ne ($3P/2$)-atoms. *Journal de Physique. Colloque*, 40(C7), 75-76.

Document status and date:

Published: 01/01/1979

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

DECAY OF METASTABLE Ne (³P₂)-ATOMS

J.W.H. Dielis, F.J. de Hoog and D.C. Schram.

Eindhoven University of Technology, Eindhoven, the Netherlands.

Investigations on the destruction rates of meta-stable atoms as a function of gas density in neon and the other noble gases have been carried out in recent years. The experimental method most frequently used is the optical absorption technique. (1,2,3, 4). A tunable dye laser was used in (5) to study non resonant fluorescence in the afterglow of a positive column in neon to determine the decay frequency of 3s-states. In another experiment a time resolved study of the vacuum U.V. emission of Ne (¹P₁) and Ne (³P₁) states was carried out to study the Ne (³P₂) decay (6). At this moment the numerous mechanisms which govern the ³P₂ decay are fairly well understood (1,5). In the present work the afterglow of a Townsend discharge (T.D.) is used for mass spectrometrical determination of the Ne (³P₂) decay at 77 and 295 K, using the Penning ionisation of N₂ impurities as a diagnostic reaction. Current densities smaller than 10⁻⁸A/cm² provide cumulative processes, e.g. dissociative recombination, not to take place.

Theory

Because of this low current density ³P₂-atoms are only destroyed by i) diffusion to the wall (D), ii) excitation with a ground state atom to the nearest ³P₁-state (a.A) and iii) three body collisions with two ground state atoms (γ). (See fig. 1). The ³P₂ decay frequency reads

$$v = \frac{D}{\Lambda^2} + \gamma N^2 + a.A.N. \left(1 - \frac{N_R}{a.N_M}\right) \quad (1)$$

with N_R and N_M the ³P₁ and ³P₂ densities, N the gas density, A the de-excitation frequency, a the ratio of excitation to de-excitation and Λ the diffusion length.

Method

The Ne (³P₂) decay rate is measured by using the Penning ionisation reaction Ne (³P₂) + N₂ → N₂⁺ + Ne + e⁻. The nitrogen density in the parent gas is so small that it does not affect the decay frequency. The rate of formation of N₂⁺ ions is proportional to the

³P₂-state density. The T.D. is pulsed from burning voltage to a lower voltage in the afterglow. N₂⁺-ions formed by the Penning reaction in the afterglow drift to the cathode which contains a small hole (10 μm). The ions are sampled, selected on mass by a Q-pole mass filter and are detected by a channeltron. The times of arrival of the N₂⁺-ions after onset of the afterglow are processed by a microprocessor. Repeatedly pulsing gives a histogram of arrival times of N₂⁺, identical to the relative density of the ³P₂-atoms as a function of time in the afterglow and hence yield the decay frequency. The T.D., placed in a cryostat, can be cooled to liquid nitrogen temperature of 77 K. Cataphoresis of the research grade neon gas provides the impurity degree to be less than a few p.p.m. (Fig.2: exp.)




Fig 1: Diagram for the decay of Ne(³P₂) atoms.

Results

The measured decay frequencies as a function of gas density at 77 and 295 K are shown in fig. 3. By means of a nonlinear least mean square procedure eq. (1) is fitted to the data, obtaining the best values for D, A and γ, as given below

Temperatùre	295 K	77 K
D(10 ²⁰ m ⁻¹ s ⁻¹)	4.8 ± 0.3	2.3 ± 0.2
A(10 ²⁰ m ³ s ⁻¹)	3.2 ± 0.1	0
γ(10 ⁻⁴⁶ m ⁶ s ⁻¹)	5.4 ± 0.3	0.52 ± 0.04

Diffusion coefficient

As can be seen in fig. 4 the agreement of the present data at 295 K is good, whereas at 77 K the present value is 25% larger than previous results. From potential curve considerations, showing a large insensibility of D on the shape and parameter values of the ($^3P_2 - ^1S_0$) interaction, this deviation is not disquieting.

De-excitation rate

Experimental (1,5,6,10) and theoretical (11) results on A are shown in fig. 5. The present result at 295 K is 50% smaller than previous data, but in good agreement with the theory of Cohen et al (11).

Three-body collision coefficient

The present results on γ at 77 and 300 K are in good agreement with previous experiments. The activation energy for this process is calculated to be 0.032 eV, in conformity with (12) and (13).




Fig 2: Experimental setup and measuring system.

Acknowledgements

The authors thank L.A.W.Gaykema and M.J.P.Vetjens for their valuable contributions and D.C.Schram and A.A.Kruithof for helpful discussions.

References

- (1) Phelps A.V., Phys. Rev. 114 (1011) 1959.
- (2) Phelps A.V. et al, Phys. Rev. 89 (1202) 1953.
- (3) Dixon J.R. et al, Phys. Rev. 107 (118) 1957.
- (4) Grant J.R. et al, Phys. Rev. 90 (59) 1953.
- (5) Steenhuysen L.W.G. et al, Proc. XIII ICPIG Berlin p. 37, 1977.
- (6) Leichner P.K. et al, Phys. Rev. A12 (2501) 1975.
- (7) Metaxas A.C. et al, J. Phys. D5 (1077) 1972.
- (8) Grant F.A., Phys. Rev. 84 (844) 1951.
- (9) Molnar J.P., Phys. Rev. 83 (945) 1951.
- (10) Biondi M.A., Phys. Rev. 88 (660) 1952.
- (11) Cohen J.S. et al, Phys. Rev. A17 (1343) 1978.
- (12) Cohen J.S. et al, J. Chem. Phys. 61 (3230) 1974.
- (13) Gritsyna V.V., Opt. Comm. 10 (320) 1974.




Fig 3: Decay frequency of Ne(3P_2) vs. gas density at 77 and 295 K
solid curve: least square fit of eq.1




Fig 4: Diffusion coefficient vs. gas temperature. 1(\circ), 4(Δ), 5(\square), 7(∇), 3(\bullet), 8(\blacktriangle), 9(\blacksquare), 10(\blacktriangledown), 2(\diamond), present(\circ).
Solid curve: theory with 16/6-potential.




Fig 5: De-excitation rate vs. gas temperature. 1(\circ), 4(Δ), 10(∇), 5(\square), 6(\blacktriangledown), present(\circ).
solid curve: theory/12/.