

# Classification of singularities at infinity of polynomials of degree 4 in two variables

**Citation for published version (APA):**

Siersma, D., & Smeltink, J. W. (1996). *Classification of singularities at infinity of polynomials of degree 4 in two variables*. (Rijksuniversiteit Utrecht. Mathematisch Instituut : preprint; Vol. 945). Utrecht University.

**Document status and date:**

Published: 01/01/1996

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

# Classification of singularities at infinity of polynomials of degree 4 in two variables

Dirk Siersma, Job Smeltink

february 6, 1996

## Introduction

During the last years there is an increasing interest in the behaviour of polynomials at infinity. In studying the family of levels curves

$$f(x, y) = t$$

one wants to know e.g the topological type of generic fibres, the set of bifurcation values, the change of topology of the fibre near bifurcation values, the monodromy at infinity, the intersection form on fibres.

For classes of polynomials (e.g. tame polynomials) the changes in the topology depend only on the affine singular points and their singularity types. But in general also other effects (due to the non properness of the polynomial function) can change the fibres. One calls this 'singular behaviour at infinity'.

Examples always played an important role in the investigation of germs of isolated hypersurface singularities. From there several effects, such as characterizations of simplicity, strange duality, etc were discovered. This is the reason that we started also in the global case to study examples in a systematic way and to classify polynomials of low degree  $d$  and a small number of variables  $n$ . In this note we give the classification in the case  $n = 2$  and  $d = 4$ , continuing the work of C.T.C Wall [Wal2] in case  $n = 2$  and  $d = 3$ .

We recall first a usual technique for studying singularities at infinity.

Let  $f(x, y)$  be a polynomial of degree  $d$ . We can write:  $f = f_0 + f_1 + \dots + f_d$  with  $f_d \neq 0$ , the decomposition in graded pieces.

We define  $\bar{f}(x, y, z) = z^d f(\frac{x}{z}, \frac{y}{z}) = f_d + z f_{d-1} + z^2 f_{d-2} + \dots + z^{d-1} f_1 + z^d f_0$  and

$$F(x, y, z; t) = \bar{f}(x, y, z) - tz^d$$

The equation  $F(x, y, z; t) = 0$  defines for each  $t$  a curve in  $P^2(\mathbb{C})$ .

Consider  $\bar{X} = \{(x, t) \in P^2(\mathbb{C}) \times \mathbb{C} \mid F(x, y, z; t) = 0\}$ .  $\bar{X}$  is the closure of the graph  $X$  of  $f$ .

Let  $t : \bar{X} \rightarrow \mathbb{C}$  be the projection on the second factor; the map  $t$  is an extension of  $f$  to the compactification  $\bar{X}$  of  $X$ . Hà-Lê [HL] used the stratification theory of this algebraic map to show that there is only a finite set of points in  $\mathbb{C}$ , above which  $t : \bar{X} \rightarrow \mathbb{C}$  fails to be a (stratified) local trivial fibre bundle. The smallest set in  $\mathbb{C}$ , above which  $f : \mathbb{C}^2 \rightarrow \mathbb{C}$  fails to be a local trivial fibre bundle is called the bifurcation set of  $f$  and is denoted by  $B = B_f$ . The fibres  $\bar{X}_t$  of  $t$  are defined by  $F(x, y, z; t) = 0$  with  $t$  fixed; the fibers of  $f$  are denoted by  $X_t$  and are defined by  $f(x, y) = t$ .

Remark that the intersection of  $\bar{X}$  with infinity ( $z = 0$ ) is given by the equation

$$f_a(x, y) = 0,$$

which defines a finite set of points  $A$  in  $\mathbb{P}^1(\mathbb{C})$ . This intersection is independent of  $t$  and  $\bar{X}$  intersects infinity in the space  $A \times \mathbb{C}$ .

We consider the family of functions

$$F_t(x, y, z) := F(x, y, z, t)$$

near each of the points  $a$  in  $A$ .

In case that  $f$  has only isolated singularities (in the affine plane) these are families of isolated hypersurface singularities. For generic values of  $t$  there is a well-defined  $\mu$ -class with Milnor number  $\mu_a^\infty$ . For a finite number of bifurcation values  $t$  this type can change and the Milnor number can drop with a value  $\lambda_a^t = \mu(\bar{X}_t, a) - \mu_a^\infty$ . In case that  $f$  has non-isolated singularities then the family of isolated singularities will jump at that critical value to a non-isolated germ at the corresponding intersection point at infinity.

Let  $\mu_p(f)$  be the local Milnor number of  $f$  at the point  $p \in \mathbb{C}$ . (For regular points, this is zero).

**Definition:** *global Milnor number:*

$$\mu(f) = \sum \mu_p(f)$$

**Definition:** *total jump:*

$$\lambda(f) = \sum_{t \in B} \sum_{a \in A} \lambda_a^t(f)$$

We now recall the global bouquet theorem as studied by Broughton [Br], Hà-Lê [HL] and in general form by Siersma-Tibar [SiTi]:

**Theorem:** *Let  $f : \mathbb{C}^2 \rightarrow \mathbb{C}$  be a polynomial with isolated singularities. Then all fibres of  $f$  have the homotopy type of a bouquet of 1-spheres and the number  $\gamma$  of spheres in a general fibre is equal to the sum  $\mu(f) + \lambda(f)$ , where  $\mu(f)$  is the total Milnor number of  $f$  and  $\lambda(f)$  is the sum of the jumps at infinity.*

*In particular  $\lambda(f)$  is invariant under diffeomorphisms of  $\mathbb{C}^n$ .*

*Moreover let  $\mu^b(f)$  denote the sum of the Milnor numbers of all the singularities of the fibre  $X_b$  and let  $\lambda^b(f)$  denote the sum of all Milnor numbers at infinity at  $A \times \{b\}$ . Then*

$$\chi(X_t) - \chi(X_b) = (-1)^{n-1}(\lambda^b(f) + \mu^b(f)),$$

where  $X_t$  is a general fibre of  $f$ .

**Remark:** There is a second formula (cf. eg [HL]), which computes the first Betti-numbers of the fibres. For the generic fibre this formula is:

$$\gamma = b_1(X_t) = d^2 - 3d + 1 - \sum_{a \in A} (\mu_a^\infty - 1)$$

where the sum runs over all points of the intersection with infinity.

List of isolated singularities: (cf [AVG], [Sie1])

<i>type</i>	<i>f</i>	<i>conditions</i>
$A_k$	$x^2 + y^{k+1}$	$k \geq 0$
$D_k$	$x^2y + y^{k-1}$	$k \geq 4$
$E_6$	$x^3 + y^4$	
$E_7$	$x^3 + xy^3$	
$E_8$	$x^3 + y^5$	
$X_9 = \tilde{E}_7$	$x^4 + y^4 + ax^2y^2$	$a^2 \neq 4$
$X_{10}$	$x^4 + y^5 + ax^2y^2$	$a \neq 0$
$J_{10} = \tilde{E}_8$	$x^3 + y^6 + axy^4$	$4a^3 + 27 \neq 0$

List of non-isolated singularities: (cf [Sie2], [dJ])

<i>type</i>	<i>f</i>	<i>transversal type</i>
$A_\infty$	$y^2$	$A_1$
$D_\infty$	$xy^2$	$A_1$
$J_{k,\infty}$ ( $k \geq 2$ )	$y^3 + x^ky^2$	$A_1$
$T_{\infty,k,2}$ ( $k \geq 4$ )	$x^2y^2 + y^k$	$A_1$
$T_{\infty,\infty,2}$	$x^2y^2$	$A_1$
$F_1A_2$	$xy^3$	$A_2$
$\mathbf{T}A_k$	$y^{k+1}$	$A_k$

In this list the singular locus is a smooth 1-dimensional subvariety, the  $x$ -axis. These singularities are called line singularities. The transversal type is the type of the singularity if one restricts  $f$  to  $H_c = \{x = c\}$  with  $c \neq 0$ .

A complete study of the local behaviour of non-isolated singularities of functions of two variables is given in [Sch], [Sie2] and [dJ].

**Conventions:** We will use the following notations:

$f \simeq g$ : in case  $f$  is (algebraic) diffeomorphic to  $g$ .

$f \cong g$ : in case  $f$  is diffeomorphic to  $g$  by an affine linear transformation.

We both use transformations in the source and target spaces. Examples are translations

$\mathbb{T}_{\alpha,\beta}$ :

$$\begin{cases} x := x - \alpha \\ y := y - \beta \end{cases}$$

and similar ones for the target. Since we are studying all fibres  $f(x, y) = t$ , we can assume that the constant term  $f_0 = 0$  and consider the actions modulo constants.

In all cases we consider first the factorisation of  $f_d$  in its linear parts and put  $f_d$  into a normal form using linear transformations of the source.

We use the following notation:

$\mathbf{A}^2\mathbf{BC}$  means that  $f_d$  has one double factor and two single factors at infinity, etc. The notation  $\mathbf{A}^3\mathbf{B}_{\gamma,\lambda}$  means that the singularities in this class have first betti number  $\gamma$  and total jump  $\lambda$ . In case  $\lambda = 0$  we omit the second number, e.g  $\mathbf{A}^3\mathbf{B}_6$ . For jumps to non-isolated singularities we use lower case stars, e.g.  $\mathbf{A}^2\mathbf{B}_{1,*}$ . The type is well-defined up to affine linear transformations.

For each singularity class we select one of its members with few monomials as its ‘typical’ form.

We next mention the types of the singular point at infinity. The notation  $A_2 \rightarrow A_3$  means that the singularity (at infinity) jumps from  $A_2$  to  $A_3$  for some bifurcation value.

The classification in degree 1 and 2 are classical. Degree 3 already has been studied by Wall [Wal2]. But we have redone this part to conform it to our notations.

**Remark:** We found the following interesting behaviour in the list. The singularity types  $\mathbf{A}^2\mathbf{BC}_{5,\lambda}$  occur for  $\lambda = 1, 2, 3, 4$  all with the same homotopy type of the general fibre. So one should consider the family

$$f = x^2(x^2 - y^2) + Kx^3 + Px^2 + Qxy + Ax = t$$

depending on  $K, P, Q, A$  and  $t$  and study its behaviour. Since  $\mu + \lambda$  is constant, some critical point is expected to go to infinity if  $\lambda$  increases. What is the effect on monodromy, etc ? The same feature occurs in other cases, and in all those examples  $\lambda \geq 1$ .

## Two variables, degree 1

Case **A**:  $f = x$

type	$f_1 = x$				
<b>A</b>	$f = x$				
	'typical' form	$(0 : 1 : 0)$	$\mu$	$\lambda$	$\gamma$
<b>A</b> <sub>0</sub>	$x$	$A_0$	0	0	0

## Two variables, degree 2

Case **AB**:  $f_2 = xy$

type	$f_2 = xy$				
<b>AB</b>	$f = xy$				
	'typical' form	two points at $\infty$	$\mu$	$\lambda$	$\gamma$
<b>AB</b> <sub>1</sub>	$xy$	$A_0, A_0$	1	0	1

Case **A<sup>2</sup>**:  $f_2 = x^2$

type	$f_2 = x^2$						
<b>A<sup>2</sup></b>	$f = x^2 + By$						
	'typical' form	$(0 : 1 : 0)$	$\neq 0$	$= 0$	$\mu$	$\lambda$	$\gamma$
<b>A<sup>2</sup></b> <sub>0</sub> $\simeq$ <b>A</b> <sub>0</sub>	$x^2 + y$	$A_0$	$B$		0	0	0
<b>A<sup>2</sup></b> <sub>*</sub>	$x^2$	$A_1 \rightarrow A_\infty$		$B$	$\infty$	*	
All jumps occur at $t = 0$							

## Two variables, degree 3

With homogeneous coordinates we can choose 3 point freely at the line of infinity which determine the whole coordinate system. Therefore  $f_3$  can be transformed into one of the following cases:

$$f_3 = xy(x + y)$$

$$f_3 = x^2y$$

$$f_3 = x^3$$

Case **ABC**:  $f_3 = xy(x + y)$

We get that

$$f(x, y) = xy(x + y) + Kx^2 + Lxy + My^2 + Px + Qy$$

By using translations  $\mathbb{T}_{M,K}$  we can assume  $K = M = 0$ .

The intersection points with infinity are  $(1 : 0 : 0)$ ,  $(0 : 1 : 0)$  and  $(1 : -1 : 0)$ . In all the cases we find that

the intersection points are nonsingular, so we have type  $A_0$ , which is always the case for non multiple factors in  $f_d$ .

type	$f_3 = xy(x+y)$				
<b>ABC</b>	$f = xy(x+y) + Lxy + Px + Qy$				
	'typical' form	three points at $\infty$	$\mu$	$\lambda$	$\gamma$
<b>ABC<sub>4</sub></b>	$x^3 + y^3$	$A_0, A_0, A_0$	4	0	4

**Case A<sup>2</sup>B:**  $f_3 = x^2y$

Let's look at the general form:

$$f(x, y) = x^2y + Kx^2 + Lxy + My^2 + Px + Qy$$

By using translations  $\mathbb{T}_{\frac{L}{2}, K}$  we can assume that  $L = K = 0$ : The intersection points with infinity are  $(0 : 1 : 0)$  and  $(1 : 0 : 0)$ .

In the case of  $(1 : 0 : 0)$  we find that the intersection point is always nonsingular, so type  $A_0$ .

The point  $(0 : 1 : 0)$  is singular iff  $M = 0$ .

Let's assume that  $M = 0$ , then the corresponding hypersurface  $F(x, y, z, t) = 0$  is given by:

$$F = x^2y + Pxz^2 + Qyz^2 - tz^3$$

On the chart  $y = 1$ , we have

$$F = x^2 + Pxz^2 + Qz^2 - tz^3$$

If  $Q \neq 0$  we have  $F \simeq x^2 + z^2$  ( $A_1$ ).

If  $Q = 0$  we can use the translation

$$\begin{cases} x := x - \frac{1}{2}Pz^2 \\ z := z \end{cases}$$

to get

$$F \simeq x^2 - tz^3 - \frac{1}{4}P^2z^4$$

So we find that if  $t \neq 0$ , then we have type  $A_2$ .

If  $t = 0$  and  $P \neq 0$ , this gives type  $A_3$ .

If  $t = P = 0$ , we have type  $A_\infty$ .

type	$f_3 = x^2y$							
<b>A<sup>2</sup>B</b>	$f = x^2y + My^2 + Px + Qy$							
	'typical' form	$(0 : 1 : 0)$	$(1 : 0 : 0)$	$\neq 0$	$= 0$	$\mu$	$\lambda$	$\gamma$
<b>A<sup>2</sup>B<sub>3</sub></b>	$x^2y + y^2$	$A_0$	$A_0$	$M$		3	0	3
<b>A<sup>2</sup>B<sub>2</sub></b>	$x^2y + y$	$A_1$	$A_0$	$Q$	$M$	2	0	2
<b>A<sup>2</sup>B<sub>1,1</sub></b>	$x^2y + x$	$A_2 \rightarrow A_3$	$A_0$	$P$	$M, Q$	0	1	1
<b>A<sup>2</sup>B<sub>1,*</sub></b>	$x^2y$	$A_2 \rightarrow A_\infty$	$A_0$		$M, Q, P$	$\infty$	*	1
All jumps occur at $t = 0$ .								

**Case A<sup>3</sup>:**  $f_3 = x^3$

Let's look at the general form

$$f(x, y) = x^3 + Kx^2 + Lxy + My^2 + Px + Qy$$

By using the translation  $\mathbb{T}_{\frac{K}{3}, 0}$  we can assume that  $K = 0$ .

The intersection point at infinity is  $(0 : 1 : 0)$ .

So the corresponding hypersurface  $F(x, y, z, t) = 0$  is given by

$$F = x^3 + Lxyz + My^2z + Pxz^2 + Qyz^2 - tz^3$$

On chart  $y = 1$  we have

$$F = x^3 + Lxz + Mz + Pxz^2 + Qz^2 - tz^3$$

So the intersection point is singular iff  $M = 0$ .

If  $L \neq 0$ , we have corank 0, so type  $A_1$ . So we can assume that  $L = 0$ .

If  $Q \neq 0$ , we have type  $A_2$ , because of the influence of  $x^3$ .

Consider the case  $M = L = Q = 0$ .

We have

$$F = x^3 + Pxz^2 - tz^3$$

If  $P \neq 0$ , we have type  $D_4$  with a jump to  $D_\infty$  if  $t^2 = -\frac{4P^3}{27}$ . So there are two jumps.

In the case  $P = 0, t \neq 0$ , we have  $D_4$ . If  $P = t = 0$ , we have  $F = x^3$  this type is called  $\mathbf{TA}_2$ . So if  $P = 0$ , we get a jump from  $D_4$  to  $\mathbf{TA}_2$  at  $t = 0$ .

type	$f_3 = x^3$						
$\mathbf{A}^3$	$f = x^3 + Lxy + My^2 + Px + Qy$						
	'typical' form	$(0 : 1 : 0)$	$\neq 0$	$= 0$	$\mu$	$\lambda$	$\gamma$
$\mathbf{A}^3_2$	$x^3 + y^2$	$A_0$	$M$		2	0	2
$\mathbf{A}^3_1 \simeq \mathbf{AB}_1$	$x^3 + xy$	$A_1$	$L$	$M$	1	0	1
$\mathbf{A}^3_0 \simeq \mathbf{A}_0$	$x^3 + y$	$A_2$	$Q$	$M, L$	0	0	0
$\mathbf{A}^3_*$	$x^3 + x$	$D_4 \rightarrow D_\infty$	$P$	$M, L, Q$	$\infty$	*	
$\mathbf{A}^3_{**}$	$x^3$	$D_4 \rightarrow \mathbf{TA}_2$		$M, L, Q, P$	$\infty$	*	
All jumps occur if $t^2 = -\frac{4P^3}{27}$							
Remark that in the last case $P = 0$ , so the jump occurs at $t = 0$							

## Two variables, degree 4

Like degree 3 we can use linear transformations to transform  $f_4$  over  $\mathbb{C}$  into one of the following cases:

$$f_4 = x^4 + y^4 + ax^2y^2, \quad \text{with } a^2 \neq 4$$

$$f_4 = x^2(x^2 - y^2)$$

$$f_4 = x^2y^2$$

$$f_4 = x^3y$$

$$f_4 = x^4$$

**Case ABCD:**  $f_4 = x^4 + y^4 + ax^2y^2$

All the intersectionpoints at infinity are regular (type  $A_0$ ).

type	$f_4 = x^4 + y^4 + ax^2y^2$				
<b>ABCD</b>	$f = f_4 + Lxy^2 + Mxy^2 + Px^2 + Qxy + Ry^2 + Ax + By$				
	'typical' form	four points at $\infty$	$\mu$	$\lambda$	$\gamma$
<b>ABCD</b> <sub>9</sub>	$x^4 + y^4$	$A_0, A_0, A_0, A_0$	9	0	9



**Case A<sup>2</sup>BC:**  $f_4 = x^2(x^2 - y^2)$

First we consider

$$f_3 = Kx^3 + Lx^2y + Mxy^2 + Ny^3$$

By using translations  $\mathbb{T}_{\frac{M}{2}, \frac{L}{2}}$  we can assume that  $L = M = 0$ . Remark that we could have put  $K$  instead of  $M$  zero.

So we get

$$f = x^2(x^2 - y^2) + Kx^3 + Ny^3 + Px^2 + Qxy + Ry^2 + Ax + By$$

The intersection points at infinity are  $(0 : 1 : 0)$ ,  $(1 : 1 : 0)$  and  $(1 : -1 : 0)$ .

The corresponding hypersurface  $F(x, y, z, t) = 0$  is given by

$$F = x^2(x^2 - y^2) + Kx^3z + Ny^3z + Px^2z^2 + Qxyz^2 + Ry^2z^2 + Axz^3 + Byz^3 - tz^4$$

On chart  $x = 1$  we get that the points  $(1 : 1 : 0)$  and  $(1 : -1 : 0)$  are always regular ( $A_0$ ).

To study  $(0 : 1 : 0)$  we consider the chart  $y = 1$ :

$$F = x^4 - x^2 + Kx^3z + Nz + Px^2 + Qxz^2 + Rz^2 + Axz^3 + Bz^3 - tz^4$$

So the point  $(0 : 1 : 0)$  is singular iff  $N = 0$ . If  $R \neq 0$  we have type  $A_1$ .

Next we assume that  $N = R = 0$ .

We have now

$$F = x^4 - x^2 + Kx^3z + Px^2z^2 + Qxz^2 + Axz^3 + Bz^3 - tz^4$$

If  $B \neq 0$ , we have type  $A_2$ .

In the case  $B = 0$  we have

$$F = -(x - \frac{1}{2}Qz^2)^2 + z^4(-t + \frac{1}{4}Q^2) + Axz^3 + Px^2z^2 + x^4 + Kx^3z$$

So if  $t \neq \frac{Q^2}{4}$ , we have type  $A_3$ .

If  $B = 0$  and  $t = \frac{Q^2}{4}$ , we get a jump to  $A_4$  iff  $AQ \neq 0$ .

So let's assume that  $AQ = 0$ . We split this into two cases:

- 1)  $Q \neq 0$
- 2)  $Q = 0$

1)  $Q \neq 0$ , so  $A = 0$ . In this case we have type  $A_3$  with a jump to  $A_5$  at  $t = \frac{Q^2}{4}$  iff  $PQ^2 \neq 0$  which is equivalent to  $P \neq 0$ .

If  $P = 0$  and  $K \neq 0$ , we have type  $A_3$  with a jump to  $A_6$  when  $t = \frac{Q^2}{4}$  and if  $P = K = 0$  we have type  $A_3$  with a jump at  $t = \frac{Q^2}{4}$  to  $A_7$ , because  $Q \neq 0$ .

2)  $Q = 0$ . We get

$$F = -x^2 + Axz^3 + x^4 + Px^2z^2 + Kx^3z - tz^4$$

So if  $A \neq 0$ , this gives type  $A_3$  with a jump to  $A_5$  at  $t = 0$ .

In case  $A = 0$ , we have

$$F = x^2(-1 + x^2 + Pz^3 + Kxz) - tz^4$$

so we have type  $A_3$  with a jump to  $A_\infty$  when  $t = 0$ .

type	$f_4 = x^2(x^2 - y^2)$								
$A^2BC$	$f = x^2(x^2 - y^2) + Kx^3 + Ny^3 + Px^2 + Qxy + Ry^2 + Ax + By$								
	'typical' form	$(0 : 1 : 0)$	$(1 : -1 : 0)$	$(1 : 1 : 0)$	$\neq 0$	$= 0$	$\mu$	$\lambda$	$\gamma$
$A^2BC_8$	$x^4 - x^2y^2 + y^3$	$A_0$	$A_0$	$A_0$	$N$		8	0	8
$A^2BC_7$	$x^4 - x^2y^2 + y^2$	$A_1$	$A_0$	$A_0$	$R$	$N$	7	0	7
$A^2BC_6$	$x^4 - x^2y^2 + y$	$A_2$	$A_0$	$A_0$	$B$	$N, R$	6	0	6
$A^2BC_{5,1}$	$x^4 - x^2y^2 + Qxy + Ax$	$A_3 \rightarrow A_4$	$A_0$	$A_0$	$Q, A$	$N, R, B$	4	1	5
$A^2BC_{5,2}$	$x^4 - x^2y^2 + Qxy + Px^2$	$A_3 \rightarrow A_5$	$A_0$	$A_0$	$Q, P$	$N, R, B, A$	3	2	5
$A^2BC_{5,3}$	$x^4 - x^2y^2 + Qxy + Kx^3$	$A_3 \rightarrow A_6$	$A_0$	$A_0$	$Q, K$	$N, R, B, A, P$	2	3	5
$A^2BC_{5,4}$	$x^4 - x^2y^2 + Qxy$	$A_3 \rightarrow A_7$	$A_0$	$A_0$	$Q$	$N, R, B, A, P, K$	1	4	5
$A^2BC_{5,2}^*$	$x^4 - x^2y^2 + x$	$A_3 \rightarrow A_5$	$A_0$	$A_0$	$A$	$N, R, B, Q$	3	2	5
$A^2BC_{5,*}$	$x^4 - x^2y^2$	$A_3 \rightarrow A_\infty$	$A_0$	$A_0$		$N, R, B, Q, A$	$\infty$		5

All jumps occur if  $t = \frac{Q^2}{4}$ .  
Remark that in the last two cases  $Q = 0$ , so the jump occurs at  $t = 0$ .

NB: The two cases  $A^2BC_{5,3}$  and  $A^2BC_{5,3}^*$  are connected !

**Case  $A^2B^2$ :**  $f_4 = x^2y^2$

First we consider

$$f_3 = Kx^3 + Lx^2y + Mxy^2 + Ny^3$$

By using translations  $\mathbb{T}_{\frac{M}{2}, \frac{L}{2}}$  we can assume  $L = M = 0$

There are two intersection points with infinity  $(1 : 0 : 0)$  and  $(0 : 1 : 0)$ .

If  $K = 0$  then  $(1 : 0 : 0)$  is a singular point and

if  $N = 0$  then  $(0 : 1 : 0)$  is a singular point.

Consider the general form:

$$f = x^2y^2 + Kx^3 + Ny^3 + Px^2 + Qxy + Ry^2 + Ax + By$$

on which we have still the possibility of permuting our coordinates  $x$  and  $y$  and multiplication with constants.

The corresponding hypersurface  $F(x, y, z, t) = 0$  is given by:

$$F = x^2y^2 + Kx^3z + Ny^3z + Px^2z^2 + Qxyz^2 + Ry^2z^2 + Axz^3 + Byz^3 - tz^4$$

First we consider the case that  $N \neq 0$ , where  $(0 : 1 : 0)$  is nonsingular, but later we'll use the same computations in case  $N = 0$ .

On the chart  $x = 1$  we have:

$$F = y^2 + Kz + Ny^3z + Pz^2 + Qyz^2 + Ry^2z^2 + Az^3 + Byz^3 - tz^4$$

We are going to discuss the type of  $F$  (depending on  $t$ ) at  $(0, 0)$ :

- 0) If  $K \neq 0$  we have the generic case: type  $A_0$  : no jumps.  
 1) If  $K = 0$  and  $P \neq 0$ : type  $A_1$  : no jumps.  
 2) If  $K = P = 0$  and  $A \neq 0$ : type  $A_2$ : no jumps.  
 3) Consider now  $K = P = A = 0$ :

$$F = y^2 + Ny^3z + Qyz^2 + Ry^2z^2 + Byz^3 - tz^4$$

If  $t \neq -Q^2/4$  we have type  $A_3$ , but if  $t = -Q^2/4$  we get a jump to type  $A_4$  if  $BQ \neq 0$ , see the next computation:

Set

$$y := y - Q^2/2$$

then we get

$$F \simeq y^2 + Ny^3z - \frac{3}{2}NQy^2z^3 + \frac{3}{4}NQ^2yz^5 - \frac{1}{8}NQ^3z^7 + Ry^2z^2 - QRyz^4 + \frac{1}{4}RQ^2z^6 + Byz^3 - \frac{1}{2}BQz^5$$

The next step splits into two cases:

4a)  $K = P = A = B = 0$ :

$$F = y^2 + Ny^3z + Qyz^2 + Ry^2z^2 - tz^4$$

If  $t \neq -Q^2/4$  we have type  $A_3$ , but if  $t = -Q^2/4$  we get a jump to type  $A_5$  if  $QR \neq 0$  (use the same substitution as above with  $B = 0$ ).

4b)  $K = P = Q = A = 0$ :

$$F = y^2 + Ny^3z + Ry^2z^2 + Byz^3 - tz^4$$

If  $t \neq 0$  we have type  $A_3$ , but if  $t = 0$  we get a jump to type  $A_5$  if  $B \neq 0$ .

5a) Next combine the two conditions from 4):  $K = P = Q = A = B = 0$ :

$$F = y^2 + Ny^3z + Ry^2z^2 - tz^4$$

If  $t \neq 0$  we have type  $A_3$ , but if  $t = 0$  we get a jump to type  $A_\infty$

5b) The case 4a) degenerates to:  $K = P = R = A = B = 0$ :

$$F = y^2 + Ny^3z + Qyz^2 - tz^4$$

If  $t \neq -Q^2/4$  we have type  $A_3$ , but if  $t = -Q^2/4$  we get a jump to type  $A_6$  if  $NQ \neq 0$ . (This is the first case where  $N \neq 0$  is really important).

6)  $K = P = Q = R = A = B = 0$ :

$$F = y^2 + Ny^3z - tz^4$$

If  $t \neq 0$  we have type  $A_3$ , but if  $t = 0$  we get a jump to type  $A_\infty$  (also if  $N = 0$ , but then the situation at the other point is changing drastically).

In the case  $N = 0$  we have that both points are singular. The computations follow the above scheme and

the answers are listed in the table below (second part):

type	$f_4 = x^2y^2$							
$\mathbf{A}^2\mathbf{B}^2$	$f = x^2y^2 + Kx^3 + Ny^3 + Px^2 + Qxy + Ry^2 + Ax + By$							
	'typical' form	$(1 : 0 : 0)$	$(0 : 1 : 0)$	$= 0$	$\neq 0$	$\mu$	$\lambda$	$\gamma$
$\mathbf{A}^2\mathbf{B}^2_7$	$x^2y^2 + x^3 + y^3$	$A_0$	$A_0$		$N, K$	7	0	7
$\mathbf{A}^2\mathbf{B}^2_6$	$x^2y^2 + x^2 + y^3$	$A_1$	$A_0$	$K$	$N, P$	6	0	6
$\mathbf{A}^2\mathbf{B}^2_5$	$x^2y^2 + x + y^3$	$A_2$	$A_0$	$K, P$	$N, A$	5	0	5
$\mathbf{A}^2\mathbf{B}^2_{4,1}$	$x^2y^2 + y^3 + Qxy + Ry^2 + By$	$A_3 \rightarrow A_4$	$A_0$	$K, P, A$	$N, Q, B, R$	3	1	4
$\mathbf{A}^2\mathbf{B}^2_{4,2}$	$x^2y^2 + y^3 + Qxy + Ry^2$	$A_3 \rightarrow A_5$	$A_0$	$K, P, A, B$	$N, Q, R$	2	2	4
$\mathbf{A}^2\mathbf{B}^2_{4,2}^*$	$x^2y^2 + y^3 + Ry^2 + By$	$A_3 \rightarrow A_5$	$A_0$	$K, P, A, Q$	$N, R, B$	2	2	4
$\mathbf{A}^2\mathbf{B}^2_{4,*}$	$x^2y^2 + y^3 + Ry^2$	$A_3 \rightarrow A_\infty$	$A_0$	$K, P, A, B, Q$	$N, R$	$\infty$	*	4
$\mathbf{A}^2\mathbf{B}^2_{4,3}$	$x^2y^2 + y^3 + Qxy$	$A_3 \rightarrow A_6$	$A_0$	$K, P, R, A, B$	$N, Q$	1	3	4
$\mathbf{A}^2\mathbf{B}^2_{4,**}$	$x^2y^2 + y^3$	$A_3 \rightarrow A_\infty$	$A_0$	$K, P, R, A, B, Q$	$N$	$\infty$	*	4
$\mathbf{A}^2\mathbf{B}^2_5^+$	$x^2y^2 + x^2 + y^2$	$A_1$	$A_1$	$K, N$	$P, R$	5	0	5
$\mathbf{A}^2\mathbf{B}^2_4^+$	$x^2y^2 + x + y^2$	$A_2$	$A_1$	$K, N, P$	$A, R$	4	0	4
$\mathbf{A}^2\mathbf{B}^2_{3,1}^+$	$x^2y^2 + Qxy + Ry^2 + By$	$A_3 \rightarrow A_4$	$A_1$	$K, N, P, A$	$B, Q, R$	2	1	3
$\mathbf{A}^2\mathbf{B}^2_{3,2}^+$	$x^2y^2 + Qxy + Ry^2$	$A_3 \rightarrow A_5$	$A_1$	$K, N, P, A, B$	$Q, R$	1	2	3
$\mathbf{A}^2\mathbf{B}^2_{3,2}^{+,*}$	$x^2y^2 + Ry^2 + By$	$A_3 \rightarrow A_5$	$A_1$	$K, N, P, A, Q$	$B, R$	1	2	3
$\mathbf{A}^2\mathbf{B}^2_{3,*}^+$	$x^2y^2 + Ry^2$	$A_3 \rightarrow A_\infty$	$A_1$	$K, N, P, A, B, Q$	$R$	$\infty$	*	3
$\mathbf{A}^2\mathbf{B}^2_3^+$	$x^2y^2 + x + y$	$A_2$	$A_2$	$K, N, P, R$	$A, B$	3	0	3
$\mathbf{A}^2\mathbf{B}^2_{2,1}^+$	$x^2y^2 + Qxy + By$	$A_3 \rightarrow A_4$	$A_2$	$K, N, P, R, A$	$B, Q$	1	1	2
$\mathbf{A}^2\mathbf{B}^2_{2,2}^+$	$x^2y^2 + By$	$A_3 \rightarrow A_5$	$A_2$	$K, N, P, R, A, Q$	$B$	0	2	2
$\mathbf{A}^2\mathbf{B}^2_{1,*}^+$	$x^2y^2 + Qxy$	$A_3 \rightarrow A_\infty$	$A_3 \rightarrow A_\infty$	$K, N, P, R, A, B$		$\infty$	*	1

All jumps occur if  $t = -Q^2/4$

**Case  $\mathbf{A}^3\mathbf{B}$ :**  $f_4 = x^3y$

First we consider

$$f_3 = Kx^3 + Lx^2y + Mxy^2 + Ny^3$$

By using translations  $\mathbb{T}_{\alpha,\beta}$  we could assume  $K = L = 0$ , but we can also arrange that  $K = B = 0$ . We make this last choice and get easier computations later.

The intersection points at infinity are  $(1 : 0 : 0)$  and  $(0 : 1 : 0)$ .

We find that  $(1 : 0 : 0)$  is always regular ( $A_0$ ) and  $(0 : 1 : 0)$  is singular iff  $N = 0$ .

Consider the general form

$$f = x^3y + Lx^2y + Mxy^2 + Px^2 + Qxy + Ry^2 + Ax$$

The corresponding hypersurface  $F(x, y, z, t) = 0$  is given by

$$F = x^3y + Lx^2yz + Mxy^2z + Px^2z^2 + Qxyz^2 + Ry^2z^2 + Axz^3 - tz^4$$

On chart  $y = 1$  we have

$$F = x^3 + Lx^2z + Mxz + Px^2z^2 + Qxz^2 + Rz^2 + Axz^3 - tz^4$$

If  $M \neq 0$  we have type  $A_1$ .

So consider the case  $N = M = 0$  and  $R \neq 0$ . We have

$$F = x^3 + Lx^2z + Px^2z^2 + Qxz^2 + Rz^2 + Axz^3 - tz^4$$

We find that  $F \simeq x^3 + z^2$ , we we have type  $A_2$ .

Consider  $N = M = R = 0$ . We have

$$f = x(x^2 + Lx + Q)y + Px^2 + Ax$$

$$F = x^3 + Lx^2z + Qxz^2 + Px^2z^2 + Axz^3 - tz^4$$

We have 3 cases; related to the behaviour of the three roots of  $x(x^2 + Lx + Q)$ :

- 1) three different roots:  $L^2 - 4Q \neq 0$  and  $Q \neq 0$
- 2) two different roots:  $L^2 - 4Q = 0$  or  $Q = 0$  but not both zero
- 3) one root:  $L = Q = 0$

In the first case, we always have type  $D_4$ .

In the second case, we can assume that the double root corresponds to  $x = 0$ , which implies  $Q = 0$ .

$$F = x^3 + Lx^2z + Px^2z^2 + Axz^3 - tz^4$$

If  $t \neq 0$  we have type  $D_5$ .

In case  $t = 0$  and  $A \neq 0$ , then we have type  $D_6$ .

But if  $A = 0$  then we have

$$F \simeq x^2(x + Lz + Pz^2)$$

So we have type  $D_\infty$  (since  $L \neq 0$ ).

In the third case, we have

$$F = x^3 + Px^2z^2 + Az^3x - tz^4$$

If  $t \neq 0$ , we have  $E_6$ .

If  $t = 0$  and  $A \neq 0$ , we have  $E_7$ .

If  $t = A = 0$  and  $P \neq 0$ , we have a non isolated point. Type  $J_{2,\infty}$ .

If  $t = A = P = 0$ , we have  $F = x^3$  that is type  $\mathbf{TA}_2$ .

type	$f_4 = x^3y$							
$\mathbf{A}^3\mathbf{B}$	$f = x^3y + Lx^2y + Ny^3 + Mxy^2 + Px^2 + Qxy + Ry^2 + Ax$							
	'typical' form	$(0 : 1 : 0)$	$(1 : 0 : 0)$	$\neq 0$	$= 0$	$\mu$	$\lambda$	$\gamma$
$\mathbf{A}^3\mathbf{B}_7$	$x^3y + y^3$	$A_0$	$A_0$	$N$		7	0	7
$\mathbf{A}^3\mathbf{B}_6$	$x^3y + xy^2$	$A_1$	$A_0$	$M$	$N$	6	0	6
$\mathbf{A}^3\mathbf{B}_5$	$x^3y + y^2$	$A_2$	$A_0$	$R$	$N, M$	5	0	5
$\mathbf{A}^3\mathbf{B}_3$	$x^3y + xy$	$D_4$	$A_0$	$L^2 - 4Q, Q$	$N, M, R$	3	0	3
$\mathbf{A}^3\mathbf{B}_{2,1}$	$x^3y + x^2y + y$	$D_5 \rightarrow D_6$	$A_0$	$L, A$	$N, M, R, Q$	1	1	2
$\mathbf{A}^3\mathbf{B}_{2,*}$	$x^3y + x^2y$	$D_5 \rightarrow D_\infty$	$A_0$	$L$	$N, M, R, Q, A$	$\infty$	*	2
$\mathbf{A}^3\mathbf{B}_{1,1}$	$x^3y + x$	$E_6 \rightarrow E_7$	$A_0$	$A$	$N, M, R, Q, L$	0	1	1
$\mathbf{A}^3\mathbf{B}_{1,*}$	$x^3y + x^2$	$E_6 \rightarrow J_{2,\infty}$	$A_0$	$P$	$N, M, R, Q, L, A$	$\infty$	*	1
$\mathbf{A}^3\mathbf{B}_{1,**}$	$x^3y$	$E_6 \rightarrow \mathbf{TA}_2$	$A_0$		$N, M, R, Q, L, A, P$	$\infty$	*	1

All jumps other occur at  $t = 0$

**Case A<sup>4</sup>:**  $f_4 = x^4$

In this case we will use a different approach. We will alter the normal form of  $f$  several times in order to get easier conditions on the constants.

Let's begin with the general case.

$$f = x^4 + Kx^3 + Lx^2y + Mxy^2 + Ny^3 + Px^2 + Qxy + Ry^2 + Ax + By$$

By using the translation  $\mathbb{T}_{\frac{K}{4}, 0}$  we can assume  $K = 0$ .

The intersection point at infinity is  $(0 : 1 : 0)$  and it is singular iff  $N = 0$ . If  $N = 0$  and  $M \neq 0$  we have type  $A_1$ .

So let's consider  $N = M = 0$  and  $R \neq 0$ .

By using the translation  $\mathbb{T}_{0, \frac{B}{2R}}$  we can put  $B = 0$ .

So the corresponding hypersurface  $F(x, y, z, t) = 0$  on the chart  $y = 1$  is given by

$$F = x^4 + L_1x^2z + R_1z^2 + P_1x^2z^2 + Q_1xz^2 + A_1z^3x - tz^4.$$

Remark that  $R_1 = R \neq 0$ ,  $L_1 = L$  and  $Q_1 = Q$ .

In the case  $L_1^2 \neq 4R_1$  we have type  $A_3$ .

So let's assume that  $L_1^2 = 4R_1$ ,

$$f = (x^2 - cy)^2 + P_1x^2 + Q_1xy + A_1x$$

where  $c = L_1/2$ . The non-linear substitution  $y := c^{-1}y - x^2$  changes the polynomial to one of degree less than 4, which is considered before. (See the information in the list)

We continue now the affine classification of this case in degree 4. Remember  $L_1^2 = 4R_1$ , We have type  $A_4$  iff  $Q_1 \neq 0$ .

In the case  $Q_1 = 0$  and  $P_1 \neq 0$ , we have type  $A_5$ .

If  $Q_1 = P_1 = 0$  and  $A_1 \neq 0$ , we have type  $A_6$ .

If  $A_1 = 0$  and  $t \neq 0$ , we have type  $A_7$  and if  $t = 0$  as well, then we have type  $A_\infty$ .

The other case is  $M = N = R = 0$ .

We have the following hypersurface:

$$\begin{aligned} F &= Lx^2z + Qxz^2 + Bz^3 + x^4 + Px^2z^2 + Axz^3 - tz^4 \\ &= z(Bz^2 + Lx^2 + Qxz) + x^4 + Px^2z^2 + Axz^3 - tz^4. \end{aligned}$$

We now split into two different cases, namely:

- 1)  $L \neq 0$
- 2)  $L = 0$

1) In the case  $Q^2 \neq 4BL$  we have type  $D_4$ .

We can from now on assume that  $Q^2 = 4BL$ .

So we have  $f = x^4 + Lx^2y + Px^2 + Qxy + Ax + By$ .

By using the transformation  $\mathbb{T}_{0, \frac{B}{L}}$  we find

$$f = x^4 + L_2x^2y + Q_2xy + A_2x + B_2y.$$

Remark that  $L_2 = L \neq 0$ ,  $Q_2 = Q$  and  $B_2 = B$ . So we have

$$F = z(L_2x^2 + B_2z^2 + Q_2xz) + x^4 + A_2xz^3 - tz^4.$$

This results into type  $D_5$  iff

$$t \neq \frac{B}{L^2} - \frac{A_2Q}{2L}.$$

If  $t = \frac{B}{L^2} - \frac{A_2Q}{2L}$ , and  $A_2L^2 - 2QB \neq 0$  we have  $D_6$  and in the other case we have  $D_\infty$ .

2) If  $L = 0$  and  $Q \neq 0$ , we can use the transformation

$$\begin{cases} x := x \\ y := \frac{-P}{Q}x + y + \frac{PB-AQ}{Q^2} \end{cases}$$

on  $f = x^4 + Px^2 + Qxy + Ax + By$  to get

$$F = z^2(Qx + Bz) + x^4 - tz^4.$$

This is type  $D_5$ , because of the influence of  $x^4$ .

The remaining case is  $L = Q = 0$ .

If  $B \neq 0$ , we can use the transformation

$$\begin{cases} x := x \\ y := y - \frac{A}{B}x \end{cases}$$

This results into the following form

$$F = Bz^3 + x^4 + Px^2z^2 - tz^4,$$

so we have type  $E_6$ .

If  $L = Q = B = 0$ , we find

$$F = x^4 + Px^2z^2 + Axz^3 - tz^4.$$

The polynomial is homogeneous of degree 4. In general there are four different solutions of the equation  $F = 0$ . In case we have a double root, the following system has to be solved.

$$\begin{cases} x^4 + Px^2 + Ax - t = 0 \\ 4x^3 + 2Px + A = 0 \end{cases}$$

So in case  $A \neq 0$  and  $27A^2 + 8P^3 \neq 0$ , we have  $\tilde{E}_7$  with jumps to  $T_{\infty,4,2}$ . We have three different jumps ( $t_1, t_2$  and  $t_3$ ). This is a result of the fact that  $t = \frac{2Px^2+3Ax}{4}$  and there are three different solutions to  $4x^3 + 2Px + A = 0$ .

When  $PA \neq 0$  and  $27A^2 + 8P^3 = 0$ , we get  $\tilde{E}_7$  with jumps to  $F_1A_2$  if  $t^3 = \frac{-27A^4}{8^4}$ .

If  $A = 0$  and  $P \neq 0$  then we have a very special situation. Here occurs two different jumps. We have  $\tilde{E}_7$  with a jump to  $T_{\infty,\infty,2}$  when  $t = \frac{P^2}{4}$ . And we have  $\tilde{E}_7$  with a jump to  $T_{\infty,4,2}$  when  $t = 0$ .

The remaining case is  $P = A = 0$ . In this case we have  $\tilde{E}_7$  with a jump to  $\mathbf{T}A_3$  if  $t = 0$ .

type	$f_4 = x^4$						
$\mathbf{A}^4$	$f = x^4 + Lx^2y + Mxy^2 + Ny^3 + Px^2 + Qxy + Ry^2 + Ax + By$						
	'typical' form	$(0 : 1 : 0)$	$\neq 0$	$= 0$	$\mu$	$\lambda$	$\gamma$
$\mathbf{A}^4_6$	$x^4 + y^3$	$A_0$	$N$		6	0	6
$\mathbf{A}^4_5$	$x^4 + xy^2$	$A_1$	$M$	$N$	5	0	5
$\mathbf{A}^4_3$	$x^4 + y^2$	$A_3$	$R, (L_1^2 - 4R_1)$	$M, N$	3	0	3
$\mathbf{A}^4_2 \simeq \mathbf{A}^3_2$	$(x^2 + y)^2 + xy$	$A_4$	$R, Q_1$	$M, N, (L_1^2 - 4R_1)$	2	0	2
$\mathbf{A}^4_1 \simeq \mathbf{A}^2\mathbf{B}_1$	$(x^2 + y)^2 + x^2$	$A_5$	$R, P_1$	$M, N, (L_1^2 - 4R_1), Q_1$	1	0	1
$\mathbf{A}^4_0 \simeq \mathbf{A}_0$	$(x^2 + y)^2 + x$	$A_6$	$R, A_1$	$M, N, (L_1^2 - 4R_1), Q_1, P_1$	0	0	0
$\mathbf{A}^4_{\star} \simeq \mathbf{A}^2_{\star}$	$(x^2 + y)^2$	$A_7 \rightarrow A_{\infty}$	$R$	$M, N, (L_1^2 - 4R_1), Q_1, P_1, A_1$	$\infty$		
$\mathbf{A}^4_2^+ \simeq \mathbf{A}^2\mathbf{B}_2$	$x^4 + x^2y + xy$	$D_4$	$(Q^2 - 4BL), Q, L$	$N, M, R$	2	0	2
$\mathbf{A}^4_{1,1}^+ \simeq \mathbf{A}^2\mathbf{B}_{1,1}$	$x^4 + x^2y + x$	$D_5 \xrightarrow{\star} D_6$	$L, (A_2L^2 - 2QB)$	$N, M, R, (Q^2 - 4BL)$	0	1	1
$\mathbf{A}^4_{1,\star}^+ \simeq \mathbf{A}^2\mathbf{B}_{1,\star}$	$x^4 + x^2y$	$D_5 \xrightarrow{\star} D_{\infty}$	$L$	$N, M, R, (Q^2 - 4BL), (A_2L^2 - 2QB)$	$\infty$	$\star$	1
$\mathbf{A}^4_1^+ \simeq \mathbf{A}^2\mathbf{B}_1$	$x^4 + xy$	$D_5$	$Q$	$N, M, R, L$	1	0	1
$\mathbf{A}^4_0^+ \simeq \mathbf{A}_0$	$x^4 + y$	$E_6$	$B$	$N, M, R, L, Q$	0	0	0
$\mathbf{A}^4_{\star 1}$	$x^4 + Px^2 + Ax$	$\tilde{E}_7 \xrightarrow{\star} T_{\infty,4,2}$	$A, (27A^2 + 8P^3)$	$N, M, R, L, Q, B$	$\infty$		
$\mathbf{A}^4_{\star 2}$	$x^4 + Px^2 + Ax$	$\tilde{E}_7 \xrightarrow{\Delta} F_{1,A_2}$	$A, P$	$N, M, R, L, Q, B, (27A^2 + 8P^3)$	$\infty$		
$\mathbf{A}^4_{\star 3}$	$x^4 + Px^2$	$\tilde{E}_7 \xrightarrow{\diamond} T_{\infty,\infty,2}$	$P$	$N, M, R, L, Q, B, A$	$\infty$		
$\mathbf{A}^4_{\star 4}$	$x^4 + Px^2$	$\tilde{E}_7 \rightarrow T_{\infty,4,2}$	$P$	$N, M, R, L, Q, B, A$	$\infty$		
$\mathbf{A}^4_{\star 5}$	$x^4$	$\tilde{E}_7 \rightarrow \mathbf{TA}_3$		$N, M, R, L, Q, B, A, P$	$\infty$		
<p>The jumps marked with a <math>\star</math> occur if <math>t = \frac{B}{L^2} - \frac{A_2Q}{2L}</math>. The jump marked with a <math>\diamond</math> occurs if <math>t = \frac{P^2}{4}</math>.</p> <p>The jump marked with a <math>\star</math> occurs at <math>t \in \{t_1, t_2, t_2\}</math>.</p> <p>The jump marked with a <math>\Delta</math> occurs when <math>t^3 = \frac{-27A^4}{8^4}</math>. All other jumps occur if <math>t = 0</math>.</p>							

### List of polynomials with $\mu(f) = 0$ .

Only the following polynomials occur in the list (up to diffeomorphism  $\cong$ ):

$\mathbf{A}_0$	$x$	
$\mathbf{A}^2\mathbf{B}_{1,1}$	$x^2y + Px$	$P \neq 0$
$\mathbf{A}^2\mathbf{B}_{2,2}^+$	$x^2y^2 + By$	$B \neq 0$
$\mathbf{A}^3\mathbf{B}_{1,1}$	$x^3y + Px^2 + Ax$	$A \neq 0$



## References

- [AVG] V. Arnol'd, A. Varchenko & S. Gussein-Zade, *Singularities of differentiable mappings, Volumes 1 and 2*, Birkhauser 1985, 1988.
- [Br] S.A. Broughton, *Milnor number and the topology of polynomial hypersurface*, *Inventiones Math.* **92** (1988), pp. 217–241.
- [dJ] Th. de Jong, *Some classes of line singularities*, *Math. Zeitschrift*, 198 (1988), 493-517.
- [HL] H.V. Hà & D.T. Lê, *Sur la topologie des polynômes complexes*, *Acta Mathematica Vietnamica*, **9** (1984), p. 21-32.
- [Sch] R. Schrauwen, *Series of singularities and their topology*, Academisch proefschrift, Utrecht, 1990.
- [Sie1] D. Siersma, *Classification and deformation of singularities*, Academisch proefschrift, Amsterdam, 1974.
- [Sie2] D. Siersma, *Isolated line singularities*, *Proceeding of Symposia in Pure Mathematics*, **40** (1983), p. 485-496.
- [SiTi] D. Siersma & M. Tibăr, *Singularities at infinity and their vanishing cycles*, To appear in *Duke Mathematical Journal*.
- [Wal1] C.T.C. Wall, *Affine cubic functions; the complex plane*, *Mathematical Proceedings of the Cambridge Philosophical Society*, **85** (1979), p. 387-401.
- [Wal2] C.T.C. Wall, *Affine cubic functions; the real plane*, *Mathematical Proceedings of the Cambridge Philosophical Society*, **87** (1980), p. 1-14.

Mathematisch Instituut  
University Utrecht  
P. O. Box 80.010  
3508 TA Utrecht, The Netherlands  
email: siersma@math.ruu.nl, job@win.tue.nl