

## Three-body recombination for vanishing scattering lengths

**Citation for published version (APA):**

Kokkelmans, S. J. J. M. F., Shotan, Z., Machtey, O., & Khaykovich, L. (2014). Three-body recombination for vanishing scattering lengths. In *Proceedings of the 46th Conference of the European Group on Atomic Systems (EGAS), 1-4 July 2014, Lille, France* (Europhysics conference abstracts; Vol. 38D). European Physical Society (EPS).

**Document status and date:**

Published: 01/01/2014

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

## Three-body recombination for vanishing scattering lengths

S. Kokkelmans<sup>1</sup>, Z. Shotan<sup>2</sup>, O. Machtey<sup>2</sup>, and L. Khaykovich<sup>2</sup>

<sup>1</sup>*Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands*

<sup>2</sup>*Department of Physics, Bar-Ilan University, Ramat-Gan, 52900 Israel*

Presenting Author: s.kokkelmans@tue.nl

Efimov physics in ultracold gases is described very well by universal scaling laws, based on the scattering length. Another important parameter in this respect is the van der Waals length, which was only recently found to define the binding energies of the Efimov trimers. While the scattering length can be tuned magnetically via a Feshbach resonance, the van der Waals length is constant and connected to the radial range of the potential. However, experimental hints at non-universal behavior, when going away from resonance, are quite badly understood. The next leading coefficient in the scattering phase shift, the effective range parameter, gives an indication of this non-universality, but at the same time it can also be strongly dependent on the magnetic field. An extreme case in the non-universal regime is the zero-crossing of a Feshbach resonance. Here the scattering length vanishes, while the effective range goes to infinity. Therefore both parameters cannot directly describe three-body recombination at this point.

We investigate both experimentally and theoretically three-body recombination, for two different zero crossings in an ultracold gas of  ${}^7\text{Li}$ . One zero crossing is associated to a broad resonance, while the other one is connected to a narrow resonance. We see a clear experimental difference for the two resonances, and we give an explanation for the behavior in terms of a new theoretical effective length parameter. This parameter is described in terms of the finite range of the two-body potential, given by the van der Waals length, and by the width of the resonance.