

Lace expansion for dummies

Citation for published version (APA):

Bolthausen, E., Van Der Hofstad, R., & Kozma, G. (2018). Lace expansion for dummies. *Annales de l'institut Henri Poincaré (B): Probability and Statistics*, 54(1), 141-153. <https://doi.org/10.1214/16-AIHP797>

DOI:

[10.1214/16-AIHP797](https://doi.org/10.1214/16-AIHP797)

Document status and date:

Published: 01/02/2018

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Lace expansion for dummies

Erwin Bolthausen^{a,1}, Remco van der Hofstad^{b,2} and Gady Kozma^{c,3}

^a*Institut für Mathematik, Universität Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland and Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan. E-mail: eb@math.uzh.ch*

^b*Department of Mathematics and Computer Science, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands. E-mail: r.w.v.d.hofstad@tue.nl*

^c*Weizmann Institute, Rehovot, 76100, Israel. E-mail: gady.kozma@weizmann.ac.il*

Received 26 April 2016; revised 27 August 2016; accepted 22 September 2016

Abstract. We show Green's function asymptotic upper bound for the two-point function of weakly self-avoiding walk in $d > 4$, revisiting a classic problem. Our proof relies on Banach algebras to analyse the lace-expansion fixed point equation and is simpler than previous approaches in that it avoids Fourier transforms.

Résumé. Nous montrons une domination asymptotique de la fonction à deux points de la marche faiblement auto-évitante en dimension $d > 4$ par la fonction de Green, revisitant ainsi un problème classique. Notre preuve s'appuie sur des techniques d'algèbres de Banach pour analyser le point fixe de l'équation de développement en lacets. Elle est plus simple que les approches précédentes car elle ne passe pas par la transformée de Fourier.

MSC: 84B41; 60K35

Keywords: Self-avoiding walk; Lace expansion; Banach algebra; Deconvolution; Edgeworth expansion

1. Introduction

The lace expansion made its debut in 1985 with a proof by Brydges and Spencer that weakly self-avoiding walk (WSAW) has “Gaussian behaviour” in dimensions 5 and above [4]. It proved to be useful way beyond its initial application, primarily in work by Hara and Slade. The technique was applied to percolation [7], lattice trees and animals [8], the contact process [12], the Ising model [13] and φ^4 [14]. Further, it was extended to finite graphs [3] and to long-range models [5]. Despite all this progress, weakly self-avoiding walk remains the simplest example to which the technique applies: lace expansion is a “perturbative” technique and it requires a small parameter. Weakly self-avoiding walk has such a small parameter naturally built-in, while for most models, the small parameter is more hidden. Consequently, it was used as a test bed for several new techniques, for example in [10], where the lace expansion was analysed using induction in time, and [2] where a Banach fixed point theorem was used. Interestingly, neither of these papers uses the so-called bootstrap analysis introduced in [15]. In our opinion, the bootstrap analysis is the most important simplification to lace expansion, replacing the difficult “moving single pole” analysis of [4]. The bootstrap analysis applies to generating functions such as the WSAW Green's function, while [2,10] prove results for WSAW with a fixed number of steps instead. Green's function asymptotics in x -space as derived here were proved

¹Supported by SNF Grant 200020_138141.

²Supported by the Netherlands Organisation for Scientific Research (NWO) through VICI Grant 639.033.806 and the Gravitation NETWORKS Grant 024.002.003.

³Supported by the Israel Science Foundation and the Jesselson Foundation, and by the CNRS during his visit to the Institut Henri Poincaré.

previously in technically more challenging settings in [9] for spread-out models and in [6] for nearest-neighbour settings. Brydges and Spencer [4] prove Gaussian limit laws for the end-to-end displacement for WSAW after n steps.

Our starting point was also an attempt to generalise lace expansion, rather than to simplify it. We wished to apply it to problems on Cayley graphs of non-commutative groups. Most of the existing approaches rely heavily on the Fourier transform, which is of course no longer available in this new setting. The approach of [1,2], though, turned out to be applicable. We realised that it can be simplified and generalised by working in an appropriate Banach algebra.

In this paper, we expose our Banach-algebra approach in the simplest possible setting: weakly self-avoiding walk on \mathbb{Z}^d , with the result being an upper bound on the critical Green's function. We repeat that related results have been proved previously, only our proof is novel.

2. Precise definitions and statement of the theorem

For a nearest-neighbour path $\gamma: \{0, \dots, n\} \rightarrow \mathbb{Z}^d$ and a $\beta \in [0, 1]$, we define its weight by

$$W(\gamma) = W^\beta(\gamma) = (1 - \beta)^{|\{0 \leq s < t \leq n: \gamma(s) = \gamma(t)\}|}, \quad (1)$$

i.e., the path is ‘‘penalised’’ by $1 - \beta$ for every self-intersection of γ . We define the weakly self-avoiding walk Green's function to be

$$G_\lambda^{\text{saw}}(x) = G_\lambda^{\beta, \text{saw}}(x) = \sum_{\gamma: 0 \rightarrow x} \lambda^{\text{len}(\gamma)} W^\beta(\gamma),$$

where the notation $\gamma: 0 \rightarrow x$ means that $\gamma: \{0, 1, \dots, n\} \rightarrow \mathbb{Z}^d$ is a nearest-neighbour path with $\gamma(0) = 0, \gamma(n) = x$, while $\text{len}(\gamma) = n$. We define λ_c to be the critical value for the finiteness of the spatial sum of G_λ^{saw} , i.e.,

$$\lambda_c = \sup \left\{ \lambda: \sum_{x \in \mathbb{Z}^d} G_\lambda^{\text{saw}}(x) < \infty \right\}.$$

Finally denote by $G^{\text{rw}}(x)$ the (critical) Green's function of simple random walk (SRW) on \mathbb{Z}^d , i.e.,

$$G^{\text{rw}}(x) = \sum_{n=0}^{\infty} p_n(x),$$

where $p_n(x)$ is the probability that simple random walk on \mathbb{Z}^d starting from 0 is at x at time n . When $d > 2$ the sum converges and

$$G^{\text{rw}}(x) = (a + o(1))|x|^{2-d} \quad \text{as } |x| \rightarrow \infty \quad (2)$$

with $a > 0$. See e.g. [17]. The result is that the WSAW Green's function is bounded by the SRW Green's function for $d > 4$:

Theorem (Green's function upper bound). *Let $d > 4$. Then there exists a β_0 such that for all $\beta < \beta_0$, β -weakly self avoiding walk satisfies*

$$G_{\lambda_c}^{\text{saw}}(x) \leq 2G^{\text{rw}}(x) \quad \forall x \in \mathbb{Z}^d.$$

Remark. We will also show a lower bound, $G_{\lambda_c}^{\text{saw}} \geq \frac{1}{2}G^{\text{rw}}$, and further that

$$G_{\lambda_c}^{\text{saw}}(x) = (1 + O(\beta))G^{\text{rw}}(x) \quad \text{as } \beta \rightarrow 0, \text{ uniformly in } x \in \mathbb{Z}^d. \quad (3)$$

See the remarks on page 152. We find the upper bound to be the more interesting and we prefer to focus on it.

The remainder of this paper is devoted to the proof of this theorem.

3. Proof

For $\mu \in \mathbb{R}$ we denote by $\Delta_\mu^{\text{rw}} : \mathbb{Z}^d \rightarrow \mathbb{R}$ the following function

$$\Delta_\mu^{\text{rw}}(x) = \begin{cases} 1, & x = 0, \\ -\mu, & x \text{ is a neighbour of } 0, \\ 0, & \text{otherwise.} \end{cases} \quad (4)$$

We say that a function $f : \mathbb{Z}^d \rightarrow \mathbb{R}$ is ‘‘symmetric to coordinate permutations and flipping’’ if for any $\sigma \in S_d$ (the group of permutations on d elements) and for any $\varepsilon_1, \dots, \varepsilon_d \in \{\pm 1\}$,

$$f(x_1, \dots, x_d) = f(\varepsilon_1 x_{\sigma(1)}, \dots, \varepsilon_d x_{\sigma(d)}).$$

We further write $f * g$ for the convolution of two functions $f, g : \mathbb{Z}^d \rightarrow \mathbb{R}$ and $\delta_0(x) = 1_{\{x=0\}}$, $x \in \mathbb{Z}^d$.

Lemma 1 (Lace expansion analysis). *Let $d > 4$. Then there exists a β_0 such that for all $\beta < \beta_0$ and for all $\lambda < \lambda_c$ the following holds. If $G_\lambda^{\text{saw}}(x) \leq 3G^{\text{rw}}(x)$ for all $x \in \mathbb{Z}^d$, then there exists a function $\Delta_\lambda^{\text{saw}} : \mathbb{Z}^d \rightarrow \mathbb{R}$ such that $G_\lambda^{\text{saw}} * \Delta_\lambda^{\text{saw}} = \delta_0$ and such that*

- (i) $\Delta_\lambda^{\text{saw}}$ is symmetric to coordinate permutations and flipping;
- (ii) $\sum_x \Delta_\lambda^{\text{saw}}(x) \geq 0$;
- (iii) There exists $\lambda' \in [0, \frac{1}{2d}]$ such that

$$|\Delta_\lambda^{\text{saw}}(x) - \Delta_{\lambda'}^{\text{rw}}(x)| \leq C\beta|x|^{-d-4}.$$

A somewhat abusive convention we adopt here and below is that $|x|^{-\alpha} = 1$ when $x = 0$, so condition (iii) in fact implies that $|\Delta_\lambda^{\text{saw}}(0) - \Delta_{\lambda'}^{\text{rw}}(0)| \leq C\beta$. C and c are used for constants that depend only on the dimension. Let us remark that in fact we simply take $\lambda' = \min(\lambda, \frac{1}{2d})$, though we will not use this fact. Another remark worth making is that in (iii) we will in fact prove, $|\Delta_\lambda^{\text{saw}}(x) - \Delta_{\lambda'}^{\text{rw}}(x)| \leq C\beta|x|^{6-3d}$ which is of course stronger than the stated estimate when $d \geq 5$. However, it will be convenient to formulate the lemma as above.

We remark that it is tempting to think about $\Delta_\lambda^{\text{saw}}$ as a generator of some random walk (with killing), but it is missing one important property of a generator: it is not true that $\Delta_\lambda^{\text{saw}}(x) < 0$ for all $x \neq 0$. This means that a lot of deconvolution techniques for random walk generators are inapplicable. The next lemma is the required deconvolution:

Lemma 2 (Deconvolution). *Let $d > 2$. Then there exists β_0 such that for all $\beta < \beta_0$ and for any $\Delta : \mathbb{Z}^d \rightarrow \mathbb{R}$ satisfying conditions (i)–(iii) of Lemma 1, there exists a function G such that $G * \Delta = \delta_0$ and $|G(x)| \leq 2G^{\text{rw}}(x)$.*

We postpone the proof of both lemmas and first show how they imply the theorem:

Proof of the theorem given Lemmas 1 and 2. Fix β to be some value sufficiently small so that both Lemmas 1 and 2 hold with this value of β . The following argument, known as a bootstrap argument, goes back to Slade [15]. Define

$$f(\lambda) = \sup_{x \in \mathbb{Z}^d} \frac{G_\lambda^{\text{saw}}(x)}{G^{\text{rw}}(x)}.$$

We first examine $f(0)$. $G_0^{\text{saw}} = \delta_0$ and of course $G^{\text{rw}} \geq \delta_0$ so $f(0) \leq 1$. Next we note that f is continuous in the interval $[0, \lambda_c)$. Indeed, λ_c is the radius of convergence of $\sum_x G_\lambda^{\text{saw}}(x)$ and hence is a lower bound for the radius of convergence of $G_\lambda^{\text{saw}}(x)$ for any x . Hence each term $G_\lambda^{\text{saw}}(x)/G^{\text{rw}}(x)$ is continuous on our interval. On the other hand, because the sum defining $G_\lambda^{\text{saw}}(x)$ contains only paths of length at least $|x|$, it also decays exponentially in x , uniformly on $[0, \lambda]$, for all $\lambda < \lambda_c$. Using (2), the same is true for the quotient $G_\lambda^{\text{saw}}(x)/G^{\text{rw}}(x)$. This means that on any $[0, \lambda]$ with $\lambda < \lambda_c$, f can be written as the supremum of a finite collection of continuous functions, and hence is continuous. Since λ can be taken arbitrarily close to λ_c , f is continuous on $[0, \lambda_c)$.

We now claim that it is not possible that $f(\lambda) \in (2, 3]$ for any $\lambda < \lambda_c$. Indeed, if $f(\lambda) \leq 3$ then $G_\lambda^{\text{saw}}(x) \leq 3G^{\text{rw}}(x)$ for all x and the condition of Lemma 1 is satisfied. We use Lemma 1 to find some $\Delta_\lambda^{\text{saw}}$ with $G_\lambda^{\text{saw}} * \Delta_\lambda^{\text{saw}} = \delta_0$ satisfying conditions (i)–(iii), and then Lemma 2 to find some G such that $G * \Delta_\lambda^{\text{saw}} = \delta_0$ and $G(x) \leq 2G^{\text{rw}}(x)$. We now claim that $G_\lambda^{\text{saw}} = G$. Indeed, both functions are in $\ell^2(\mathbb{Z}^d)$, G_λ^{saw} by the assumption $G_\lambda^{\text{saw}}(x) \leq 3G^{\text{rw}}(x)$ and G by the conclusion of Lemma 2 – and so is $\Delta_\lambda^{\text{saw}}$ by condition (iii). In ℓ^2 , deconvolution can be performed by Fourier transform and hence is unique. We get that $G_\lambda^{\text{saw}}(x) = G(x) \leq 2G^{\text{rw}}(x)$ so $f(\lambda) \leq 2$. We conclude that $f(\lambda) \notin (2, 3]$ for any $\lambda < \lambda_c$.

Now, if f is continuous, starts below 1 and cannot traverse the interval $(2, 3]$, then it must be that $f(\lambda) \leq 2$ for all $\lambda < \lambda_c$, i.e., $G_\lambda^{\text{saw}}(x) \leq 2G^{\text{rw}}(x)$ for all x and all $\lambda < \lambda_c$. Finally, by monotone convergence, $G_{\lambda_c}^{\text{saw}}(x) = \lim_{\lambda \nearrow \lambda_c} G_\lambda^{\text{saw}}(x)$, so that also $G_{\lambda_c}^{\text{saw}}(x) \leq 2G^{\text{rw}}(x)$ for all $x \in \mathbb{Z}^d$. \square

We move to the proofs of Lemmas 1 and 2. Lemma 1 essentially relies on the same lace-expansion argument as performed by Brydges and Spencer [4] – we include the proof for completeness, but we will be a little brief. Lemma 2 is the new ingredient of our paper.

Proof of Lemma 1. We follow [10, Appendix A] closely for the derivation of the lace expansion, and [9] for the analysis of the coefficients arising in it. As β and λ are fixed, let us remove them from the notation and denote our functions by G^{saw} and Δ^{saw} . We start by finding a formula for Δ^{saw} (or rather, a representation as an infinite sum). Recall the weight $W(\gamma)$ defined in (1). We define

$$U_{st}(\gamma) = \begin{cases} 0, & \text{when } \gamma(s) \neq \gamma(t), \\ -\beta, & \text{when } \gamma(s) = \gamma(t), \end{cases} \quad \forall 0 \leq s < t \leq \text{len}(\gamma),$$

so that

$$W(\gamma) = \prod_{s < t} (1 + U_{st}(\gamma)).$$

Given an interval $I = [a, b]$ of integers with $0 \leq a \leq b$, we refer to a pair $\{s, t\}$ ($s < t$) of elements of I as an *edge*. To abbreviate the notation, we write st for $\{s, t\}$. A set of edges is called a *graph*. A graph Γ on $[a, b]$ is said to be *connected* if both a and b are endpoints of edges in Γ and if, in addition, for any $c \in (a, b)$ there is an edge $st \in \Gamma$ such that $s < c < t$ (note that this is unrelated to the usual definition of graph connectivity). The set of all graphs on $[a, b]$ is denoted $\mathcal{B}[a, b]$, and the subset consisting of all connected graphs is denoted $\mathcal{C}[a, b]$.

For integers $0 \leq a < b$, define

$$K[a, b](\gamma) = \prod_{a \leq s < t \leq b} (1 + U_{st}(\gamma)) \quad \text{so that } G^{\text{saw}}(x) = \sum_{\gamma: 0 \rightarrow x} \lambda^{\text{len}(\gamma)} K[0, \text{len}(\gamma)](\gamma), \quad (5)$$

where the sum is over all simple random walk paths from 0 to x . Expanding the product in the definition of $K[a, b](\gamma)$, we get

$$K[a, b](\gamma) = \sum_{\Gamma \in \mathcal{B}[a, b]} \prod_{st \in \Gamma} U_{st}(\gamma). \quad (6)$$

For $0 \leq a < b$ we define an analogous quantity, in which the sum over graphs is restricted to connected graphs, namely,

$$J[a, b](\gamma) = \sum_{\Gamma \in \mathcal{C}[a, b]} \prod_{st \in \Gamma} U_{st}(\gamma). \quad (7)$$

We claim that, for $n \geq 1$,

$$K[0, n] = K[1, n] + \sum_{m=2}^n J[0, m]K[m, n]. \quad (8)$$

To see this, we note from (6) that the contribution to $K[0, n]$ from all graphs Γ for which 0 is not in an edge is exactly $K[1, n]$. To resum the contribution from the remaining graphs, we proceed as follows. When Γ does contain an edge ending at 0, we let $m(\Gamma) > 0$ denote the smallest number that is not crossed by an edge, i.e., there is no $st \in \Gamma$ such that $s < m(\Gamma) < t$. We lose nothing by taking $m \geq 2$, since $U_{a, a+1} = 0$ for all a . Resummation over graphs on $[m, n]$ and (7) proves (8).

Let us now define the key quantities in the lace expansion, which are

$$\Pi_\lambda(x) = \sum_{\gamma: 0 \rightarrow x} \lambda^{\text{len}(\gamma)} J[0, \text{len}(\gamma)](\gamma) \quad (9)$$

and

$$\Delta_\lambda^{\text{saw}}(x) := \Delta_\lambda^{\text{rw}}(x) - \Pi_\lambda(x). \quad (10)$$

λ will be fixed through the rest of the proof of Lemma 1 and we usually drop it from the notation. The key to the proof of the lemma is the estimate

$$\sum_{n=0}^{\infty} \sum_{\substack{\gamma: 0 \rightarrow x \\ \text{len } \gamma = n}} |\lambda^n J[0, n](\gamma)| < C\beta|x|^{6-3d}, \quad (11)$$

which we will prove below. We first prove that it implies the claims in the lemma. Of course it implies

$$|\Pi(x)| \leq C\beta|x|^{6-3d}. \quad (12)$$

To conclude from the definitions and estimate above that $G^{\text{saw}} * \Delta^{\text{saw}} = \delta_0$ note that, by (8),

$$\begin{aligned} G^{\text{saw}}(x) &\stackrel{(5)}{=} \sum_{\gamma: 0 \rightarrow x} \lambda^{\text{len}(\gamma)} K[0, \text{len}(\gamma)](\gamma) \\ &\stackrel{(8)}{=} \delta_0 + \sum_{\gamma: 0 \rightarrow x} \lambda^{\text{len}(\gamma)} \left(K[1, \text{len}(\gamma)](\gamma) + \sum_{m=2}^{\text{len}(\gamma)} J[0, m](\gamma) K[m, \text{len}(\gamma)](\gamma) \right) \\ &= \delta_0 + \lambda \sum_{y: \|y\|=1} G^{\text{saw}}(x-y) + \sum_y \Pi(y) G^{\text{saw}}(x-y), \end{aligned} \quad (13)$$

where the last equality is derived as follows: the $K[1, \text{len}(\gamma)](\gamma)$ terms we separate according to $\gamma(1)$, which we denote by y . Translation invariance gives that each term is exactly $G^{\text{saw}}(x-y)$. The terms containing J are split according to $\gamma(m)$, which we denote by y , and again by translation invariance the sum over K gives $G^{\text{saw}}(x-y)$. Finally, the change of order of summation is justified by (11) and $G^{\text{saw}}(x) \leq C|x|^{2-d}$. This proves (13). Rearranging (13) gives $G^{\text{saw}} * \Delta_\lambda^{\text{rw}} = \delta_0 + G^{\text{saw}} * \Pi$, i.e. $G^{\text{saw}} * \Delta_\lambda^{\text{saw}} = \delta_0$, as required.

We move to prove properties (i)–(iii) of Δ^{saw} . The symmetry of Δ^{saw} is immediate from the construction, and the property that $\sum \Delta^{\text{saw}}(x) \geq 0$ comes from summing the relation $(G^{\text{saw}} * \Delta^{\text{saw}})(x) = \delta_0(x)$ over $x \in \mathbb{Z}^d$, which gives that

$$\sum_{x \in \mathbb{Z}^d} \Delta^{\text{saw}}(x) = \frac{1}{\sum_{x \in \mathbb{Z}^d} G^{\text{saw}}(x)}, \quad (14)$$

and the last term is clearly non-negative as well as finite since $\lambda < \lambda_c$ which means that $G^{\text{saw}}(x)$ decays exponentially as $x \rightarrow \infty$.

For (iii), we need to choose $\lambda' \in (0, \frac{1}{2d}]$ and estimate $\Delta_\lambda^{\text{saw}} - \Delta_{\lambda'}^{\text{rw}} \stackrel{(10)}{=} \Delta_\lambda^{\text{saw}} - \Delta_{\lambda'}^{\text{rw}} - \Pi_\lambda$. Observe first that by (12), we can take $\lambda' = \lambda$ if $\lambda \leq 1/2d$. If $\lambda > 1/2d$, we obtain from (12) and the already proved (ii)

$$1 - 2d\lambda = \sum_x \Delta_\lambda^{\text{rw}}(x) \stackrel{(10)}{=} \sum_x (\Pi_\lambda(x) + \Delta_\lambda^{\text{saw}}(x)) \stackrel{(ii)}{\geq} \sum_x \Pi_\lambda(x) \stackrel{(12)}{\geq} - \sum_x C\beta|x|^{6-3d} = -C\beta,$$



Fig. 1. Laces.

and therefore $\lambda - 1/2d \leq C\beta/2d$, and so we can take $\lambda' = 1/2d$ in this case. Therefore, $\lambda' = \min(\lambda, \frac{1}{2d})$ is doing the job.

It remains to prove (11).

We next rewrite (9) in a form that can be used to obtain good bounds on $\Pi(x)$. For this, we start by introducing the laces that give the lace expansion its name. A *lace* is a minimally connected graph, i.e., a connected graph for which the removal of any edge would result in a disconnected graph. The set of laces on $[a, b]$ is denoted $\mathcal{L}[a, b]$, and the set of laces on $[a, b]$ consisting of exactly N edges is denoted $\mathcal{L}^{(N)}[a, b]$. Given a connected graph Γ , the following prescription associates to Γ a unique lace L_Γ : The lace L_Γ consists of edges a_1b_1, a_2b_2, \dots , with $a_1, b_1, b_2, a_2, \dots$ determined, in that order, by

$$b_1 = \max\{t : at \in \Gamma\}, \quad a_1 = a,$$

$$b_i = \max\{t : \exists z < b_{i-1} \text{ such that } zt \in \Gamma\}, \quad a_i = \min\{z : zb_i \in \Gamma\}.$$

See Figure 1. Given a lace L , the set of all edges $st \notin L$ such that $L_{L \cup \{st\}} = L$ is denoted $\mathcal{C}(L)$. Edges in $\mathcal{C}(L)$ are said to be *compatible* with L . Now, $L_\Gamma = L$ if and only if $L \subset \Gamma$ and all edges in $\Gamma \setminus L$ are compatible with L . This allows to write

$$\sum_{\Gamma: L_\Gamma=L} \prod_{st \in \Gamma \setminus L} U_{st} = \prod_{st \in \mathcal{C}(L)} (1 + U_{st})$$

and then partially resum the right-hand side of (7), to obtain

$$J[a, b] = \sum_{L \in \mathcal{L}[a, b]} \sum_{\Gamma: L_\Gamma=L} \prod_{st \in L} U_{st} \prod_{s't' \in \Gamma \setminus L} U_{s't'} = \sum_{L \in \mathcal{L}[a, b]} \prod_{st \in L} U_{st} \prod_{s't' \in \mathcal{C}(L)} (1 + U_{s't'}). \quad (15)$$

For $0 \leq a < b$, we define $J^{(N)}[a, b]$ to be the contribution to (15) coming from laces consisting of exactly N edges:

$$J^{(N)}[a, b] = \sum_{L \in \mathcal{L}^{(N)}[a, b]} \prod_{st \in L} (-U_{st}) \prod_{s't' \in \mathcal{C}(L)} (1 + U_{s't'}), \quad N \geq 1. \quad (16)$$

Then, by (9),

$$J[a, b] = \sum_{N=1}^{\infty} (-1)^N J^{(N)}[a, b] \quad \text{and} \quad \Pi(x) = \sum_{N=1}^{\infty} (-1)^N \Pi^{(N)}(x), \quad (17)$$

where we define

$$\begin{aligned} \Pi^{(N)}(x) &= \sum_{\gamma: 0 \rightarrow x} \lambda^{\text{len}(\gamma)} J^{(N)}[0, \text{len}(\gamma)](\gamma) \\ &= \sum_{\gamma: 0 \rightarrow x} \lambda^{\text{len}(\gamma)} \sum_{L \in \mathcal{L}^{(N)}[0, \text{len}(\gamma)]} \prod_{st \in L} (-U_{st}(\gamma)) \prod_{s't' \in \mathcal{C}(L)} (1 + U_{s't'}(\gamma)). \end{aligned} \quad (18)$$

We will now show that the sum over N converges *absolutely*, which will show (11) and finish the lemma. The product over $st \in \mathcal{C}(L)$ will be easier to handle when we restrict it. Let therefore $\mathcal{D}(L)$ be the set of edges st such that the open interval (s, t) does not contain an a_i or b_i for any $(a_i, b_i) \in L$. Clearly $\mathcal{D}(L) \subseteq \mathcal{C}(L)$ and therefore

$\prod_{\mathcal{L}(L)}(1 + U_{st}) \leq \prod_{\mathcal{D}(L)}(1 + U_{st})$. Once we restrict, the sum over γ becomes independent between any two consecutive elements of L . For a lace $L = \{a_1b_1, a_2b_2, \dots, a_Nb_N\}$ we write c_1, c_2, \dots, c_{2N} for the ordered sequence of vertices of the bonds of L , i.e. $c_1 = a_1, c_2 = a_2, c_3 = b_1, \dots$. There can be matches among the c 's, e.g. for $N = 3$, b_1 can be equal to a_3 . We denote by $\gamma_i, 1 \leq i \leq 2N - 1$, the γ -path restricted to the interval $\{c_i, c_i + 1, \dots, c_{i+1}\}$, and we get, for $N \geq 2$,

$$\prod_{st \in \mathcal{D}(L)} (1 + U_{st}(\gamma)) = \prod_{i=1}^{2N-1} W(\gamma_i).$$

For $N = 1$, the left-hand side misses the bond from 0 to $\text{len}(\gamma)$, so that in this case, the right-hand side has to be divided by $(1 - \beta)$. Inserting this into the definition of $\Pi^{(N)}$ gives

$$\begin{aligned} \Pi^{(1)}(x) &\begin{cases} \leq \frac{\beta}{1-\beta} G^{\text{saw}}(0) & \text{for } x = 0, \\ = 0 & \text{for } x \neq 0, \end{cases} \\ |\Pi^{(2)}(x)| &\leq \beta^2 G^{\text{saw}}(x)^3, \end{aligned} \quad (19)$$

and, for $N \geq 2$,

$$|\Pi^{(N)}(x)| \leq \beta^N \sum_{x_2, \dots, x_{N-1}} G^{\text{saw}}(x_1 - x_2) \prod_{j=2}^N [G^{\text{saw}}(x_j - x_{j-1}) G^{\text{saw}}(x_{j+1} - x_{j-1})], \quad (20)$$

with $x_1 = 0, x_N = x_{N+1} = x$ (see again Figure 1).

To derive (20), observe that the terms U_{st} in (18) give the factor β^N as well as restrictions $\gamma(a_i) = \gamma(b_i)$ for all i . Under this restrictions γ breaks into paths γ_i which are independent given their endpoints, so their sum gives G^{saw} . This implies (20).

With these estimates in hand, we can bound Π .

$\Pi^{(1)}$ clearly poses no problems. For $\Pi^{(2)}$ we have $|\Pi^{(2)}(x)| \leq \beta^2 G^{\text{saw}}(x)^3$. By our assumptions $G^{\text{saw}}(x)^3 \leq 27G^{\text{rw}}(x)^3 \leq C|x|^{6-3d}$, as required. For the terms with $N \geq 3$, we need the following lemma:

Lemma 3. *Let $d > 4$. For any $u, v \in \mathbb{Z}^d$,*

$$\sum_{w \in \mathbb{Z}^d} |w|^{4-2d} |w - u|^{2-d} |w - v|^{2-d} \leq C|u|^{2-d} |v|^{2-d}.$$

Proof. By Cauchy–Schwarz

$$\left(\sum_{w \in \mathbb{Z}^d} |w|^{4-2d} |w - u|^{2-d} |w - v|^{2-d} \right)^2 \leq \left(\sum_{w \in \mathbb{Z}^d} |w|^{4-2d} |w - u|^{4-2d} \right) \left(\sum_{w \in \mathbb{Z}^d} |w|^{4-2d} |w - v|^{4-2d} \right). \quad (21)$$

For $d > 4$, each term can be estimated simply by splitting the sum to $|w| > |u|/2$ and $|w| \leq |u|/2$, see a detailed calculation in [9, Proposition 1.7(i)]. We get

$$\sum_{w \in \mathbb{Z}^d} |w|^{4-2d} |w - u|^{4-2d} \leq C|u|^{4-2d}. \quad \square$$

To use this lemma, define $A^{(N)}(x)$ by the right-hand side of (20) without the factor β^N , and $G(z)$ replaced by $|z|^{2-d}$. We prove by induction on $N \geq 2$ that

$$A^{(N)}(x) \leq C^N |x|^{6-3d}. \quad (22)$$

As $A^{(2)}(x) = (|x|^{2-d})^3$, the statement is true for $N = 2$. Let $N > 3$. In the expressions which are summed in $A^{(N)}$, we take out everything which depends on x_2 which is the factor

$$|x_2|^{4-2d} |x_3 - x_2|^{2-d} |x_4 - x_2|^{2-d}. \quad (23)$$

The rest of the product is

$$|x_3|^{2-d} \prod_{j=4}^N [|x_j - x_{j-1}|^{2-d} |x_{j+1} - x_{j-1}|^{2-d}]. \quad (24)$$

For $N = 3$, the product $\prod_{j=4}^N$ is absent and $x_3 = x$. Summing over x_2 in (23) we can estimate this part with Lemma 3 by $C|x_3|^{2-d}|x_4|^{2-d}$ which together with (24) and summed over the other variables is $CA_{N-1}(x)$. (Remember that we had $x_{N+1} = x_N = x$.) This proves (22).

Using (22) together with $G^{\text{saw}}(x) \leq 3G^{\text{rw}}(x) \leq C|x|^{2-d}$ proves $|\Pi^{(N)}(x)| \leq (C\beta)^N |x|^{6-3d}$ for $N \geq 2$. Together with (19), and summing over N , for sufficiently small β , this proves Lemma 1. \square

4. Proof of Lemma 2, with Banach algebras

We start by defining a norm on $f: \mathbb{Z}^d \rightarrow \mathbb{R}$ by

$$\|f\| = \max \left\{ \sum_{x \in \mathbb{Z}^d} |f(x)|, \sup_{x \in \mathbb{Z}^d} |f(x)| \cdot |x|^d \right\} \in [0, \infty],$$

where $|x|$ denotes, say, the ℓ^2 norm in \mathbb{Z}^d . Our norm is a Banach algebra norm with respect to convolution, up to a constant. Indeed, let f and g satisfy that $\|f\|, \|g\| < \infty$. Then

$$\sum_{x \in \mathbb{Z}^d} |(f * g)(x)| \leq \left(\sum_x |f(x)| \right) \left(\sum_x |g(x)| \right) \leq \|f\| \cdot \|g\|, \quad (25)$$

and for every $x \in \mathbb{Z}^d$,

$$|(f * g)(x)| \leq \sum_{y \in \mathbb{Z}^d} |f(y)| |g(x-y)| = \sum_{|y| > |x-y|} |f(y)| |g(x-y)| + \sum_{|y| \leq |x-y|} |f(y)| |g(x-y)|.$$

For the first term, whenever $|y| > |x-y|$, we have $|y| > |x|/2$ and hence

$$\sum_{|y| > |x-y|} |f(y)| |g(x-y)| \leq \sup_{|y| > \frac{1}{2}|x|} |f(y)| \cdot \sum_{|y| > |x-y|} |g(x-y)| \leq \left(\frac{1}{2}|x| \right)^{-d} \|f\| \cdot \|g\|.$$

A similar estimate holds for the other term, now using that $|x-y| \geq |x|/2$ when $|y| \leq |x-y|$, and we get

$$|(f * g)(x)| \leq 2^{d+1} |x|^{-d} \|f\| \|g\|.$$

With (25) we get

$$\|f * g\| \leq 2^{d+1} \|f\| \cdot \|g\|.$$

In particular $B = \{f: \|f\| < \infty\}$ has a Banach algebra structure. In fact, if we define $\|f\|' = 2^{d+1} \|f\|$ then B equipped with $\|\cdot\|'$ is a Banach algebra (see also [11, Theorem VIII.1.4] for a proof that there exists an equivalent norm satisfying that the norm of the unit element is 1). But let us stick to using the norm $\|\cdot\|$. The only property of

Banach algebras that we will use is that if $\|f - \delta_0\| < 2^{-d-1}$, then f is invertible. This follows because the inverse can be defined by the convergent Neumann series

$$f^{-1} = \sum_{n=0}^{\infty} (\delta_0 - f)^n.$$

In addition we get the norm estimate

$$\|f^{-1} - \delta_0\| \leq \sum_{n=1}^{\infty} \|(\delta_0 - f)^n\| \leq \sum_{n=1}^{\infty} (2^{d+1})^{n-1} \|\delta_0 - f\|^n = \frac{\|\delta_0 - f\|}{1 - 2^{d+1} \|\delta_0 - f\|}. \quad (26)$$

We will not need anything else from the theory of Banach algebras.

The following lemma forms the heart of our analysis:

Lemma 4. Fix $d > 2$. Assume $\rho: \mathbb{Z}^d \rightarrow \mathbb{R}$ satisfies

- (i) ρ is symmetric to coordinate permutations and flipping,
- (ii) $\sum_x \rho(x) = 0$,
- (iii) $|\rho(x)| \leq |x|^{-d-4}$.

Then $\|\rho * G^{\text{rw}}\| \leq C$.

Proof. By [17], the random walk Green's function has an expansion of the form

$$G^{\text{rw}}(x) = a|x|^{2-d} + b|x|^{-d} + O(|x|^{-d-2}) \quad (27)$$

(such an expansion is sometimes called an ‘‘Edgeworth expansion’’). Therefore, using (iii) and (27) (in fact here the simpler (2) is enough), the sum defining $(\rho * G^{\text{rw}})(x)$ converges absolutely for every $x \in \mathbb{Z}^d$ so this function is well defined. Write

$$(\rho * G^{\text{rw}})(x) = \sum_y \rho(y)G^{\text{rw}}(x-y) = \sum_{|y| < \frac{1}{2}|x|} \rho(y)G^{\text{rw}}(x-y) + \sum_{|y| \geq \frac{1}{2}|x|} \rho(y)G^{\text{rw}}(x-y) = I + II.$$

We start with I and write it as $I = I_1 + I_2 + I_3$, I_j from the three parts on the RHS of (27). For I_1 we Taylor expand $|x-y|^{2-d}$ around x to order 3 and get

$$\begin{aligned} \frac{1}{a}I_1 &= \sum_{|y| < \frac{1}{2}|x|} \rho(y)|x-y|^{2-d} = \sum_{|y| < \frac{1}{2}|x|} \rho(y) \left[|x|^{2-d} + \sum_{i=1}^d y_i \cdot (2-d)x_i |x|^{-d} \right. \\ &\quad \left. + \sum_{i,j=1}^d y_i y_j (-(2-d)d \cdot x_i x_j |x|^{-d-2} + \delta_{ij}(2-d)|x|^{-d}) + O(|y|^3|x|^{-d-1}) \right]. \end{aligned}$$

We now bound these terms. For the first we write

$$\left| \sum_{|y| < \frac{1}{2}|x|} \rho(y)|x|^{2-d} \right| = |x|^{2-d} \left| \sum_{|y| < \frac{1}{2}|x|} \rho(y) \right| \leq |x|^{2-d} \sum_{|y| \geq \frac{1}{2}|x|} |y|^{-d-4} \leq C|x|^{-d-2},$$

where in the equality we have used that $\sum_x \rho(x) = 0$. For the second, we use the symmetry of ρ to flipping of y_i to conclude that

$$\sum_{|y| < \frac{1}{2}|x|} \rho(y)y_i = 0$$

and similarly for the off-diagonal second-order terms, i.e., for $\sum_{|y| < \frac{1}{2}|x|} \rho(y) y_i y_j$ for $i \neq j$. The on-diagonal terms are equal to

$$\sum_{i=1}^d \sum_{|y| < \frac{1}{2}|x|} \rho(y) y_i^2 ((2-d)|x|^{-d} - (2-d)d \cdot x_i^2 |x|^{-d-2}),$$

and the symmetry of ρ to coordinate permutations shows that $\sum_{|y| < \frac{1}{2}|x|} \rho(y) y_i^2$ does not depend on i . We take it out of the sum and see that

$$\sum_{i=1}^d ((2-d)|x|^{-d} - (2-d)d \cdot x_i^2 |x|^{-d-2}) = 0. \quad (28)$$

Finally, the third order terms are bounded by

$$\sum_{|y| < \frac{1}{2}|x|} |\rho(y)| |y|^3 |x|^{-d-1} \leq |x|^{-d-1} \sum_{|y| < \frac{1}{2}|x|} |y|^{-d-4} |y|^3 \leq C|x|^{-d-1}.$$

Putting all these estimates together gives that

$$|I_1| \leq C|x|^{-d-1}.$$

The estimates of I_2 , I_3 and II are much simpler. To estimate $|I_2|$, we Taylor expand $|x-y|^{-d}$ to first order, i.e. $|x-y|^{-d} = |x|^{-d} + O(|y||x|^{-d-1})$. A similar argument shows that $|I_2| \leq C|x|^{-d-1}$. (We don't need the harmonicity of $|x|^{2-d}$ which is the true reason for the cancellation in (28) above.) For I_3 , we bound, again using that $|x-y| > \frac{1}{2}|x|$ when $|y| < \frac{1}{2}|x|$,

$$|I_3| \leq C \sum_{|y| < \frac{1}{2}|x|} |\rho(y)| |x-y|^{-d-2} \leq C|x|^{-d-2} \sum_{y \in \mathbb{Z}^d} |\rho(x)| \leq C|x|^{-d-2}.$$

For II , we split $II = II_1 + II_2$, depending on whether $|x-y| \leq |x|$ or $|x-y| > |x|$ and use $|y| \geq \frac{1}{2}|x|$, $|\rho(y)| \leq |y|^{-d-4}$ together with $G^{\text{rw}}(x) \leq C|x|^{2-d}$ to bound

$$|II_1| \leq 2^{d+4} |x|^{-d-4} \sum_{y: |x-y| \leq |x|} G^{\text{rw}}(x-y) \leq C|x|^{-d-2},$$

while for $|x-y| > |x|$,

$$|II_2| \leq C|x|^{2-d} \sum_{|y| \geq \frac{1}{2}|x|} |\rho(y)| \leq C|x|^{2-d} |x|^{-4} = C|x|^{-d-2}.$$

We conclude that $|(\rho * G^{\text{rw}})(x)| \leq C|x|^{-d-1}$. This proves the lemma. \square

Remark. It seems as if Lemma 4 makes a stringent requirement on the types of random walks for which the argument can be applied, as Edgeworth expansions are not easy to get. For example, if one wishes to apply the argument for the Cayley graph of, say, the Heisenberg group, then the natural analog of an Edgeworth expansion is not known. We have a more roundabout proof of Lemma 4 that only uses the local central limit theorem. This argument will be presented elsewhere.

In the following lemma, we extend Lemma 4 to the subcritical SRW Green's function G_μ^{rw} , i.e., $G_\mu^{\text{rw}}(x) = \sum_{n \geq 0} (2d\mu)^n p_n(x)$. Note that $G_\mu^{\text{rw}} * \Delta_\mu^{\text{rw}} = \delta_0$.

Lemma 5. Let $d > 2$. Let ρ be as in Lemma 4 and let $\mu \in [-\frac{1}{4d}, \frac{1}{2d}]$. Then

$$\|\rho * G_\mu^{\text{rw}}\| \leq C,$$

where C does not depend on μ .

Proof. Write

$$\rho * G_\mu^{\text{rw}} = \rho * G^{\text{rw}} * \Delta^{\text{rw}} * G_\mu^{\text{rw}}$$

(recall that $G^{\text{rw}} = G_{1/(2d)}^{\text{rw}}$ and $\Delta^{\text{rw}} = \Delta_{1/(2d)}^{\text{rw}}$). Notice that we do not need to put any parenthesis in this expression as associativity follows from the fact that all sums converge absolutely, which can be easily seen from the upper bounds for the various terms. Since we already know that $\|\rho * G^{\text{rw}}\| \leq C$ by Lemma 4, we need only bound $\|G_\mu^{\text{rw}} * \Delta^{\text{rw}}\|$ (note again that this is not Δ_μ^{rw} but rather $\Delta_{1/(2d)}^{\text{rw}}$). Noting that $p_n * p_m = p_{n+m}$, we get

$$G_\mu^{\text{rw}} * \Delta^{\text{rw}} = \left(\sum_{n=0}^{\infty} (2d\mu)^n p_n \right) * (\delta_0 - p_1) = \delta_0 - (1 - 2d\mu) \sum_{n=1}^{\infty} (2d\mu)^{n-1} p_n.$$

For $\mu = 1/(2d)$, this is identically equal to δ_0 , and there is nothing to prove. Thus, we can assume that $\mu \in [-\frac{1}{4d}, \frac{1}{2d})$. Now, since $p_n(x) \leq Cn^{-d/2}e^{-c|x|^2/n}$ we get that $\|p_n\| \leq C$. Hence,

$$\|G_\mu^{\text{rw}} * \Delta^{\text{rw}}\| \leq 1 + (1 - 2d\mu) \sum_{n=1}^{\infty} (2d\mu)^{n-1} \|p_n\| \leq C,$$

and the lemma is proved. \square

Proof of Lemma 2. Recall that the input of the lemma is a function Δ satisfying conditions (i)–(iii) of Lemma 1. These conditions are quite close to the conditions on ρ in Lemmas 4 and 5, only a linear map is required to pass from one to the other. We define μ by

$$\mu = \frac{1}{2d} \left(1 - \sum_{x \in \mathbb{Z}^d} \Delta(x) \right),$$

which implies that

$$\sum_{x \in \mathbb{Z}^d} (\Delta - \Delta_\mu^{\text{rw}})(x) = 0. \quad (29)$$

(This choice is closely related to the choice of constants λ, μ in [9, (2.29)].) To use Lemma 5 we need to justify $\mu \in [-\frac{1}{4d}, \frac{1}{2d}]$. The upper bound $\mu \leq \frac{1}{2d}$ is automatic since $\sum_x \Delta(x) \geq 0$. For the lower bound $\mu \geq -\frac{1}{4d}$, we need to show that $\sum_x \Delta(x) \leq \frac{3}{2}$. This follows for small enough β by our assumption that Δ satisfies (iii) of Lemma 1 and $\sum_x \Delta_{\lambda'}^{\text{rw}}(x) \leq 1$.

Next we note that

$$|\Delta(x) - \Delta_\mu^{\text{rw}}(x)| \leq C_2\beta|x|^{-d-4}. \quad (30)$$

Indeed, at every x that is not a neighbour of 0 this is an immediate corollary from our condition (iii) of Lemma 1. For the neighbours, we note that

$$|\lambda' - \mu| = \frac{1}{2d} \left| \sum_{x \in \mathbb{Z}^d} (\Delta_{\lambda'}^{\text{rw}} - \Delta_\mu^{\text{rw}})(x) \right| \stackrel{(29)}{=} \frac{1}{2d} \left| \sum_{x \in \mathbb{Z}^d} (\Delta_{\lambda'}^{\text{rw}} - \Delta)(x) \right| \leq \frac{C\beta}{2d}, \quad (31)$$

where the last inequality is again from condition (iii) of Lemma 1. Thus, for x a neighbour of the origin, we conclude that

$$\Delta(x) - \Delta_\mu^{\text{rw}}(x) = \mu - \lambda' + \Delta(x) - \Delta_{\lambda'}^{\text{rw}}(x) = O(\beta),$$

by (31) and condition (iii) of Lemma 1. This shows (30).

We next define

$$\rho = \frac{1}{C_2\beta}(\Delta - \Delta_\mu^{\text{rw}}),$$

with C_2 being the constant from (30). This ρ has the properties (i)–(iii) of Lemma 4, so that

$$\|\rho * G_\mu^{\text{rw}}\| \leq C.$$

In turn, this implies

$$\|(\Delta - \Delta_\mu^{\text{rw}}) * G_\mu^{\text{rw}}\| \leq C\beta.$$

But this is exactly $\Delta * G_\mu^{\text{rw}} - \delta_0$. This means that $\Delta * G_\mu^{\text{rw}}$ is invertible if β is sufficiently small (recall (26)), and further that we have $(\Delta * G_\mu^{\text{rw}})^{-1} = \delta_0 + E$ with $\|E\| \leq C\beta$. Our required function is now

$$G = (\Delta * G_\mu^{\text{rw}})^{-1} * G_\mu^{\text{rw}},$$

which is clearly an inverse for Δ . To see that $G(x) \leq 2G^{\text{rw}}(x)$ write

$$G = G_\mu^{\text{rw}} + E * G_\mu^{\text{rw}}. \quad (32)$$

Since $G_\mu^{\text{rw}}(x) \leq G^{\text{rw}}(x)$, because $\mu \leq 1/(2d)$, we need only estimate $E * G_\mu^{\text{rw}}$. We write

$$(E * G_\mu^{\text{rw}})(x) = \sum_y G_\mu^{\text{rw}}(y)E(x-y) = \sum_{|y| < \frac{1}{2}|x|} G_\mu^{\text{rw}}(y)E(x-y) + \sum_{|y| \geq \frac{1}{2}|x|} G_\mu^{\text{rw}}(y)E(x-y) = I + II. \quad (33)$$

For I , we use that $|y| < \frac{1}{2}|x|$ implies that $|x-y| \geq \frac{1}{2}|x|$ so $|E(x-y)| \leq C\beta|x|^{-d}$ and hence

$$|I| \leq C\beta|x|^{-d} \sum_{|y| < \frac{1}{2}|x|} |G_\mu^{\text{rw}}(y)| \leq C\beta|x|^{-d} \sum_{|y| < \frac{1}{2}|x|} |y|^{2-d} \leq C\beta|x|^{2-d}. \quad (34)$$

For II , we have

$$|II| \leq \left(\max_{|y| \geq \frac{1}{2}|x|} |G_\mu^{\text{rw}}(y)| \right) \cdot \sum_y |E(x-y)| \leq C \left(\frac{1}{2}|x| \right)^{2-d} \cdot C\beta = C\beta|x|^{2-d}. \quad (35)$$

We get that $|(E * G_\mu^{\text{rw}})| \leq C\beta|x|^{2-d}$, which means that for β sufficiently small, it is less than G^{rw} . This shows that $G(x) \leq 2G^{\text{rw}}(x)$, and thus completes the proof of Lemma 2. \square

Remarks

(1) Examining (32)–(35) in the proof of the last lemma shows that in fact we got that

$$G_\lambda^{\text{saw}}(x) = G_\mu^{\text{rw}}(x) + O(\beta|x|^{2-d}).$$

Together with (27), this would prove (3), if only we could show that $\mu = \frac{1}{2d}$ for λ_c , the critical λ . This is a classical fact, let us sketch its proof for the convenience of the reader. Since $\mu = \frac{1}{2d}(1 - \sum_x \Delta(x))$, it is equivalent to showing

that $\chi(\lambda) \rightarrow \infty$ as $\lambda \nearrow \lambda_c$ where $\chi(\lambda) = \sum_x G_\lambda^{\text{saw}}(x)$ (recall (14)). Let $c_n = \sum_{\text{len}(\gamma)=n} W^\beta(\gamma)$ so $\chi(\lambda) = \sum_n \lambda^n c_n$. Then, c_n is submultiplicative i.e. $c_n \leq c_m c_{n-m}$, so $c_n \geq \lambda_c^n$. Summing gives $\chi(\lambda) \geq 1/(\lambda_c - \lambda)$. (The reader who finds this sketch too dense may see more details in, say, [16, Theorem 2.3].)

(2) The result of the theorem is known as an “infrared bound.” It implies the finiteness of the so-called *bubble diagram*, which in turn implies various critical exponents. See again [16, Theorem 2.3].

(3) Let us remark on the exponent $-d-4$ appearing in the inequality $|\Delta^{\text{rw}} - \Delta^{\text{saw}}| \leq C\beta|x|^{-d-4}$ of Lemma 1. We in fact prove a stronger bound with exponent $-3(2-d)$, recall (12). Most of the proof of Lemma 2 actually needs less, $|x|^{-d-2-\varepsilon}$ would have been enough. The only place where the stronger estimate $|x|^{-d-4}$ is used is in Lemma 5, in order to justify the associativity of the convolution in the expression $\rho * G^{\text{rw}} * \Delta^{\text{rw}} * G_\mu^{\text{rw}}$. There are certainly ways to justify associativity at that point under the weaker assumption $|\rho(x)| \leq |x|^{-d-2-\varepsilon}$, but an additional argument would be needed.

Acknowledgements

This work was performed in part during a visit of RvdH to the Weizmann Institute, and when the authors met in Eurandom, Oberwolfach and the Institut Henri Poincaré. We thank these institutions for their hospitality.

References

- [1] L. Avena, E. Bolthausen and C. Ritzmann. A local CLT for convolution equations with an application to weakly self-avoiding random walks. *Ann. Probab.* **44** (1) (2016) 206–234. [MR3456336](#)
- [2] E. Bolthausen and C. Ritzmann. A central limit theorem for convolution equations and weakly self-avoiding walks. Unpublished manuscript, 2001. Available at [arXiv:math/0103218](#).
- [3] C. Borgs, J. T. Chayes, R. van der Hofstad, G. Slade and J. Spencer. Random subgraphs of finite graphs. II. The lace expansion and the triangle condition. *Ann. Probab.* **33** (5) (2005) 1886–1944. [MR2165583](#)
- [4] D. Brydges and T. Spencer. Self-avoiding walk in 5 or more dimensions. *Comm. Math. Phys.* **97** (1–2) (1985) 125–148. [MR0782962](#)
- [5] L.-C. Chen and A. Sakai. Critical two-point functions for long-range statistical-mechanical models in high dimensions. *Ann. Probab.* **43** (2) (2015) 639–681. [MR3306002](#)
- [6] T. Hara. Decay of correlations in nearest-neighbor self-avoiding walk, percolation, lattice trees and animals. *Ann. Probab.* **36** (2) (2008) 530–593. [MR2393990](#)
- [7] T. Hara and G. Slade. Mean-field critical behaviour for percolation in high dimensions. *Comm. Math. Phys.* **128** (2) (1990) 333–391. [MR1043524](#)
- [8] T. Hara and G. Slade. On the upper critical dimension of lattice trees and lattice animals. *J. Stat. Phys.* **59** (5–6) (1990) 1469–1510. [MR1063208](#)
- [9] T. Hara, R. van der Hofstad and G. Slade. Critical two-point functions and the lace expansion for spread-out high-dimensional percolation and related models. *Ann. Probab.* **31** (2) (2003) 349–408. [MR1959796](#)
- [10] R. van der Hofstad, F. den Hollander and G. Slade. A new inductive approach to the lace expansion for self-avoiding walks. *Probab. Theory Related Fields* **111** (2) (1990) 253–286. [MR1633582](#)
- [11] Y. Katznelson. *An Introduction to Harmonic Analysis*, 3rd edition. *Cambridge Mathematical Library*. Cambridge University Press, Cambridge, 2004. [MR2039503](#)
- [12] A. Sakai. Mean-field critical behavior for the contact process. *J. Stat. Phys.* **104** (1–2) (2001) 111–143. [MR1851386](#)
- [13] A. Sakai. Lace expansion for the Ising model. *Comm. Math. Phys.* **272** (2) (2007) 283–344. [MR2300246](#)
- [14] A. Sakai. Application of the lace expansion to the φ^4 model. *Comm. Math. Phys.* **336** (2) (2015) 619–648. [MR3322382](#)
- [15] G. Slade. The diffusion of self-avoiding random walk in high dimensions. *Comm. Math. Phys.* **110** (4) (1987) 661–683. [MR0895223](#)
- [16] G. Slade. The lace expansion and its applications. In *Lectures from the 34th Summer School on Probability Theory Held in Saint-Flour, July 6–24*. J. Picard (Ed). *Lecture Notes in Mathematics* **1879**. Springer-Verlag, Berlin, 2004. [MR2239599](#)
- [17] K. Uchiyama. Green’s functions for random walks on \mathbb{Z}^N . *Proc. Lond. Math. Soc.* (3) **77** (1) (1998) 215–240. [MR1625467](#)